



Impact of dust addition on the microbial food web under present and future conditions of pH and temperature

Julie Dinasquet, Estelle Bigeard, Frédéric Gazeau, Farooq Azam, Cécile Guieu, Emilio Marañón, Céline Ridame, France van Wambeke, Ingrid Obernosterer, Anne-Claire Baudoux

► To cite this version:

Julie Dinasquet, Estelle Bigeard, Frédéric Gazeau, Farooq Azam, Cécile Guieu, et al.. Impact of dust addition on the microbial food web under present and future conditions of pH and temperature. 2021. insu-03636650v1

HAL Id: insu-03636650

<https://hal.sorbonne-universite.fr/insu-03636650v1>

Preprint submitted on 15 Jun 2021 (v1), last revised 12 Apr 2022 (v2)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



1 **Impact of dust addition on the microbial food web under present and future**
2 **conditions of pH and temperature**

3

4 Julie Dinasquet^{1,2*}, Estelle Bigeard³, Frédéric Gazeau⁴, Farooq Azam¹, Cécile Guieu⁴, Emilio
5 Marañón⁵, Céline Ridame⁶, France Van Wambeke⁷, Ingrid Obernosterer² and Anne-Claire
6 Baudoux³

7 ¹ Marine Biology Research Division, Scripps Institution of Oceanography, UCSD, USA

8 ² Sorbonne Université, CNRS, Laboratoire d'Océanographie Microbienne, LOMIC, France

9 ³ Sorbonne Université, CNRS, Station Biologique de Roscoff, UMR 7144 Adaptation et
10 Diversité en Milieu Marin, France

11 ⁴ Sorbonne Université, CNRS, Laboratoire d'Océanographie de Villefranche, LOV, 06230
12 Villefranche-sur-Mer, France

13 ⁵ Department of Ecology and Animal Biology, Universidade de Vigo, Spain

14 ⁶ CNRS-INSU/IRD/MNH/UPMC, LOCEAN: Laboratoire d'Océanographie et du Climat:
15 Expérimentation et Approches Numériques, UMR 7159

16 ⁷ Aix-Marseille Université, CNRS/INSU, Université de Toulon, IRD, Mediterranean Institute of
17 Oceanography, UM110, France

18 *Corresponding: jdinasquet@ucsd.edu, present address: Center for Aerosol Impact on Chemistry
19 of the Environment (CAICE), Scripps Institution of Oceanography, UCSD, USA

20 **Keywords:** bacteria, microeukaryotes, virus, community composition, top-down



21 Abstract

22 In the oligotrophic waters of the Mediterranean Sea, during the stratification period, the
23 microbial loop relies on pulsed inputs of nutrients through atmospheric deposition of aerosols
24 from both natural (Saharan dust) and anthropogenic origins. While the influence of dust
25 deposition on microbial processes and community composition is still not fully constrained, the
26 extent to which future environmental conditions will affect dust inputs and the microbial
27 response is not known. The impact of atmospheric wet dust deposition was studied both under
28 present and future (warming and acidification) environmental conditions through experiments in
29 300 L climate reactors. Three dust addition experiments were performed with surface seawater
30 collected from the Tyrrhenian Sea, Ionian Sea and Algerian basin in the Western Mediterranean
31 Sea during the PEACETIME cruise in May-June 2017. Top-down controls on bacteria, viral
32 processes and community, as well as microbial community structure (16S and 18S rDNA
33 amplicon sequencing) were followed over the 3-4 days experiments. Different microbial and
34 viral responses to dust were observed rapidly after addition and were most of the time higher
35 when combined to future environmental conditions. The input of nutrients and trace metals
36 changed the microbial ecosystem from bottom-up limited to a top-down controlled bacterial
37 community, likely from grazing and induced lysogeny. The composition of mixotrophic
38 microeukaryotes and phototrophic prokaryotes was also altered. Overall, these results suggest
39 that the effect of dust deposition on the microbial loop is dependent on the initial microbial
40 assemblage and metabolic state of the tested water, and that predicted warming, and acidification
41 will intensify these responses, affecting food web processes and biogeochemical cycles.



42 1. Introduction

43 Input of essential nutrients and trace metals through aerosol deposition is crucial to the ocean
44 surface water biogeochemistry and productivity (at the global scale: *e.g.*, Mahowald et al., 2017;
45 in the Mediterranean Sea: *e.g.*, Guieu and Ridame, 2020) with episodic fertilization events
46 driving microbial processes in oligotrophic regions such as the Pacific Ocean, the Southern
47 Ocean and the Mediterranean Sea.

48 The summer Mediterranean food web is characterized by low primary production (PP) and
49 heterotrophic prokaryotic production (more classically abbreviated as BP for bacterial
50 production) constrained by nutrient availability further limiting dissolved organic matter (DOM)
51 utilization and export, resulting in DOM accumulation. Therefore, inputs of bioavailable
52 nutrients through deposition of atmospheric particles are essential to this microbial ecosystem.
53 Indeed, these nutrient pulses have been shown to support microbial processes but the degree to
54 which the microbial food web is affected might be dependent on the degree of oligotrophy of the
55 water (Marín-Beltrán et al., 2019; Maraňon et al., 2010).

56 In the Mediterranean Sea, dust deposition stimulates PP and N₂ fixation (Guieu et al., 2014;
57 Ridame et al., 2011) but also BP, bacterial respiration, virus production, grazing activities, and
58 can alter the composition of the microbial community (*e.g.*, Pulido-Villena et al., 2014; Tsiola et
59 al., 2017; Guo et al., 2016; Pitta et al., 2017; Marín-Beltrán et al., 2019). Overall, in such
60 oligotrophic system, dust deposition appears to predominantly promote heterotrophic activity
61 which will increase respiration rates and CO₂ release.

62 Anthropogenic CO₂ emissions are projected to induce an increase in seawater temperature
63 and an accumulation of CO₂ in the ocean, leading to its acidification and an alteration of ocean
64 carbonate chemistry (IPCC, 2014). In response to ocean warming and increased stratification,



low nutrient low chlorophyll (LNLC) regions such as the Mediterranean Sea, are projected to expand in the future (Durrieu de Madron et al., 2011). Moreover, dust deposition is also expected to increase due to desertification (Moulin and Chiapello, 2006). Hence, in the future ocean, the microbial food web might become even more dependent on atmospheric deposition of nutrients. Expected increased temperature and acidification might have complex effects on the microbial loop by modifying microbial and viral and community (*e.g.*, Highfield et al., 2017; Krause et al., 2012; Hu et al., 2021; Allen et al., 2020; Malits et al., 2021). While increasing temperature in combination with nutrient input might enhance heterotrophic bacterial growth (Degerman et al., 2012; Morán et al., 2020) more than PP (Marañón et al., 2018), future environmental conditions could push even further this microbial community towards heterotrophy. But so far, the role of dust on the microbial food web in future climate scenarios is unknown.

Here, we studied the response of Mediterranean microbial and viral communities (*i.e.*, viral strategies, microbial growth, and controls, as well as community composition) to simulated wet Saharan dust deposition during onboard minicosm experiments conducted in three different basins of the Western and Central Mediterranean Sea under present and future projected conditions of temperature and pH. To our knowledge, this is the first study assessing the effect of atmospheric deposition on the microbial food web under future environmental conditions.



82 2. Material & Method

83 2.1 Experimental set-up

84 During the ProcEss studies at the Air-sEa Interface after dust deposition in the
 85 Mediterranean sea project cruise (PEACETIME), onboard the R/V “Pourquoi Pas ?” in
 86 May/June 2017, three experiments were conducted in 300 L climate reactors (minicosms) filled
 87 with surface seawater collected at three different stations (Table 1), in the Tyrrhenian Sea (TYR),
 88 Ionian Sea (ION) and in the Algerian basin (FAST). The experimental set-up is described in
 89 details in Gazeau et al. (2020). Briefly, the experiments were conducted for 3 days (TYR and
 90 ION) and 4 days (FAST) in trace metal free conditions, under light, temperature and pH-
 91 controlled conditions following ambient or future projected conditions of temperature and pH.
 92 For each experiment, the biogeochemical evolution of the water, after dust deposition, under
 93 present and future environmental conditions was followed in three duplicate treatments: i)
 94 CONTROL (C1, C2) with no dust addition and under present pH and temperature conditions, ii)
 95 DUST (D1, D2) with dust addition under present environmental conditions and iii)
 96 GREENHOUSE (G1, G2) with dust addition under projected temperature and pH for 2100
 97 (IPCC, 2014; ca. +3 °C and -0.3 pH units). The same dust analog was used as during the DUNE
 98 2009 experiments described in Desboeufs et al. (2014) and the same dust wet flux of 10 g m⁻²
 99 was simulated. Such deposition event represents a high but realistic scenario, as several studies
 100 reported even higher short deposition events in this area of the Mediterranean Sea (Ternon et al.,
 101 2010; Bonnet and Guieu, 2006; Loÿe-Pilot and Martin, 1996). After mixing the dust analog (3.6
 102 g) in 2 L of ultrahigh-purity water, this solution was sprayed at the surface of the dust amended
 103 treatments (D1, D2 and G1, G2; Gazeau et al., 2020).



104 Samples were taken at t-12h (while filling the tanks), t0 (just before dust addition), t1h,
 105 t6h, t12h, t24h, t48h, t72h and t96h (after dust addition, and t96h only for FAST).

106 2.2. Growth rates, mortality, and top down controls

107 BP was estimated at all sampling points from rates of ^3H -Leucine incorporation
 108 (Kirchman et al., 1985; Smith and Azam, 1992) as described in Gazeau et al. (2021). Briefly,
 109 triplicate 1.5 mL samples and one blank were incubated in the dark for 1-2 h in two temperature-
 110 controlled incubators maintained respectively at ambient temperature for C1, C2, D1 and D2 and
 111 at ambient temperature +3 °C for G1 and G2. HB, *Synechococcus*, picoeukaryotes and
 112 heterotrophic nanoflagellates (HNF) abundances were measured by flow cytometry as described
 113 in Gazeau et al. (2020). Bacterial cell specific growth rates were estimated assuming exponential
 114 growth and a carbon to cell ration of 20 fg C cell⁻¹ (Lee and Fuhrman, 1987). Net growth rates
 115 (h⁻¹) were calculated from the exponential phase of growth of BP, abundances of *Synechococcus*
 116 and picoeukaryotes cells, observable from at least three successive sampling points. Mortality
 117 was estimated as the difference between HB present between two successive sampling points and
 118 those produced during that time.

119 2.3. Viral abundance, production and life strategy

120 Virus abundances were determined on glutaraldehyde fixed samples (0.5% final
 121 concentration, Grade II, Sigma Aldrich, St Louis, MO, USA) stored at -80 °C until analysis. Flow
 122 cytometry analysis was performed as described by Brussaard (2004). Briefly, samples were
 123 thawed at 37 °C, diluted in 0.2 µm filtered autoclaved TE buffer (10:1 Tris-EDTA, pH 8) and
 124 stained with SYBR-Green I (0.5×10^{-4} of the commercial stock, Life Technologies, Saint-Aubin,
 125 France) for 10 min at 80 °C. Virus particles were discriminated based on their green fluorescence



126 and SSC during 1 min analyses (Fig. S1). All cytogram analyses were performed with the Flowing
 127 Software freeware (Turku Center of Biotechnology, Finland).

128 Viral production and bacterial losses due to phages were assessed by the virus reduction approach
 129 (Weinbauer et al., 2010) at t0, t24 h and t48h in all six minicosms. Briefly, 3 L of seawater were-
 130 filtered through 1.2-µm-pore-size polycarbonate filter (Whatman©), and heterotrophic
 131 prokaryotes (HB, filtrate) were concentrated by ultrafiltration (0.22 µm pore size, Vivaflow 200©
 132 polyethersulfone, PES) down to a volume of 50 mL. Virus-free water was obtained by filtering 1
 133 L of seawater through a 30 kDa pore-size cartridge (Vivaflow 200©, PES). Six mixtures of HB
 134 concentrate (2 mL) diluted in virus-free water (23 mL) were prepared and distributed into 50 mL
 135 Falcon tubes. Three of the tubes were incubated as controls, while the other three were inoculated
 136 with mitomycin C (Sigma-Aldrich, 1 µg mL⁻¹ final concentration) as inducing agent of the lytic
 137 cycle in lysogenic bacteria. All tubes were incubated in darkness in two temperature-controlled
 138 incubators maintained respectively at ambient temperature for C1, C2, D1 and D2 and at ambient
 139 temperature +3 °C for G1 and G2. Samples for HB and viral abundances were collected every 6 h
 140 for a total incubation period of 18 h.

141 The estimation of virus-mediated mortality of HB was performed according to Weinbauer et al.
 142 (2002) and Winter et al. (2004). Briefly, increase in virus abundance in the control tubes represents
 143 lytic viral production (VPL), and an increase in mitomycin C treatments represents total (VPT),
 144 *i.e.*, lytic plus lysogenic, viral production. The difference between VPT and VPL represents
 145 lysogenic production (VPLG). The frequency of lytically infected cells (FLIC) and the frequency
 146 of lysogenic cells (FLC) were calculated as:

147
$$FLIC = 100 \times VPL / BS \times HB_i \quad (1)$$

148
$$FLC = 100 \times VPLG / BS \times HB_i \quad (2)$$



149 where HB_i is the initial HB abundance in the viral production experiment and BS is a theoretical
 150 burst size of 20 viruses per infected cell (averaged BS in marine oligotrophic waters, Parada et al.,
 151 2006).

152

153 2.4 DNA sampling, sequencing and sequence analysis

154 To study the temporal dynamics of the microbial diversity, water samples (3 L) were
 155 collected in acid-washed containers from each minicosm at t_0 , t_{24h} , and at the end of the
 156 experiments (t_{72h} at TYR and ION and t_{96h} at FAST). Samples were filtered onto 0.2 μm PES
 157 filters (Sterivex©) and stored at $-80^\circ C$ until DNA extraction. Nucleic acids were extracted from
 158 the filters using a phenol-chloroform method and DNA was then purified using filter columns from
 159 NucleoSpin® PlantII kit (Macherey-Nagel©) following a modified protocol. DNA extracts were
 160 quantified and normalized at $5ng\ \mu L^{-1}$ and used as templates for PCR amplification of the V4
 161 region of the 18S rRNA (~380 bp) using the primers TAREuk454FWD1 and TAREukREV3
 162 (Stoeck et al., 2010) and the V4-V5 region of the 16S rRNA (~411 bp) using the primers 515F-Y
 163 (5'-GTGYCAGCMGCCGCGGTAA) and 926R-R (5'-CCGYCAATTYMTTTRAGTTT) (Parada
 164 et al., 2016). Following polymerase chain reactions, DNA amplicons were purified, quantified and
 165 sent to Genotoul (<https://www.genotoul.fr/>, Toulouse, France) for high throughput sequencing
 166 using paired-end 2x250bp Illumina MiSeq. Note that although we used universal primer, Archaea
 167 were mostly not detected and the prokaryotic heterotrophic communities corresponded essentially
 168 to Eubacteria, therefore the taxonomic description referred to the general term 'bacterial
 169 communities'

170 All reads were processed using the Quantitative Insight Into Microbial Ecology 2 pipeline
 171 (QIIME2 v2020.2, Bolyen et al., 2019). Reads were truncated 350bp based on sequencing



172 quality, denoised, merged and chimera-checked using DADA2 (Callahan et al., 2016). A total of
173 714 and 3070 amplicon sequence variants (ASVs) were obtained for 16S and 18S respectively.
174 Taxonomy assignments were made against the database SILVA 132 (Quast et al., 2013) for 16S
175 and PR2 (Guillou et al., 2013) for 18S. All sequences associated with this study have been
176 deposited under the BioProject ID: PRJNA693966.

177 2.5 Statistics

178 Alpha and beta-diversity indices for community composition were estimated after
179 randomized subsampling to 26000 reads for 16S rDNA and 19000 reads for 18S rDNA. Analysis
180 were run in QIIME 2 and in Primer v.6 software package (Clarke and Warwick, 2001).
181 Differences between the samples richness and diversity were assessed using Kruskal-Wallis
182 pairwise test. Beta diversity were run on Bray Curtis dissimilarity. Differences between samples'
183 beta diversity were tested using PERMANOVA (Permutational Multivariate Analysis of
184 Variance) with pairwise test and 999 permutations. The sequences contributing most to the
185 dissimilarity between clusters were identified using SIMPER (similarity percentage). A linear
186 mixed model was performed using the R software (R Core Team, 2020) using the nlme package
187 (Pinheiro et al., 2014) to test if the amended treatments differed from the controls at t24h and
188 t72h or t96h.



189 3. Results

190 3.1. Microbial growth, mortality and top-down controls

191 Significant increases in heterotrophic bacterial cell specific growth rates were observed in
 192 all experiments with dust under D and G (Fig. 1, $p \leq 0.016$ after 24 h and 72 h) relative to C, the
 193 highest growth rates relative to C were observed already 24 h after dust seeding. Bacterial net
 194 growth rates were also higher in D and especially in G relative to C (Table 2). *Synechococcus*
 195 and picoeukaryotes net growth rates showed a similar trend (Table 2). Heterotrophic bacterial
 196 mortality was also higher than in C especially at TYR and in G at ION and FAST (Fig. 1). Over
 197 the course of the three experiments, the slope of the linear regression between bacterial biomass
 198 and bacterial production was below 0.4 in the three treatments suggesting a weak bottom up
 199 control (Fig. 2A; Ducklow, 1992). The slope decreased in D and G relative to C. Overall, the top
 200 down index, as described by Morán et al. (2017), was higher in G (0.92) relative to C and D
 201 (0.80). The relationship between log transformed HNF and bacterial abundance (Fig. 3B), plotted
 202 according to the model in Gasol (1994), showed that HNF were below the MRA (Mean realized
 203 HNF abundance) in all treatments, suggesting a top down control of HNF abundance. HNF and
 204 bacteria were weakly coupled in all treatments. The relationship between total viruses and
 205 bacterial abundance was weaker in D and G relative to C (Fig. S2).

206

207 3.2. Viral dynamics and processes

208 The abundance and production of virus-like particles (VLP) increased following an east
 209 to west gradient (Table 1). Viral strategy (lysogenic vs. lytic replication) was also different
 210 between stations, with a higher frequency of lysogenic cells (FLC) at TYR and ION (23 and



211 19%, respectively Table 1) and a higher frequency of lytically infected cells (FLIC) at FAST
 212 (43%, Table 1).

213 During TYR and ION experiments, the relative contribution of VLP populations was similar
 214 and stable over time with Low DNA viruses representing over 80% of the community (Figs. 3
 215 and S3). The Low DNA VLP abundance was however slightly higher in D and G relative to C
 216 after 24 h at TYR and significantly higher at ION after 48h ($p = 0.037$; Fig. S3). In contrast to
 217 the other two stations, at FAST, Giruses were also present and increased in all treatments but
 218 especially in G where they made up to 9% of the viral community at the end of the experiment
 219 (Figs. 3 and S3). The abundance of high DNA viruses at FAST also increased independent of
 220 treatments and accounted for 16 – 18% of the community at the end of the experiment (Figs. 3
 221 and S3).

222 The sampling strategy for production and life strategies of HB viruses allowed to
 223 discriminate independently the effect of i) greenhouse conditions (sampling at T0 before dust
 224 addition), ii) dust addition (sampling at T24) and the combined effects of dust addition and
 225 greenhouse. Lytic viral production (VPL) increased significantly at T0 in G at TYR and ION
 226 compared to C ($p \leq 0.036$). The addition of dust induced higher VPL in D at TYR compared to C
 227 (Fig.1). No significant impact of dust on VPL was observed in G compared to D after 24h for
 228 any of the experiments. Changes in viral infection strategy were observed with G conditions at
 229 T0 where, FLC decreased relative to the non-G treatments at TYR and ION, and especially at
 230 FAST ($p = 0.047$). FLIC increased slightly in G at TYR and ION already at T0. Dust addition
 231 had no detectable significant effect on this parameter for any experiments. Looking at the
 232 relative share between lytic and lysogenic infection, dust addition favored lytic infection at TYR
 233 (no lysogenic bacteria were observed after 24h) but the contribution of both infection strategies



remained unchanged compared to C at ION and FAST. Greenhouse conditions also favored replication through lytic cycle already at T0 for all three experiments and this trend was not impacted by dust addition.

3.3. Microbial community composition

Microbial community structure, bacteria and micro-eukaryotes from 16S rDNA and 18S rDNA sequencing respectively, responded to dust addition in all three experiments relative to C (Figs. 4 and 5). After quality controls, reads were assigned to 714 and 1443 ASVs for 16S and 18S respectively.

3.3.1. Bacterial community composition

The initial community composition (t-12h) was significantly different at the three stations (PERMANOVA; $p = 0.001$, Fig. S4a, S5). Rapid and significant changes in the bacterial community composition were observed already 24 h after dust addition (Fig. 4). Despite the initial different communities, the three stations appeared to converge towards a closer community composition in response to dust addition (Fig. S5). At TYR, communities in D and G significantly changed 24 h after dust addition (PERMANOVA; $p = 0.001$). This cluster presented no significant differences between treatments (D and G) or time (24 and 72 h). The differences between C and D/G were attributed to a relative increase of ASVs related to different *Alteromonas* sp., OM60 and *Pseudophaeobacter* sp. and *Erythrobacter* sp.; contribution of ASVs related to SAR11 and Verrucomicrobia and *Synechococcus* decreased (Table S1a). At ION, the bacterial community composition significantly changed 24 h after dust addition (PERMANOVA; $p = 0.001$) and was significantly different between D and G (PERMANOVA; $p = 0.032$). As observed at TYR, no further change occurred between 24 h and the end of the



experiment (72 h; Fig. 4). The difference between the controls and dust amended minicosms were assigned to an increase of ASVs related to different *Alteromonas* sp., *Erythrobacter* sp., *Dokdonia* sp. and OM60, and a decrease of ASVs related to SAR11, *Synechococcus*, Verrucomicrobia, Rhodospirillales and some Flavobacteria (Table S1b). Several ASVs related to *Alteromonas* sp., *Synechococcus* sp. and *Erythrobacter* sp. were further enriched in G compared D while *Dokdonia* sp. was mainly present in D. At FAST, the bacterial community after 24 h only significantly changed in G (PERMANOVA; $p = 0.011$; Fig. 4). However, after 96 h, the community in D and G were similar and appeared to transition back to the initial state at 96 h (PERMANOVA; $p = 0.077$). The higher relative abundance in *Erythrobacter* sp., *Synechococcus* sp., different ASVs related to *Alteromonas* sp. and Flavobacteria appeared to contribute mainly to the difference between C and D/G (Table S1) while ASVs related to SAR11, Verrucomicrobia, *Celeribacter* sp. *Thalassobius* sp. and Rhodospirillales were mainly present in C (Table S1c).

3.3.2 Nano- and micro-eukaryotes community composition

The diversity of initial community was large (Fig. S5) and significantly different at the three stations (PERMANOVA; $p = 0.001$; Fig. S4b). At TYR, the nano- and micro-eukaryotes community responded rapidly (24 h) to dust addition (PERMANOVA; $p = 0.003$). This initial high diversity disappeared after 72 h, with similar communities in all minicosms (Fig. S5). They were significantly different from initial and t24h communities ($p = 0.002$ and 0.03 respectively; Fig 5) in D/G. The variations at t24h were attributed to changes in the dinoflagellate communities in particular to an increase in ASVs related to *Heterocapsa rotundata*, Gymnodiniales and Gonyaulacales as well as to an increase in Chlorophyta (Table S2a). At ION, no significant changes were observed between C and D/G after 24 h. However, after 72 h, the



279 communities were significantly different in D ($p = 0.018$) and G ($p = 0.05$) compared to the
280 communities at t24h in these treatments (Table S2B). In D, diversity was significantly higher at
281 t72h compared to t24h and to C at the same sampling time ($p = 0.036$). In contrast, diversity in G
282 at t72h was lower than at t24h and lower to the one observed in C at the same sampling time ($p =$
283 0.066 ; Fig S6). These differences were mainly attributed to changes in ASVs related to
284 dinoflagellates and to the increase at t72h of *Emiliana huxleyi* and Chlorophyta in D and G,
285 respectively (Table S2b). At FAST, significant differences were observed between the controls
286 and initial communities compared to the dust amended (D and G) treatments at t24h ($p = 0.036$).
287 No major differences were observed between D/G at t24h and t96h ($p = 0.06$). The differences
288 were mainly attributed to changes in dinoflagellates ASVs and to an increase in Acantharea and
289 *Emiliana huxleyi* in D and G treatments at t96h (Table S2c).



290 **4. Discussion**

291 Pulsed inputs of essential nutrients and trace metals through aerosol deposition are crucial to
292 surface microbial communities in LNLC regions such as the Mediterranean Sea (reviewed in
293 Guieu and Ridame, 2020). Here we assessed the impact of dust deposition on the late spring
294 microbial loop under present and future environmental conditions on the surface water of three
295 different Mediterranean basins (Tyrrhenian, TYR; Ionian, ION; and Algerian, FAST). The initial
296 conditions at the three sampled stations for the onboard experiments are described in more
297 details in Gazeau et al. (2020). Briefly, very low levels of dissolved inorganic nutrients were
298 measured at all three stations, highlighting the oligotrophic status of the waters, typical of the
299 stratified conditions observed in the Mediterranean Sea in late spring/early summer (*e.g.*, Bosc et
300 al., 2004; D'Ortenzio et al., 2005). Despite similar total chl. *a* concentrations at the three stations
301 (Gazeau et al., 2020), PP was higher at FAST (Table 1, Gazeau et al., 2021; Marañón et al.,
302 2021). The initial microbial communities differed substantially between the three stations as
303 shown by pigments (Gazeau et al., 2020), 18S and 16S rDNA sequencing (this study). DOC
304 concentrations were slightly higher at TYR where PP was the lowest (Gazeau et al., 2021). HB,
305 HNF abundances (Gazeau et al., 2020), as well as viral abundance and production increased
306 following the east to west gradient of the initial water conditions.

307 The dust addition induced similar nitrate + nitrite (NO_x) and dissolved inorganic phosphate
308 (DIP) release during all three experiments. Rapid changes were observed on plankton stocks and
309 metabolisms, suggesting that the impact of dust deposition is constrained by the initial
310 composition and metabolic state of the investigated community (Gazeau et al., 2020; 2021).
311 While no direct effect of warming and acidification was observed on the amount of nutrient
312 released from dust, Gazeau et al., (2020, 2021) showed that biological processes were generally



enhanced by these conditions and suggested that deposition may weaken the biological pump in future climate conditions. Here we are further investigating how dust addition in present and future conditions affected, on a short-term scale (≤ 4 days), the microbial trophic interactions and community composition.

4.1. Trophic interactions after dust addition under present and future conditions

Parallel nutrient enrichment incubations conducted in darkness showed that *in situ* heterotrophic bacterioplankton communities, were N, P co-limited at TYR, mainly P limited at ION and N limited at FAST (Van Wambeke et al., 2020). However, the HB appeared to be weakly bottom up controlled (Ducklow, 1992) in our experiment especially in D and G (Fig 2a). Such top-down control on the bacterioplankton has been previously observed in the Mediterranean Sea (Siokou-Frangou et al., 2010) and might increase under future conditions as suggested by the higher top-down index in G ($G = 0.92$ vs. $C/D = 0.80$, Morán et al., 2017).

Bacterial mortality increased relative to controls in D and G at TYR, and only in G at ION and FAST. The weak coupling between bacteria and viruses, as well as the increased virus production and relative abundance of lytic cells (see below), only explained a small fraction of the estimated bacterial mortality (max. 17%), suggesting an additional grazing pressure on bacteria. HNF abundances increased in D at TYR and at all stations in G (Gazeau et al., 2020), which could explain the increased bacterial mortality. Increased grazing rate by HNF on bacteria with dust addition has been previously reported in the Eastern Mediterranean Sea (Tsiola et al., 2017). While our results suggest a strong grazing pressure on bacteria, HNF appeared to be top-down controlled as well (Gasol, 1994, Fig 3b), potentially by the increasing populations of mixotrophic dinoflagellates or Giruses (see below). It is also possible that HB were grazed by



335 mixotrophic nanoflagellates or by larger protozoans, or that the HNF abundance was
336 underestimated by flow cytometry.

337 Considering the seasonal impact of grazing and viral mortality in the Mediterranean Sea,
338 where higher grazing pressure and lysogeny were observed in the stratified nutrient-limited
339 waters in summer (Sánchez et al., 2020), it will be important to further study the seasonal impact
340 of dust deposition on trophic interactions and indirect cascading impact on microbial dynamics
341 and community composition.

342

343 4.2. Viral processes and community during dust enrichment in present and future conditions

344 Viruses represent pivotal components of the marine food web, influencing genome evolution,
345 community dynamics, and ecosystem biogeochemistry (Suttle, 2007). The environmental and
346 evolutionary implications of viral infection differ depending on whether viruses establish lytic or
347 lysogenic infections. Lytic infections produce virion progeny and result in cell destruction while
348 viruses undergoing lysogenic infections can replicate as “dormant” prophages without producing
349 virions or can switch to a lytic productive cycle upon an induction event. Understanding how
350 viral processes are impacted by changes in environmental conditions, is thus crucial to better
351 constrain microbial mortality and cascading impacts on marine ecosystems. Aerosol deposition
352 was already identified as a factor that stimulates virus production and viral induced mortality of
353 bacteria in the Mediterranean Sea (Pulido-Villena et al., 2014; Tsiola et al., 2017) while the
354 impact of future environmental conditions remains more controversial (Larsen et al., 2008;
355 Brussaard et al., 2013; Maat et al., 2014; Vaqué et al., 2019; Malits et al., 2021). The combined



356 effect of aerosol deposition and future conditions of temperature and pH on the viral
357 compartment has, to our knowledge, never been investigated.

358 The rapid changes in viral production and lifestyle observed in all three experiments support the
359 idea that the viral component is sensitive to the environmental variability even on short (hourly)-
360 time scales. The dynamics in viral activities was however impacted differently depending on the
361 treatments and the experiments. Viral production increased in D and G at TYR and only in G at
362 ION and FAST. Regarding the G treatments, increase in viral production was detected before
363 dust addition for all three experiments and remained mostly unchanged for the remaining of the
364 incubation. This suggests that water warming, and acidification were responsible for most
365 changes in viral activities while dusts had no detectable impact in such conditions regardless of
366 the studied station. Based on our results, the most likely explanation for observed changes in
367 viral production is an activation of a lysogenic to lytic switch. The factors that result in prophage
368 induction are still not well constrained, but nutrients pulses and elevated temperatures have been
369 identified as potential stressors (Danovaro et al., 2011 and references therein). Consistent with
370 the observation of N, P co-limited bacterial community at TYR, it is likely that nutrients released
371 from dust upon deposition to surface water activate the productive cycle of temperate viruses at
372 this station. Such mechanism was also speculated during another dust addition study (Pulido-
373 Villena et al., 2014). Under future conditions (G), the low proportion of lysogens was associated
374 to higher frequency of lytically infected cells relative to C and D at TYR and ION. These trends
375 probably reflect an indirect effect of enhanced bacterial growth with increased temperature not
376 only on prophage induction (Danovaro et al., 2011; Vaqué et al., 2019; Mojica and Brussaard,
377 2014) but also on the kinetics of lytic infections. Intriguingly, the enhanced viral production did
378 not translate into marked changes in viral abundance. The abundance of Low DNA virus



379 population, which typically comprises virus of bacteria, actually decreased between t0 and t48h
 380 pointing to possible viral decay, potentially related to an adsorption onto dust particles
 381 (Weinbauer et al., 2009; Yamada et al., 2020) and the potential export of viral particle to deeper
 382 water layers (Van Wambeke et al. 2020). While recurrent patterns emerged from this study, the
 383 amplitude of viral responses varied between the experiments. At TYR, where heterotrophic
 384 metabolism was higher, the dust addition induced higher viral production relative to controls
 385 than at the two other sites, which suggests that viral processes, as other microbial processes, are
 386 dependent on the initial metabolic status of the water.

387 Overall, no marked changes were observed for viral communities and abundances after dust
 388 addition, both under present and future conditions relative to controls, except at FAST where the
 389 abundance of Girus population increased significantly in G from t24h until the end of the
 390 experiment. Giruses typically comprise large double stranded DNA viruses that infect
 391 nanoeukaryotes including photosynthetic (microalgae) and heterotrophic (HNF, amoeba,
 392 choanoflagellate) organisms (Brussaard and Martinez, 2008; Needham et al., 2019; Fischer et al.,
 393 2010; Martínez et al., 2014). The presence of Giruses at FAST in this treatment might be
 394 explained by the increase in nano-eukaryote abundances at t72h and their decline after 96 h of
 395 incubation (Gazeau et al., 2020). The coccolithophore *Emiliania huxleyi* appears as one of the
 396 potential host candidates for these Giruses. The abundance of *E. huxleyi* increased in D and G at
 397 this station and this phytoplankter is known to be infected by such giant viruses (Jacquet et al.,
 398 2002; Schroeder et al., 2002; Pagarete et al., 2011). It is not clear from our results whether
 399 increased Girus abundance is due to the greenhouse effect only (as discussed above for viruses of
 400 HB) or the combination of dust addition and greenhouse effects. While temperature warming
 401 was shown to accelerate viral production in several virus – phytoplankton systems (Mojica and



402 Brussaard 2014, Demory et al. 2017), a temperature-induced resistance to viral infection was
 403 specifically observed in *E. huxleyi* (Kendrick et al., 2014). Previous experiments have also
 404 reported a negative impact of acidification on *E. huxleyi* virus dynamics (Larsen et al., 2008). By
 405 contrast, nutrient release following dust seeding could indirectly stimulate *E. huxleyi* virus
 406 production (Bratbak et al., 1993) or induced switching between non-lethal temperate to lethal
 407 lytic stage (Knowles et al., 2020) under future conditions. Targeted analyses are of course
 408 required to identify the viral populations selected in G and the outcomes of their infection.
 409 Nonetheless, this is the first time, to our knowledge, that dust deposition and enhanced
 410 temperature and acidification have been shown to induce the proliferation of G viruses. The impact
 411 of dust deposition under future environmental conditions on the viral infections processes could
 412 have significant consequences for microbial evolution, food web processes, biogeochemical
 413 cycles, and carbon sequestration.

414

415 4.3 Microbial community dynamic after dust addition under present and future conditions

416 While changes in bacterial community composition during various type of dust addition
 417 experiments have shown only minor transient responses (e.g., Maraňon et al., 2010; Hill et al.,
 418 2010; Laghdass et al., 2011; Pulido-Villena et al., 2014; Marín-Beltrán et al., 2019), here
 419 microbial community structure showed quick, significant and sustained changes in response to
 420 dust addition in all three experiments. Similar to other parameters observed during these
 421 experiments (discussed above and in Gazeau et al., 2020; Gazeau et al., 2021), the degree of
 422 response in terms of community composition was specific to the tested waters.



423 At TYR, where primary production was low, only transient changes after 24 h of incubation
424 were observed, before the micro-eukaryotes community converged back close to initial
425 conditions. In contrast, the bacterial community significantly and rapidly changed after 24 h and
426 remained different after 72 h. At FAST, where the addition of dust appeared to promote
427 autotrophic processes, the micro-eukaryotes community responded quickly 24 h after dust
428 addition, while minor and delayed changes, probably related to the lower BP growth rates
429 compared to the other tested waters, were observed in the bacterial community. At ION both
430 eukaryotes and bacterial community responded to dust addition. The delayed response of micro-
431 eukaryotes after 72 h compared to the quick bacterial response at 24 h suggests that HB were
432 better at competing for nutrient inputs at this station and that autotrophic processes may be
433 responding to bacterial nutrient regeneration after a lag phase, further suggesting the tight
434 coupling between heterotrophic bacteria and phytoplankton at this station. The combined effect
435 of decreased pH and elevated temperature on marine microbes is not yet well understood
436 (reviewed in O'Brien et al., 2016). The absence of significant community changes at TYR and
437 FAST while changes were observed at ION, suggests that the response might be dependent on
438 other environmental factors, which need to be further studied.

439 Dust addition likely favors certain group of micro-organisms, suggesting a quicker response
440 of fast growing/copiotrophic groups as well as the increase of specialized functional groups (Guo
441 et al., 2016; Westrich et al., 2016; Maki et al., 2016). Potential toxicity effects of metals released
442 from dust/aerosols on certain micro-organisms have also been reported (Paytan et al., 2009;
443 Rahav et al., 2020). Here, the micro-eukaryotic community was dominated by a diverse group of
444 dinoflagellates which were responsible for the main variations between treatments at all stations.
445 The overwhelming abundance of dinoflagellates sequences over other micro-eukaryotes could be



446 biased by the large genomes and multiple ribosomal gene copies per genome found in
447 dinoflagellates (Zhu et al., 2005) or due to their preferential amplification. However, the
448 dominance of dinoflagellates in surface water at this time of the year in the Mediterranean Sea is
449 not uncommon (García-Gómez et al., 2020) and was also observed in surface waters of the three
450 sampled stations by Imaging Flow Cytobot (Marañón et al., 2021). While pigment data suggest
451 an increase of haptophytes and pelagophytes in D (Gazeau et al., 2020), the sequencing data only
452 show the presence of *Emiliana huxleyi* as responsible for some of the community changes after
453 dust addition at ION and FAST. These pigments could also indicate the presence of
454 dinoflagellates through tertiary endosymbiosis, in particular *Karlodinium* sp. (Yoon et al., 2002;
455 Zapata et al., 2012), which is an important mixotrophic dinoflagellate (Calbet et al., 2011)
456 observed in D and G at ION and FAST. The variations in dinoflagellate groups might have
457 important trophic impacts due to their diverse mixotrophic states (Stoecker et al., 2017) and the
458 effect of dust addition on mixotrophic interactions should be further studied to better understand
459 the cascading impact of dust on food webs and the biological pump.

460 Positive to toxic impacts on cyanobacteria have been reported from atmospheric deposition
461 experiments (e.g., Paytan et al., 2009; Zhou et al., 2021). Here, *Synechococcus* appeared to be
462 inhibited at TYR while it was enhanced at ION and FAST, especially under future conditions
463 (this study, Gazeau et al., 2020). The same ASVs appeared to be inhibited at TYR and ION
464 while promoted at FAST and a different ASVs increased at ION. *Synechococcus* has recently
465 been shown to be stimulated by wet aerosol addition in P-limited conditions but inhibited in N-
466 limited conditions, in the South China Sea (Zhou et al., 2021). It was also shown to be repressed
467 by dust addition in nutrient limited tropical Atlantic (Marañón et al., 2010). This suggests that



different *Synechococcus* ecotypes (Sohm et al., 2016) might respond differently to dust addition depending on the initial biogeochemical conditions of the water.

In the three experiments, the main bacterial ASVs responsible for the differences between the control and treatments were closely related to different *Alteromonas* strains. *Alteromonas* are ubiquitous in marine environment and can respond rapidly to nutrient pulses (López-Pérez and Rodríguez-Valera, 2014). Some *Alteromonas* are capable to grow on a wide range of carbon compounds (Pedler et al., 2014). They can produce iron binding ligands (Hogle et al., 2016) to rapidly assimilate Fe released from dust. Thus, they could have significant consequences for the marine carbon and Fe cycles during dust deposition events. Other copiotrophic γ -Proteobacteria, such as *Vibrio*, have been observed to bloom after dust deposition in the Atlantic Ocean (Westrich et al., 2016). Guo et al. (2016) using RNA sequencing, also show that γ -Proteobacteria quickly outcompete α -Proteobacteria (mainly SAR11 and Rhodobacterales) that were initially more active. Here, while SAR11 relative abundance decreased in all experiments after 24h, other α -Proteobacteria related to the aerobic anoxygenic phototroph (AAP) *Erythrobacter* sp., increased in response to dust, in particular under future conditions. Other AAP, such as OM60, also responded to dust addition in our experiment and in the Eastern Mediterranean Sea (Guo et al., 2016). Fast growing AAP might quickly outcompete other HB by supplementing their growth with light derived energy (e.g., Koblížek, 2015). They have also been shown to be stimulated by higher temperature (Sato-Takabe et al., 2019). AAP response to dust and future conditions could have a significant role in marine biogeochemical cycles.

5. Conclusion

The microbial food web response to dust addition was dependent on the initial state of the microbial community in the tested waters. A different response in trophic interactions and



community composition of the microbial food web, to the wet dust addition, was observed at each station. Generally greater changes were observed in future conditions. Pulsed input of nutrients and trace metals changed the microbial ecosystem from bottom-up limited to a top-down controlled bacterial community, likely from grazing and induced lysogeny. The composition of mixotrophic microeukaryotes and phototrophic prokaryotes was also altered.

Overall, the impact of such simulated pulsed nutrient deposition will depend on the initial biogeochemical conditions of the ecosystem, with likely possible large impact on microbial trophic interactions and community structure. All effects might be generally enhanced in future climate scenarios. The impact of dust deposition on metabolic processes and consequences for the carbon and nitrogen cycles and the biological pump based on these minicosm experiments are further discussed in Gazeau et al. (2021), and the *in situ* effect of a wet dust deposition event is explored in Van Wambeke et al. (2020), in this special issue.

6. Data availability

Guieu et al., Biogeochemical dataset collected during the PEACETIME cruise. SEANOE. <https://doi.org/10.17882/75747> (2020). All sequences associated with this study have been deposited under the BioProject ID: PRJNA693966.

7. Author contributions

FG and CG designed the experiment. All authors participated in sampling or sample processes. JD analyzed the data and wrote the paper with contributions from all authors.

8. Competing interests

The authors declare that they have no conflict of interest.



513 **9. Special issue statement**

514 This article is part of the special issue ‘Atmospheric deposition in the low-nutrient–low-
515 chlorophyll (LNLC) ocean: effects on marine life today and in the future (ACP/BG inter-
516 journal SI)’. It is not associated with a conference.

517 **10. Financial support**

518 Part of this research was funded by the ANR CALYPSO attributed to ACB (ANR-15-CE01-
519 0009). EM was supported by the Spanish Ministry of Science, Innovation and Universities
520 through grant PGC2018-094553B-I00. JD was funded by a Marie Curie Actions-International
521 Outgoing Fellowship (PIOF-GA-2013-629378).

522
523 **11. Acknowledgements**

524 This study is a contribution to the PEACETIME project (<http://peacetime-project.org>,
525 <https://doi.org/10.17600/17000300>), a joint initiative of the MERMEX and ChArMEx
526 components supported by CNRS-INSU, IFREMER, CEA, and Météo-France as part of the
527 programme MISTRALS coordinated by INSU. PEACETIME was endorsed as a process study
528 by GEOTRACES and is also a contribution to SOLAS. We gratefully acknowledge the onboard
529 support from the captain and crew of the RV Pourquoi Pas? and of our chief scientists C. Guieu
530 and K. Desboeufs. We also thank K. Djaoudi for her assistance in sampling the minicosms, P.
531 Catala, B. Marie and M. Perez-Lorenzo with their assistance in measuring microbial abundance,
532 DOC concentration and primary production.

533
534 **12. References**



535 Allen, R., Hoffmann, L. J., Law, C. S., and Summerfield, T. C.: Subtle bacterioplankton
 536 community responses to elevated CO₂ and warming in the oligotrophic South Pacific gyre, *Env*
 537 *Microbiol Rep*, 12, 377-386, <https://doi.org/10.1111/1758-2229.12844>, 2020.

538 Bolyen, E., Rideout, J. R., Dillon, M. R., Bokulich, N. A., Abnet, C. C., Al-Ghalith, G. A.,
 539 Alexander, H., Alm, E. J., Arumugam, M., Asnicar, F., Bai, Y., Bisanz, J. E., Bittinger, K.,
 540 Brejnrod, A., Brislawn, C. J., Brown, C. T., Callahan, B. J., Caraballo-Rodríguez, A. M.,
 541 Chase, J., Cope, E. K., Da Silva, R., Diener, C., Dorrestein, P. C., Douglas, G. M., Durall, D.
 542 M., Duvallet, C., Edwardson, C. F., Ernst, M., Estaki, M., Fouquier, J., Gauglitz, J. M.,
 543 Gibbons, S. M., Gibson, D. L., Gonzalez, A., Gorlick, K., Guo, J., Hillmann, B., Holmes, S.,
 544 Holste, H., Huttenhower, C., Huttley, G. A., Janssen, S., Jarmusch, A. K., Jiang, L., Kaehler,
 545 B. D., Kang, K. B., Keefe, C. R., Keim, P., Kelley, S. T., Knights, D., Koester, I., Kosciulek,
 546 T., Kreps, J., Langille, M. G. I., Lee, J., Ley, R., Liu, Y.-X., Loftfield, E., Lozupone, C.,
 547 Maher, M., Marotz, C., Martin, B. D., McDonald, D., McIver, L. J., Melnik, A. V., Metcalf, J.
 548 L., Morgan, S. C., Morton, J. T., Naimey, A. T., Navas-Molina, J. A., Nothias, L. F.,
 549 Orchanian, S. B., Pearson, T., Peoples, S. L., Petras, D., Preuss, M. L., Priesse, E., Rasmussen,
 550 L. B., Rivers, A., Robeson, M. S., Rosenthal, P., Segata, N., Shaffer, M., Shiffer, A., Sinha, R.,
 551 Song, S. J., Spear, J. R., Swafford, A. D., Thompson, L. R., Torres, P. J., Trinh, P., Tripathi,
 552 A., Turnbaugh, P. J., Ul-Hasan, S., van der Hooft, J. J. J., Vargas, F., Vázquez-Baeza, Y.,
 553 Vogtmann, E., von Hippel, M., Walters, W., Wan, Y., Wang, M., Warren, J., Weber, K. C.,
 554 Williamson, C. H. D., Willis, A. D., Xu, Z. Z., Zaneveld, J. R., Zhang, Y., Zhu, Q., Knight, R.,
 555 and Caporaso, J. G.: Reproducible, interactive, scalable and extensible microbiome data
 556 science using QIIME 2, *Nat Biotechnol*, 37, 852-857, [10.1038/s41587-019-0209-9](https://doi.org/10.1038/s41587-019-0209-9), 2019.



- 557 Bonnet, S., and Guieu, C.: Atmospheric forcing on the annual iron cycle in the Western
 558 Mediterranean Sea: A 1-year survey, *J Geophys Res-Oceans*, 111,
 559 <https://doi.org/10.1029/2005JC003213>, 2006.
- 560 Bosc, E., Bricaud, A., and Antoine, D.: Seasonal and interannual variability in algal biomass and
 561 primary production in the Mediterranean Sea, as derived from 4 years of SeaWiFS
 562 observations, *Global Biogeochem Cy*, 18, <https://doi.org/10.1029/2003GB002034>, 2004.
- 563 Bratbak, G., Egge, J. K., and Haldal, M.: Viral mortality of the marine alga *Emiliania huxleyi*
 564 (Haptophyceae) and termination of algal blooms, *Mar Ecol Prog Ser*, 39-48, 1993.
- 565 Brussaard, C., Noordeloos, A., Witte, H., Collenteur, M., Schulz, K. G., Ludwig, A., and
 566 Riebesell, U.: Arctic microbial community dynamics influenced by elevated CO₂ levels,
 567 *Biogeosciences*, 10, 719-731, 2013.
- 568 Brussaard, C. P., and Martinez, J. M.: Algal bloom viruses, *Plant Viruses*, 2, 1-13, 2008.
- 569 Brussaard, C. P. D.: Optimization of Procedures for Counting Viruses by Flow Cytometry, *Appl*
 570 *Environ Microb*, 70, 1506-1513, [10.1128/aem.70.3.1506-1513.2004](https://doi.org/10.1128/aem.70.3.1506-1513.2004), 2004.
- 571 Calbet, A., Bertos, M., Fuentes-Grünwald, C., Alacid, E., Figueroa, R., Renom, B., and Garcés,
 572 E.: Intraspecific variability in *Karlodinium veneficum*: growth rates, mixotrophy, and lipid
 573 composition, *Harmful Algae*, 10, 654-667, 2011.
- 574 Callahan, B. J., McMurdie, P. J., Rosen, M. J., Han, A. W., Johnson, A. J. A., and Holmes, S. P.:
 575 DADA2: High-resolution sample inference from Illumina amplicon data, *Nat Methods*, 13,
 576 581, [10.1038/nmeth.3869](https://doi.org/10.1038/nmeth.3869), 2016.
- 577 Clarke, K. R., and Warwick, P. E.: *Change in Marine Communities: An Approach to Statistical*
 578 *Analysis and Interpretation*, Plymouth, Ltd ed., 2001.



579 D'Ortenzio, F., Iudicone, D., de Boyer Montegut, C., Testor, P., Antoine, D., Marullo, S.,
 580 Santoleri, R., and Madec, G.: Seasonal variability of the mixed layer depth in the
 581 Mediterranean Sea as derived from *in situ* profiles, *Geophys Res Let*, 32,
 582 <https://doi.org/10.1029/2005GL022463>, 2005.

583 Danovaro, R., Corinaldesi, C., Dell'Anno, A., Fuhrman, J. A., Middelburg, J. J., Noble, R. T.,
 584 and Suttle, C. A.: Marine viruses and global climate change, *FEMS Microbiol Rev*, 35, 993-
 585 1034, [10.1111/j.1574-6976.2010.00258.x](https://doi.org/10.1111/j.1574-6976.2010.00258.x), 2011.

586 Degerman, R., Dinasquet, J., Riemann, L., Sjostedt de Luna, S., and Andersson, A.: Effect of
 587 resource availability on bacterial community responses to increased temperature, *Aquat Microb*
 588 *Ecol*, 68: 131-142, 2012.

589 Desboeufs, K., Leblond, N., Wagener, T., Bon Nguyen, E., and Guieu, C.: Chemical fate and
 590 settling of mineral dust in surface seawater after atmospheric deposition observed from dust
 591 seeding experiments in large mesocosms, *Biogeosciences*, 11, 5581-5594, [10.5194/bg-11-](https://doi.org/10.5194/bg-11-5581-2014)
 592 [5581-2014](https://doi.org/10.5194/bg-11-5581-2014), 2014.

593 Ducklow, H.: Factors regulating bottom-up control of bacteria biomass in open ocean plankton
 594 communities, *Arch. Hydrobiol. Beih. Ergebn. Limnol*, 37, 207-217, 1992.

595 Durrieu de Madron, X., Guieu, C., Sempéré, R., Conan, P., Cossa, D., D'Ortenzio, F., Estournel,
 596 C., Gazeau, F., Rabouille, C., Stemmann, L., Bonnet, S., Diaz, F., Koubbi, P., Radakovitch, O.,
 597 Babin, M., Baklouti, M., Bancon-Montigny, C., Belviso, S., Bensoussan, N., Bonsang, B.,
 598 Bouloubassi, I., Brunet, C., Cadiou, J. F., Carlotti, F., Chami, M., Charmasson, S., Charrière,
 599 B., Dachs, J., Doxaran, D., Dutay, J. C., Elbaz-Poulichet, F., Eléaume, M., Eyrolles, F.,
 600 Fernandez, C., Fowler, S., Francour, P., Gaertner, J. C., Galzin, R., Gasparini, S., Ghiglione, J.
 601 F., Gonzalez, J. L., Goyet, C., Guidi, L., Guizien, K., Heimbürger, L. E., Jacquet, S. H. M.,



602 Jeffrey, W. H., Joux, F., Le Hir, P., Leblanc, K., Lefèvre, D., Lejeusne, C., Lemé, R., Loÿe-
 603 Pilot, M. D., Mallet, M., Méjanelle, L., Mélin, F., Mellon, C., Mérigot, B., Merle, P. L., Migon,
 604 C., Miller, W. L., Mortier, L., Mostajir, B., Mousseau, L., Moutin, T., Para, J., Pérez, T.,
 605 Petrenko, A., Poggiale, J. C., Prieur, L., Pujo-Pay, M., Pulido, V., Raimbault, P., Rees, A. P.,
 606 Ridame, C., Rontani, J. F., Ruiz Pino, D., Sicre, M. A., Taillandier, V., Tamburini, C., Tanaka,
 607 T., Taupier-Letage, I., Tedetti, M., Testor, P., Thébault, H., Thouvenin, B., Touratier, F.,
 608 Tronczynski, J., Ulses, C., Van Wambeke, F., Vantrepotte, V., Vaz, S., and Verney, R.: Marine
 609 ecosystems' responses to climatic and anthropogenic forcings in the Mediterranean, Prog
 610 Oceanog, 91, 97-166, <https://doi.org/10.1016/j.pocean.2011.02.003>, 2011.

611 Fischer, M. G., Allen, M. J., Wilson, W. H., and Suttle, C. A.: Giant virus with a remarkable
 612 complement of genes infects marine zooplankton, P Natl Acad Sci 107, 19508-19513, 2010.

613 García-Gómez, C., Yebra, L., Cortés, D., Sánchez, A., Alonso, A., Valcárcel-Pérez, N., Gómez-
 614 Jakobsen, F., Herrera, I., Johnstone, C., and Mercado, J. M.: Shifts in the protist community
 615 associated with an anticyclonic gyre in the Alboran Sea (Mediterranean Sea), FEMS Microbiol
 616 Ecol, 96, 10.1093/femsec/fiaa197, 2020.

617 Gasol, J. M.: A framework for the assessment of top-down vs bottom-up control of heterotrophic
 618 nanoflagellate abundance, Mar ecol prog ser. 113, 291-300, 1994.

619 Gazeau, F., Ridame, C., Van Wambeke, F., Alliouane, S., Stolpe, C., Irisson, J. O., Marro, S.,
 620 Grisoni, J. M., De Liège, G., Nunige, S., Djaoudi, K., Pulido-Villena, E., Dinasquet, J.,
 621 Obernosterer, I., Catala, P., and Guieu, C.: Impact of dust enrichment on Mediterranean
 622 plankton communities under present and future conditions of pH and temperature: an
 623 experimental overview, Biogeosciences Discuss., 2020, 1-81, 10.5194/bg-2020-202, 2020.



- 624 Gazeau, F., Van Wambeke, F., Marañón, E., Pérez-Lorenzo, M., Alliouane, S., Stolpe, C.,
 625 Blasco, T., Leblond, N., Zäncker, B., Engel, A., Marie, B., Dinasquet, J., and Guieu, C.: Impact
 626 of dust addition on the metabolism of Mediterranean plankton communities and carbon export
 627 under present and future conditions of pH and temperature, *Biogeosciences Discuss.*,
 628 <https://doi.org/10.5194/bg-2021-20>, in review, 2021.
- 629 Guieu, C., Aumont, O., Paytan, A., Bopp, L., Law, C. S., Mahowald, N., Achterberg, E. P.,
 630 Marañón, E., Salihoglu, B., Crise, A., Wagener, T., Herut, B., Desboeufs, K., Kanakidou, M.,
 631 Olgun, N., Peters, F., Pulido-Villena, E., Tovar-Sanchez, A., and Völker, C.: The significance
 632 of the episodic nature of atmospheric deposition to Low Nutrient Low Chlorophyll regions,
 633 *Global Biogeochem Cy*, 28, 1179-1198, 10.1002/2014GB004852, 2014.
- 634 Guieu, C. and Ridame, C.: Impact of atmospheric deposition on marine chemistry and
 635 biogeochemistry, in *Atmospheric Chemistry in the Mediterranean Region: Comprehensive
 636 Diagnosis and Impacts*, edited by F. Dulac, S. Sauvage, and E. Hamonou, Springer, Cham,
 637 Switzerland, 2020.
- 638 Guillou, L., Bachar, D., Audic, S., Bass, D., Berney, C., Bittner, L., Boutte, C., Burgaud, G., de
 639 Vargas, C., Decelle, J., del Campo, J., Dolan, J. R., Dunthorn, M., Edvardsen, B., Holzmann,
 640 M., Kooistra, W. H. C. F., Lara, E., Le Bescot, N., Logares, R., Mahé, F., Massana, R.,
 641 Montresor, M., Morard, R., Not, F., Pawlowski, J., Probert, I., Sauvadet, A.-L., Siano, R.,
 642 Stoeck, T., Vaulot, D., Zimmermann, P., and Christen, R.: The Protist Ribosomal Reference
 643 database (PR2): a catalog of unicellular eukaryote Small Sub-Unit rRNA sequences with
 644 curated taxonomy, *Nucleic Acids Res*, 41, D597-D604, 10.1093/nar/gks1160, 2013.
- 645 Guo, C., Xia, X., Pitta, P., Herut, B., Rahav, E., Berman-Frank, I., Giannakourou, A., Tsiola, A.,
 646 Tsagaraki, T. M., and Liu, H.: Shifts in Microbial Community Structure and Activity in the



647 Ultra-Oligotrophic Eastern Mediterranean Sea Driven by the Deposition of Saharan Dust and
 648 European Aerosols, *Front Mar Sci*, 3, 10.3389/fmars.2016.00170, 2016.

649 Highfield, A., Joint, I., Gilbert, J. A., Crawford, K. J., and Schroeder, D. C.: Change in *Emiliana*
 650 *huxleyi* Virus Assemblage Diversity but Not in Host Genetic Composition during an Ocean
 651 Acidification Mesocosm Experiment, *Viruses*, 9, 41, 2017.

652 Hill, P. G., Zubkov, M. V., and Purdie, D. A.: Differential responses of *Prochlorococcus* and
 653 SAR11-dominated bacterioplankton groups to atmospheric dust inputs in the tropical Northeast
 654 Atlantic Ocean, *FEMS Microbiol Let*, 306, 82-89, 10.1111/j.1574-6968.2010.01940.x, 2010.

655 Hogle, S. L., Bundy, R. M., Blanton, J. M., Allen, E. E., and Barbeau, K. A.: Copiotrophic
 656 marine bacteria are associated with strong iron-binding ligand production during
 657 phytoplankton blooms, *Limnol Oceanogr Let*, 10.1002/lol2.10026, 2016.

658 Hu, C., Li, X., He, M., Jiang, P., Long, A., and Xu, J.: Effect of Ocean Acidification on Bacterial
 659 Metabolic Activity and Community Composition in Oligotrophic Oceans, Inferred From Short-
 660 Term Bioassays, *Front Microbiol*, 12, 10.3389/fmicb.2021.583982, 2021.

661 IPCC: Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to
 662 the Fifth Assessment Report of the Intergovernmental Panel on Climate Change, Cambridge
 663 University Press, Cambridge, United Kingdom and New York, NY, USA, 1535 pp., 2014.

664 Jacquet, S., Heldal, M., Iglesias-Rodriguez, D., Larsen, A., Wilson, W., and Bratbak, G.: Flow
 665 cytometric analysis of an *Emiliana huxleyi* bloom terminated by viral infection, *Aquat Microb*
 666 *Ecol*, 27, 111-124, 2002.

667 Kendrick, B. J., DiTullio, G. R., Cyronak, T. J., Fulton, J. M., Van Mooy, B. A., and Bidle, K.
 668 D.: Temperature-induced viral resistance in *Emiliana huxleyi* (Prymnesiophyceae), *PLoS One*,
 669 9, e112134, 2014.



670 Kirchman, D., Knees, E., and Hodson, R.: Leucine Incorporation and Its Potential As A Measure
 671 of Protein-Synthesis by Bacteria in Natural Aquatic Systems, *Appl Environ Microbiol*, 49,
 672 599-607, 1985.

673 Knowles, B., Bonachela, J. A., Behrenfeld, M. J., Bondoc, K. G., Cael, B., Carlson, C. A.,
 674 Cieslik, N., Diaz, B., Fuchs, H. L., and Graff, J. R.: Temperate infection in a virus–host system
 675 previously known for virulent dynamics, *Nat comms*, 11, 1-13, 2020.

676 Koblížek, M.: Ecology of aerobic anoxygenic phototrophs in aquatic environments, *FEMS*
 677 *Microbiol Rev*, 39, 854-870, 10.1093/femsre/fuv032, 2015.

678 Krause, E., Wichels, A., Giménez, L., Lunau, M., Schilhabel, M. B., and Gerdts, G.: Small
 679 Changes in pH Have Direct Effects on Marine Bacterial Community Composition: A
 680 Microcosm Approach, *PLOS ONE*, 7, e47035, 10.1371/journal.pone.0047035, 2012.

681 Laghdass, M., Blain, S., Besseling, M., Catala, P., Guieu, C., and Obernosterer, I.: Effects of
 682 Saharan dust on the microbial community during a large *in situ* mesocosm experiment in the
 683 NW Mediterranean Sea, *Aquat Microb Ecol*, 62, 201-213, 2011.

684 Larsen, J. B., Larsen, A., Thyrraug, R., Bratbak, G., and Sandaa, R.-A.: Response of marine
 685 viral populations to a nutrient induced phytoplankton bloom at different $p\text{CO}_2$ levels,
 686 *Biogeosciences*, 5, 523-533, 2008.

687 Lee, S. H., and Fuhrman, J. A.: Relationships between biovolume and biomass of naturally
 688 derived marine bacterioplankton, *Appl Environ Microbiol*, 53, 1298-1303, 1987.

689 Loýe-Pilot, M., and Martin, J.: Saharan dust input to the western Mediterranean: an eleven years
 690 record in Corsica, in: *The impact of desert dust across the Mediterranean*, Springer, 191-199,
 691 1996.



692 López-Pérez, M., and Rodríguez-Valera, F.: The Family Alteromonadaceae, in: The Prokaryotes:
 693 Gammaproteobacteria, edited by: Rosenberg, E., DeLong, E. F., Lory, S., Stackebrandt, E., and
 694 Thompson, F., Springer Berlin Heidelberg, Berlin, Heidelberg, 69-92, 2014.

695 Maat, D. S., Crawford, K. J., Timmermans, K. R., and Brussaard, C. P.: Elevated carbon dioxide
 696 and phosphorus limitation favor *Micromonas pusilla* through stimulated growth and reduced
 697 viral impact, aspects of algal host-virus interactions in a changing ocean, 80, 29, 2014.

698 Mahowald, N. M., Scanza, R., Brahney, J., Goodale, C. L., Hess, P. G., Moore, J. K., and Neff,
 699 J.: Aerosol Deposition Impacts on Land and Ocean Carbon Cycles, Current Climate Change
 700 Reports, 3, 16-31, 10.1007/s40641-017-0056-z, 2017.

701 Maki, T., Ishikawa, A., Mastunaga, T., Pointing, S. B., Saito, Y., Kasai, T., Watanabe, K., Aoki,
 702 K., Horiuchi, A., Lee, K. C., Hasegawa, H., and Iwasaka, Y.: Atmospheric aerosol deposition
 703 influences marine microbial communities in oligotrophic surface waters of the western Pacific
 704 Ocean, Deep Sea Research Part I: Oceanographic Research Papers, 118, 37-45,
 705 <https://doi.org/10.1016/j.dsr.2016.10.002>, 2016.

706 Malits, A., Boras, J.A., Balagué, V., Calvo, E., Gasol, J.M., Marrasé, C., Pelejero, C., Pinhassi,
 707 J., Sala, M.M. and Vaqué, D. Viral-Mediated Microbe Mortality Modulated by Ocean
 708 Acidification and Eutrophication: Consequences for the Carbon Fluxes Through the Microbial
 709 Food Web. *Front. Microbiol.* 12:635821. doi: 10.3389/fmicb.2021.635821, 2021

710 Marañón, E., Fernández, A., Mouriño-Carballido, B., Martínez-García, S., Teira, E., Cermeño,
 711 P., Chouciño, P., Huete-Ortega, M., Fernández, E., Calvo-Díaz, A., Morán, X. A. G., Bode, A.,
 712 Moreno-Ostos, E., Varela, M. M., Patey, M. D., and Achterberg, E. P.: Degree of oligotrophy
 713 controls the response of microbial plankton to Saharan dust, Limnology and Oceanography, 55,
 714 2339-2352, <https://doi.org/10.4319/lo.2010.55.6.2339>, 2010.



715 Marañón, E., Lorenzo, M. P., Cermeño, P., and Mouriño-Carballido, B.: Nutrient limitation
 716 suppresses the temperature dependence of phytoplankton metabolic rates, *The ISME Journal*,
 717 12, 1836-1845, 10.1038/s41396-018-0105-1, 2018.

718 Marañón, E., Van Wambeke, F., Uitz, J., Boss, E. S., Pérez-Lorenzo, M., Dinasquet, J.,
 719 Haëntjens, N., Dimier, C., and Taillandier, V.: Deep maxima of phytoplankton biomass,
 720 primary production and bacterial production in the Mediterranean Sea during late spring,
 721 *Biogeosciences*, 18, 1749–1767, <https://doi.org/10.5194/bg-18-1749-2021>, 2021

722 Marín-Beltrán, I., Logue, J. B., Andersson, A. F., and Peters, F.: Atmospheric Deposition Impact
 723 on Bacterial Community Composition in the NW Mediterranean, *Frontiers in Microbiology*,
 724 10, 10.3389/fmicb.2019.00858, 2019.

725 Martínez, J. M., Swan, B. K., and Wilson, W. H.: Marine viruses, a genetic reservoir revealed by
 726 targeted viromics, *The ISME Journal*, 8, 1079-1088, 10.1038/ismej.2013.214, 2014.

727 Mojica, K. D., and Brussaard, C. P.: Factors affecting virus dynamics and microbial host–virus
 728 interactions in marine environments, *FEMS microbiology ecology*, 89, 495-515, 2014.

729 Morán, X. A. G., Gasol, J. M., Pernice, M. C., Mangot, J.-F., Massana, R., Lara, E., Vaqué, D.,
 730 and Duarte, C. M.: Temperature regulation of marine heterotrophic prokaryotes increases
 731 latitudinally as a breach between bottom-up and top-down controls, *Global Change Biology*,
 732 23, 3956-3964, <https://doi.org/10.1111/gcb.13730>, 2017.

733 Morán, X. A. G., Baltar, F., Carreira, C., and Lønborg, C.: Responses of physiological groups of
 734 tropical heterotrophic bacteria to temperature and dissolved organic matter additions: food
 735 matters more than warming, *Environmental Microbiology*, 22, 1930-1943,
 736 <https://doi.org/10.1111/1462-2920.15007>, 2020.



737 Moulin, C., and Chiapello, I.: Impact of human-induced desertification on the intensification of
 738 Sahel dust emission and export over the last decades, *Geophysical Research Letters*, 33,
 739 <https://doi.org/10.1029/2006GL025923>, 2006.

740 Needham, D. M., Yoshizawa, S., Hosaka, T., Poirier, C., Choi, C. J., Hehenberger, E., Irwin, N.
 741 A., Wilken, S., Yung, C.-M., and Bachy, C.: A distinct lineage of giant viruses brings a
 742 rhodopsin photosystem to unicellular marine predators, *Proceedings of the National Academy*
 743 *of Sciences*, 116, 20574-20583, 2019.

744 O'Brien, P. A., Morrow, K. M., Willis, B. L., and Bourne, D. G.: Implications of Ocean
 745 Acidification for Marine Microorganisms from the Free-Living to the Host-Associated,
 746 *Frontiers in Marine Science*, 3, 10.3389/fmars.2016.00047, 2016.

747 Pagarete, A., Le Corguillé, G., Tiwari, B., Ogata, H., de Vargas, C., Wilson, W. H., and Allen,
 748 M. J.: Unveiling the transcriptional features associated with coccolithovirus infection of natural
 749 *Emiliana huxleyi* blooms, *FEMS Microbiology Ecology*, 78, 555-564, 10.1111/j.1574-
 750 6941.2011.01191.x, 2011.

751 Parada, A. E., Needham, D. M., and Fuhrman, J. A.: Every base matters: assessing small subunit
 752 rRNA primers for marine microbiomes with mock communities, time series and global field
 753 samples, *Environmental Microbiology*, 18, 1403-1414, 10.1111/1462-2920.13023, 2016.

754 Parada, V., Herndl, G. J., and Weinbauer, M. G.: Viral burst size of heterotrophic prokaryotes in
 755 aquatic systems, *JMBA-Journal of the Marine Biological Association of the United Kingdom*,
 756 86, 613, 2006.

757 Paytan, A., Mackey, K. R. M., Chen, Y., Lima, I. D., Doney, S. C., Mahowald, N., Labiosa, R.,
 758 and Post, A. F.: Toxicity of atmospheric aerosols on marine phytoplankton, *Proceedings of the*
 759 *National Academy of Sciences*, 106, 4601-4605, 10.1073/pnas.0811486106, 2009.



760 Pedler, B. E., Aluwihare, L. I., and Azam, F.: Single bacterial strain capable of significant
 761 contribution to carbon cycling in the surface ocean, *Proceedings of the National Academy of*
 762 *Sciences of the United States of America*, 111, 7202-7207, 2014.

763 Pinheiro, J., Bates, D., DebRoy, S., and Sarkar, D.: R Core Team. nlme: linear and nonlinear
 764 mixed effects models. R package version 3.1-117, Available at [http://CRAN.R-project.](http://CRAN.R-project.org/package=nlme)
 765 [org/package= nlme](http://CRAN.R-project.org/package=nlme), 2014.

766 Pitta, P., Kanakidou, M., Mihalopoulos, N., Christodoulaki, S., Dimitriou, P.D., Frangoulis, C.,
 767 Giannakourou, A., Kagiorgi, M., Lagaria, A., Nikolaou, P., Papageorgiou, N., Psarra, S., Santi,
 768 I., Tsapakis, M., Tsiola, A., Violaki, K., and Petihakis, G. Saharan Dust Deposition Effects on
 769 the Microbial Food Web in the Eastern Mediterranean: A Study Based on a Mesocosm
 770 Experiment. *Front. Mar. Sci.* 4:117. doi: 10.3389/fmars.2017.00117, 2017

771 Pulido-Villena, E., Baudoux, A. C., Obernosterer, I., Landa, M., Caparros, J., Catala, P.,
 772 Georges, C., Harmand, J., and Guieu, C.: Microbial food web dynamics in response to a
 773 Saharan dust event: results from a mesocosm study in the oligotrophic Mediterranean Sea,
 774 *Biogeosciences*, 11, 5607-5619, 10.5194/bg-11-5607-2014, 2014.

775 Quast, C., Pruesse, E., Yilmaz, P., Gerken, J., Schweer, T., Yarza, P., Peplies, J., and Glöckner,
 776 F. O.: The SILVA ribosomal RNA gene database project: improved data processing and web-
 777 based tools, *Nucleic Acids Research*, 41, D590-D596, 10.1093/nar/gks1219, 2013.

778 R Core Team: R: A language and environment for statistical computing. , R Foundation for
 779 Statistical Computing, Vienna, Austria., <https://www.R-project.org/>. 2020.

780 Rahav, E., Paytan, A., Mescioglou, E., Bar-Zeev, E., Martínez Ruiz, F., Xian, P., and Herut, B.:
 781 Bio-Aerosols Negatively Affect *Prochlorococcus* in Oligotrophic Aerosol-Rich Marine
 782 Regions, *Atmosphere*, 11, 540, 2020.



783 Ridame, C., Le Moal, M., Guieu, C., TERNON, E., Biegala, I. C., L'Helguen, S., and Pujo-Pay, M.:
 784 Nutrient control of N₂ fixation in the oligotrophic Mediterranean Sea and the impact of
 785 Saharan dust events, *Biogeosciences*, 8, 2773-2783, 10.5194/bg-8-2773-2011, 2011.
 786 Sánchez, O., Ferrera, I., Mabrito, I., Gazulla, C. R., Sebastián, M., Auladell, A., Marín-Vindas,
 787 C., Cardelús, C., Sanz-Sáez, I., Pernice, M. C., Marrasé, C., Sala, M. M., and Gasol, J. M.:
 788 Seasonal impact of grazing, viral mortality, resource availability and light on the group-
 789 specific growth rates of coastal Mediterranean bacterioplankton, *Scientific Reports*, 10, 19773,
 790 10.1038/s41598-020-76590-5, 2020.
 791 Sato-Takabe, Y., Hamasaki, K., and Suzuki, S.: High temperature accelerates growth of aerobic
 792 anoxygenic phototrophic bacteria in seawater, *MicrobiologyOpen*, 8, e00710-e00710,
 793 10.1002/mbo3.710, 2019.
 794 Schroeder, D., Oke, J., Malin, G., and Wilson, W.: Coccolithovirus (Phycodnaviridae):
 795 characterisation of a new large dsDNA algal virus that infects *Emiliana huxleyi*, *Archives of*
 796 *virology*, 147, 1685-1698, 2002.
 797 Siokou-Frangou, I., Christaki, U., Mazzocchi, M. G., Montresor, M., Ribera d'Alcalá, M., Vaqué,
 798 D., and Zingone, A.: Plankton in the open Mediterranean Sea: a review, *Biogeosciences*, 7,
 799 1543-1586, 10.5194/bg-7-1543-2010, 2010.
 800 Smith, D. C., and Azam, F.: A simple, economical method for measuring bacterial protein
 801 synthesis rates in seawater using ³H-leucine, *Marine microbial food webs*, 6, 102-114, 1992.
 802 Sohm, J. A., Ahlgren, N. A., Thomson, Z. J., Williams, C., Moffett, J. W., Saito, M. A., Webb,
 803 E. A., and Rocap, G.: Co-occurring *Synechococcus* ecotypes occupy four major oceanic
 804 regimes defined by temperature, macronutrients and iron, *The ISME journal*, 10, 333-345,
 805 10.1038/ismej.2015.115, 2016.



806 Stoeck, T., Bass, D., Nebel, M., Christen, R., Jones, M. D. M., Breiner, H. W., and Richards, T.
 807 A.: Multiple marker parallel tag environmental DNA sequencing reveals a highly complex
 808 eukaryotic community in marine anoxic water, *Molecular Ecology*, 19, 21-31, 2010.

809 Stoecker, D. K., Hansen, P. J., Caron, D. A., and Mitra, A.: Mixotrophy in the Marine Plankton,
 810 *Annual Review of Marine Science*, 9, 311-335, 10.1146/annurev-marine-010816-060617,
 811 2017.

812 Suttle, C. A.: Marine viruses — major players in the global ecosystem, *Nature Reviews*
 813 *Microbiology*, 5, 801-812, 10.1038/nrmicro1750, 2007.

814 Ternon, E., Guieu, C., Loÿe-Pilot, M. D., Leblond, N., Bosc, E., Gasser, B., Miquel, J. C., and
 815 Martín, J.: The impact of Saharan dust on the particulate export in the water column of the
 816 North Western Mediterranean Sea, *Biogeosciences*, 7, 809-826, 10.5194/bg-7-809-2010, 2010.

817 Tsiola, A., Tsagaraki, T. M., Giannakourou, A., Nikolioudakis, N., Yücel, N., Herut, B., and
 818 Pitta, P.: Bacterial Growth and Mortality after Deposition of Saharan Dust and Mixed Aerosols
 819 in the Eastern Mediterranean Sea: A Mesocosm Experiment, *Frontiers in Marine Science*, 3,
 820 10.3389/fmars.2016.00281, 2017.

821 Van Wambeke, F., Taillandier, V., Deboeufs, K., Pulido-Villena, E., Dinasquet, J., Engel, A.,
 822 Marañón, E., Ridame, C., and Guieu, C.: Influence of atmospheric deposition on
 823 biogeochemical cycles in an oligotrophic ocean system, *Biogeosciences Discuss.*, 2020, 1-51,
 824 10.5194/bg-2020-411.

825 Vaqué, D., Lara, E., Arrieta, J. M., Holding, J., Sà, E. L., Hendriks, I. E., Coello-Camba, A.,
 826 Alvarez, M., Agustí, S., Wassmann, P. F., and Duarte, C. M.: Warming and CO₂ Enhance
 827 Arctic Heterotrophic Microbial Activity, *Frontiers in Microbiology*, 10,
 828 10.3389/fmicb.2019.00494, 2019.



- 829 Weinbauer, M., Bettarel, Y., Cattaneo, R., Luef, B., Maier, C., Motegi, C., Peduzzi, P., and Mari,
 830 X.: Viral ecology of organic and inorganic particles in aquatic systems: avenues for further
 831 research, *Aquatic Microbial Ecology*, 57, 321-341, 2009.
- 832 Weinbauer, M. G., Winter, C., and Höfle, M. G.: Reconsidering transmission electron
 833 microscopy based estimates of viral infection of bacterio-plankton using conversion factors
 834 derived from natural communities, *Aquatic Microbial Ecology*, 27, 103-110, 2002.
- 835 Weinbauer, M. G., Rowe, J. M., and Wilhelm, S.: Determining rates of virus production in
 836 aquatic systems by the virus reduction approach, 2010.
- 837 Westrich, J. R., Ebling, A. M., Landing, W. M., Joyner, J. L., Kemp, K. M., Griffin, D. W., and
 838 Lipp, E. K.: Saharan dust nutrients promote *Vibrio* bloom formation in marine surface waters,
 839 *Proceedings of the National Academy of Sciences*, 113, 5964-5969, 10.1073/pnas.1518080113,
 840 2016.
- 841 Winter, C., Herndl, G. J., and Weinbauer, M. G.: Diel cycles in viral infection of
 842 bacterioplankton in the North Sea, *Aquatic Microbial Ecology*, 35, 207-216, 2004.
- 843 Yamada, Y., Guillemette, R., Baudoux, A.-C., Patel, N., and Azam, F.: Viral Attachment to
 844 Biotic and Abiotic Surfaces in Seawater, *Applied and Environmental Microbiology*, 86,
 845 e01687-01619, 10.1128/aem.01687-19, 2020.
- 846 Yoon, H. S., Hackett, J. D., and Bhattacharya, D.: A single origin of the peridinin- and
 847 fucoxanthin-containing plastids in dinoflagellates through tertiary endosymbiosis, *Proceedings*
 848 *of the National Academy of Sciences*, 99, 11724-11729, 10.1073/pnas.172234799, 2002.
- 849 Zapata, M., Fraga, S., Rodríguez, F., and Garrido, J. L.: Pigment-based chloroplast types in
 850 dinoflagellates, *Marine Ecology Progress Series*, 465, 33-52, 2012.



851 Zhou, W., Li, Q. P., and Wu, Z.: Coastal phytoplankton responses to atmospheric deposition
852 during summer, *Limnol Oceanogr*, 66: 1298-1315, 2021.

853 Zhu, F., Massana, R., Not, F., Marie, D., and Vaultot, D.: Mapping of picoeucaryotes in marine
854 ecosystems with quantitative PCR of the 18S rRNA gene, *FEMS microbiology ecology*, 52,
855 79-92, 2005.

856

857



858 Tables and Figures

859 **Table1:** Initial conditions (t-12h) at the three stations sampled for the dust addition experiments. Other
 860 parameters are presented in more details in Gazeau et al. (2020; 2021)

Variables	TYR	ION	FAST
Location	Tyrrhenian Basin	Ionian Basin	Algerian Basin
Coordinates	39.34N, 12.60E	35.49N, 19.78E	37.95N, 2.90E
Temperatures (°C)	20.6	21.2	21.5
DOC (μM) ²	72.2	70.2	69.6
Chlorophyll <i>a</i> (μg L ⁻¹) ¹	0.063	0.066	0.072
BP (ng C L ⁻¹ h ⁻¹) ²	11.6	15.1	34.6
Bacterial abundance (x10 ⁵ cells mL ⁻¹) ¹	4.79	2.14	6.15
Viral abundance (x 10 ⁶ VLP mL ⁻¹)	3.01	1.44	2.79
% Lysogenic bacteria FLC	22.7	19.4	7.8
% Lytic bacteria FLIC	17.5	37.2	42.7
Viral production (x 10 ⁴ VLP mL ⁻¹ h ⁻¹)	2.05	1.36	7.99
HNF abundance (cells mL ⁻¹) ¹	110	53	126
Diatoms (cells L ⁻¹) ¹	340	900	1460
Dinoflagellates (cells L ⁻¹) ¹	2770	3000	3410
Ciliates (cells L ⁻¹) ¹	270	380	770

861

862 DOC: dissolved organic carbon, * BP: heterotrophic prokaryotic production, HNF: Heterotrophic
 863 nanoflagellates

864 ¹Results presented in Gazeau et al. 2020

865 ²Results presented in Gazeau et al. 2021

866

867



Table 2. Net growth rates (h^{-1}) calculated from the exponential phase of growth of BP, abundances of *Synechococcus* and picoeukaryotes cells, observable from at least three successive sampling points. Values $\pm$ standard error are shown, as well as the period of exponential phase (period, in days). nd: no significant exponential phase noted.

			C1	C2	D1	D2	G1	G2
TYR	$\mu_{\text{BP}}^{\text{app}}$	mean $\pm$ sd	0.076 ± 0.025	0.066 ± 0.018	0.116 ± 0.008	0.194 ± 0.02	0.164 ± 0.019	0.1503 ± 0.003
		period	0 - 0.5	0 - 0.5	0 - 0.5	0 - 0.5	0 - 0.5	0 - 0.5
TYR	$\mu_{\text{syn}}^{\text{app}}$	mean $\pm$ sd	nd	nd	nd	nd	0.014 ± 0.05	0.033 ± 0.003
		Period					2 - 3	2 - 3
TYR	$\mu_{\text{picoeuk}}^{\text{app}}$	mean $\pm$ sd	nd	nd	nd	nd	0.024 ± 0.004	nd
		period					2 - 3	
ION	$\mu_{\text{BP}}^{\text{app}}$	mean $\pm$ sd	0.042 ± 0.007	0.041 ± 0.005	0.09 ± 0.02	0.14 ± 0.006	0.13 ± 0.01	0.14 ± 0.03
		Period	0 - 0.5	0 - 0.5	0 - 0.5	0 - 0.5	0 - 0.5	0 - 0.5
ION	$\mu_{\text{syn}}^{\text{app}}$	mean $\pm$ sd	nd	nd	0.011 ± 0.001	0.015 ± 0.001	0.038 ± 0.002	0.045 ± 0.008
		Period			0.5 - 2	0.5 - 2	0.5 - 2	0.5 - 2
ION	$\mu_{\text{picoeuk}}^{\text{app}}$	mean $\pm$ sd	0.018 ± 0.001	0.012 ± 0.007	0.043 ± 0.014	0.034 ± 0.014	0.057 ± 0.012	0.053 ± 0.008
		Period	0.5 - 3	0.5 - 2	0.5 - 2	0.5 - 2	0.5 - 2	0.5 - 2
FAST	$\mu_{\text{BP}}^{\text{app}}$	mean $\pm$ sd	0.020 ± 0.002	0.026 ± 0.003	0.089 ± 0.014	0.090 ± 0.007	0.12 ± 0.005	0.16 ± 0.014
		Period	0 - 0.5	0 - 0.5	0 - 0.5	0 - 0.5	0 - 0.5	0 - 0.5
FAST	$\mu_{\text{syn}}^{\text{app}}$	mean $\pm$ sd	0.022 ± 0.002	0.024 ± 0.002	0.039 ± 0.001	0.045 ± 0.003	0.064 ± 0.001	0.063 ± 0.001
		Period	0.5 - 2	0.5 - 2	0.5 - 2	0.5 - 2	0.5 - 2	0.5 - 2
FST	$\mu_{\text{picoeuk}}^{\text{app}}$	mean $\pm$ sd	0.020 ± 0.002	0.012 ± 0.001	0.023 ± 0.004	0.026 ± 0.001	0.040 ± 0.002	0.034 ± 0.005
		Period	0.5 - 2	0.5 - 2	0.5 - 2	0.5 - 2	0.5 - 2	0.5 - 2



881 Figure legends:

882 **Figure 1.** Bacterial and viral parameters in the three experiments (TYR, ION and FAST) in each minicosm
 883 (D1, D2, G1 and G2). The values are normalized to the controls: the data are presented as the difference
 884 between the treatments and the mean value of the duplicate controls. The first row represents the
 885 bacterial cell specific growth rates and relative mortality rates at t24h after dust addition. The second
 886 row represents the relative viral productions at t24h and at T0 for the G treatments. The last row
 887 represents the viral strategies: the percentages of lytic (FLIC) or lysogenic (FLC) cells at t24h and at T0 for
 888 the G treatments.

889 **Figure 2.** (A) Log-log linear regression between bacterial biomass and bacterial production, dotted lines
 890 represent linear regressions for each treatment. (B) Relationships between log HNF abundance and log
 891 bacterial prey abundance. Solid black and dotted black lines corresponds to the Mean Realized HNF
 892 Abundance (MRA) and theoretical Maximum Attainable HNF Abundance line (MAA) respectively. The
 893 samples are grouped per treatments.

894 **Figure 3.** Relative abundance of viral populations at the initial (*in situ*: at t-12h before dust addition) and
 895 final time points in all minicosms (C1, C2, D1, D2, G1 and G2) during the three experiments (TYR, ION
 896 and FAST).

897 **Figure 4.** nMDS plot of bacterial community composition over the course of the three experiments
 898 based on Bray-Curtis dissimilarities of 16S rDNA sequences. Samples clustering at different level of
 899 similarity are circled together. All circles represent clusters which are significantly different from each
 900 other ($p < 0.05$) based on a PERMANOVA test.


901 **Figure 5.** nMDS plot of micro-eukaryotes community composition over the course of the three
 902 experiments based on Bray-Curtis dissimilarities of 18S rDNA sequences. Samples clustering at different
 903 level of similarity are circled together. All circles represent clusters which are significantly different ($p <$
 904 0.05) from each other based on a PERMANOVA test.

905

906




907



908

909 **Figure 1.**



910 **Figure 2.**

911

912




Figure 3.




Figure 4.




Figure 5.