



**HAL**  
open science

## Mass, velocity anisotropy, and pseudo phase-space density profiles of Abell 2142 (Corrigendum)

E. Munari, A. Biviano, G. A. Mamon

► **To cite this version:**

E. Munari, A. Biviano, G. A. Mamon. Mass, velocity anisotropy, and pseudo phase-space density profiles of Abell 2142 (Corrigendum). *Astronomy and Astrophysics - A&A*, EDP Sciences, 2015, 574, 10.1051/0004-6361/201322450e . insu-03644775

**HAL Id: insu-03644775**

**<https://hal-insu.archives-ouvertes.fr/insu-03644775>**

Submitted on 29 Apr 2022

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# Mass, velocity anisotropy, and pseudo phase-space density profiles of Abell 2142 (Corrigendum)

E. Munari<sup>1</sup>, A. Biviano<sup>2,3</sup>, and G. A. Mamon<sup>3</sup>

<sup>1</sup> Astronomy Unit, Department of Physics, University of Trieste, via Tiepolo 11, 34131 Trieste, Italy

e-mail: munari@oats.inaf.it

<sup>2</sup> INAF/Osservatorio Astronomico di Trieste, via Tiepolo 11, 34131 Trieste, Italy

e-mail: biviano@oats.inaf.it

<sup>3</sup> Institut d'Astrophysique de Paris (UMR 7095: CNRS & UPMC), 98bis Bd Arago, 75014 Paris, France

e-mail: gam@iap.fr


A&A 566, A68 (2014), DOI: 10.1051/0004-6361/201322450

**Key words.** methods: observational – galaxies: clusters: general – galaxies: kinematics and dynamics – galaxies: general – errata, addenda

The pseudo phase-space density profile of Abell 2142, defined with the galaxy number density profile  $\nu$  instead of the cluster mass density profile  $\rho$ , was shown in Fig. 12 of Munari et al. (2014). This graph was erroneous (because we had incorrectly considered the projected number density profile instead of the 3D profile). The correct figure is shown below. The values in Table 4 are changed, and the correct ones are reported in the table below.

While Fig. 12 of Munari et al. (2014) indicated that the PPSDs computed with the number density profile are significantly shallower than the theoretical relation of Dehnen & McLaughlin (2005), the corrected version of the figure shown here indicates that the PPSDs computed with the number density profile are now either consistent with the relation of Dehnen & McLaughlin ( $Q(r)$  for BLUE sample) or only slightly shallower, but not less consistent with that relation than found for the PPSDs computed with the mass density profile.

Therefore, the statement in Munari et al. (2014) that the mass density profile represents the PPSD and  $\beta - \gamma$  relations better is no longer correct. Indeed, Figs. 11 and 13 of Munari et al. show that the  $\beta - \gamma$  relations obtained using the mass density or tracer number density are indistinguishable for the RED and ALL samples. And therefore, the entire discussion of the greater relevance of the mass density profile relative to the galaxy number density profile must be dismissed.



**Fig. 12.** Same as Fig. 10 of Munari et al. (2014), but now using the radial profiles of galaxy number density instead of total mass density to estimate the PPSD.

## References

- Dehnen, W., & McLaughlin, D. E. 2005, MNRAS, 363, 1057  
 Munari, E., Biviano, A., & Mamon, G. A. 2014, A&A, 566, A68

**Table 4.** Best-fit parameters of the PPSD profile.

		$Q(r)$		$Q_r(r)$	
	A	B	A	B	
	$[M_\odot \text{ Mpc}^{-3} \text{ km}^{-3} \text{ s}^3]$		$[M_\odot \text{ Mpc}^{-3} \text{ km}^{-3} \text{ s}^3]$		
Fixed slope					
ALL	$5534 \pm 314$	-1.84	$25\,071 \pm 3341$	-1.92	
RED	$7727 \pm 391$	-1.84	$38\,484 \pm 5622$	-1.92	
BLUE	$1753 \pm 294$	-1.84	$3998 \pm 1084$	-1.92	
Free slope					
ALL	$6342 \pm 367$	$-2.28 \pm 0.11$	$29\,175 \pm 4223$	$-2.27 \pm 0.24$	
RED	$8034 \pm 411$	$-2.00 \pm 0.09$	$38\,881 \pm 5665$	$-1.77 \pm 0.23$	
BLUE	$3121 \pm 793$	$-2.97 \pm 0.50$	$5413 \pm 1810$	$-2.60 \pm 0.67$	
		$Q(r)$ GAL		$Q_r(r)$ GAL	
	A	B	A	B	
	$[10^{-9} \text{ Mpc}^{-3} \text{ km}^{-3} \text{ s}^3]$		$[10^{-9} \text{ Mpc}^{-3} \text{ km}^{-3} \text{ s}^3]$		
Fixed slope					
ALL	$3.7 \pm 0.18$	-1.84	$17. \pm 2.2$	-1.92	
RED	$2.9 \pm 0.14$	-1.84	$13. \pm 1.9$	-1.92	
BLUE	$0.37 \pm 0.056$	-1.84	$0.68 \pm 0.18$	-1.92	
Free slope					
ALL	$3.6 \pm 0.86$	$-1.72 \pm 0.10$	$17. \pm 2.4$	$-1.72 \pm 0.23$	
RED	$2.9 \pm 0.59$	$-1.75 \pm 0.09$	$14. \pm 2.0$	$-1.52 \pm 0.23$	
BLUE	$0.36 \pm 0.39$	$-1.74 \pm 0.48$	$0.62 \pm 0.21$	$-1.39 \pm 0.66$	

**Notes.** The PPSD profile is parametrized as  $Q(r) = A r^B$ . The first panel at the top shows the results of the fit of  $Q(r)$  and  $Q_r(r)$  for the different samples, both when keeping the exponent fixed to the values suggested by [Dehnen & McLaughlin \(2005\)](#), and when considering the exponent as a free parameter. In the bottom panel (the one identified by  $Q(r)$  GAL and  $Q_r(r)$  GAL) the same quantities are shown, but they refer to the PPSD computed using the galaxy number density profile instead of the total matter density profile.