



HAL
open science

Plan β : core or cusp?

Thomas D. Richardson, Douglas Spolyar, Matthew D. Lehnert

► **To cite this version:**

Thomas D. Richardson, Douglas Spolyar, Matthew D. Lehnert. Plan β : core or cusp?. Monthly Notices of the Royal Astronomical Society, 2014, 440, pp.1680-1689. 10.1093/mnras/stu383. insu-03645653

HAL Id: insu-03645653

<https://insu.hal.science/insu-03645653>

Submitted on 25 Apr 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Plan β : core or cusp?

Thomas D. Richardson,¹★ Douglas Spolyar^{2,3} and Matthew D. Lehnert²

¹*Department of Physics, Kings College London, Strand, London WC2R 2LS, UK*

²*Institut d'Astrophysique de Paris, 98 bis boulevard Arago, F-75014 Paris, France*

³*GRAPPA Institute, University of Amsterdam, Science Park 904, NL-1090 GL Amsterdam, the Netherlands*

Accepted 2014 February 25. Received 2014 February 24; in original form 2013 November 20

ABSTRACT

The inner profile of dark matter (DM) haloes remains one of the central problems in small-scale cosmology. At present, the problem cannot be resolved in dwarf spheroidal galaxies due to a degeneracy between the DM profile and the velocity anisotropy β of the stellar population. We discuss a method which can break the degeneracy by exploiting 3D positions and 1D line-of-sight (LOS) velocities. With the full 3D spatial information, we can determine precisely what fraction of each stars LOS motion is in the radial and tangential direction. This enables us to infer the anisotropy parameter β directly from the data. The method is particularly effective if the galaxy is highly anisotropic. Finally, we argue that such a test could be applied to Sagittarius and potentially other dwarfs with RR Lyrae providing the necessary depth information.

Key words: galaxies: dwarf – galaxies: kinematics and dynamics – Local Group – dark matter.

1 INTRODUCTION

A central prediction of cold dark matter (CDM) simulations (Navarro, Frenk & White 1996) is the formation of a cusp at the centre of a DM halo. Mounting (though still debated) observational evidence (see Gentile et al. 2004 for spiral galaxies) contrarily prefers a cored halo. This issue has been dubbed the ‘cusp versus core problem’. If indeed haloes are cored, then our present understanding of CDM haloes requires modification from either astrophysical effects or new particle physics. On the astrophysical side, numerous explanations (Read & Gilmore 2005; Tonini, Lapi & Salucci 2006; Governato et al. 2012) have been given that may reduce the cusp of a halo. On the other hand, if DM has an internal force, then the self-interaction (Spergel & Steinhardt 2000) can induce a core (Zavala, Vogelsberger & Walker 2013).

At a practical level, the DM profile is also an important uncertainty for indirect detection experiments. Dwarf galaxies are a primary target for indirect experiments, e.g. the *FERMI* gamma-ray telescope (Ackermann et al. 2011) looking for a DM annihilation signal. The potential signal is highly sensitive to the DM profile with the annihilation rate going like the density squared $\sim \rho^2$. Reducing this astrophysical uncertainty will improve indirect detection constraints on the particle physics properties of DM. Hence, determining the density profile of a dwarf galaxy is a critical benchmark in deciphering the nature of DM and structure formation in general.

Unfortunately, the density of DM at the centre of distant spheroidal systems is very poorly constrained by the simplest and most widely used method in galactic dynamics – the Jeans equation. The failure of this method to address the cusp/core issue goes some way

to explaining why there is still no consensus. The fundamental limitation of the Jeans equation is a degeneracy between the density profile and the β parameter which measures the anisotropy between radial and tangential velocity dispersions. A direct measurement of this quantity is not possible if the observer only has access to the line-of-sight (LOS) component of each stars velocity and a 2D projection of each stars radius. This is the kinematic data that is available for Milky Way dwarf galaxies at present. As a shorthand, we call it the 2+1 scenario as we have access to two of the three position coordinates and one of the three velocity coordinates for each star in the sample.

In practice, the flat LOS velocity dispersion profiles found in real (2+1) dwarf spheroidal data sets are fit equally well (Dejonghe & Merritt 1992; Charbonnier et al. 2011) by solutions to the Jeans equation encompassing a huge range of density profiles simply by tuning the anisotropy parameter. More precisely, Wolf et al. (2010) has recently shown that mass estimates of dwarf spheroidal galaxies are approximately independent of the anisotropy parameter β at the radius of half-light. If β does not undergo sharp transitions within the stellar extent then an inference of the mass at this radius is primarily limited by statistical noise in the velocity measurement. At smaller and larger radii, a free choice of β permits a wider range of masses that can fit velocity dispersion measurements. Unless one has some a priori intuition for the anisotropy parameter, the mass-anisotropy degeneracy masks the density slope of DM at the galactic centre. Under such circumstances, even a broad measurement of the anisotropy parameter (e.g. ruling out $\beta < 0.2$) could offer a significant improvement.

There have been several interesting attempts to mitigate the above uncertainty for 2+1 data sets. Here, we provide a brief summary of the simplest analytic methods and guide the interested reader to Battaglia, Helmi & Breddels (2013) for a detailed review of

* E-mail: thomas.d.richardson@kcl.ac.uk

Table 1. Summary of observational scenarios: X+Y indicates that we can measure X position coordinates and Y velocity coordinates.

Method	Description
2+1	Only the projected radius $R^2 = x^2 + y^2$ and LOS velocity v_z is available for each star (standard technique). Large data sets (500+ stars) are currently available for the classical Milky Way dwarf spheroidal population. See the Introduction for details of techniques that use 2+1 data.
2+2(3)	Proper motions of each star are added to R (and v_z). As discussed in An, Evans & Deason (2012), the <i>Gaia</i> satellite's spectrograph has a limiting magnitude (in the G band) of $G \sim 17$ whilst astrometric measurements with microarcsecond precision can be made for stars with $G \sim 20$. The <i>Gaia</i> satellite will therefore be able to measure the proper motion of stars before LOS velocities are available in systems such as Galactic globular clusters (at a distance of ~ 50 kpc or less). Mass estimators for this 2+2 scenario are provided in An et al. (2012). See also Wilkinson et al. (2002) and Strigari, Bullock & Kaplinghat (2007) for dynamic techniques that can be applied to 2+3 data. In these two works, the authors consider the scenario where proper motions can be added to existing LOS velocity measurements in dwarf spheroidal galaxies.
3+1	In this scenario, the LOS depth z is available before the proper motions. This could arise if the variable nature of a star is used to determine its distance from the observer. The LOS depth can be added to R to calculate the deprojected 3D radius $r^2 = R^2 + z^2$. This is the situation that we investigate in unprecedented detail in this paper.
3+3	The full 6D information of each star is known. By performing a simple coordinate transformation, one has the radial and tangential velocities. The anisotropy parameter $\beta(r)$ can then be read off directly.

the more sophisticated numerical techniques. With multiple stellar populations, one has multiple half-light radii for which one can estimate the mass. Given mass estimates at multiple points, the mass gradient (Walker & Penarrubia 2011) can be calculated. Similarly, in the case of Sculptor, Agnello & Evans (2012) have also exploited multiple stellar populations to place constraints on cusped profiles with the Viral theorem.

Without using multiple populations, higher moments of the LOS velocity distribution (Łokas, Mamon & Prada 2005) can also break the degeneracy between β and the DM density profile. A complete generalization of the classic Jeans equation analysis has very recently been derived in Richardson & Fairbairn (2013). The kurtosis (fourth moment divided by the dispersion squared) of LOS velocity data is very sensitive (Łokas 2002) to the anisotropy parameter. By simultaneously fitting to the kurtosis and the velocity dispersions of the data we have an additional constraint on the density profile. This has become an increasingly attractive option as dwarf (2+1) data sets have grown. Recent sets of 2+1 kinematic data (Walker, Mateo & Olszewski 2009) for Sculptor and Fornax still exceed 1000 stars after applying sophisticated interloper removal schemes.

In order to obtain maximum information from discrete data however, one would like to avoid the binning process altogether and to instead evaluate the likelihood of the fit on a star-by-star basis with the full phase space distribution function $f(\mathbf{x}, \mathbf{v})$ (see Chakrabarty & Portegies Zwart 2004 for equations that relate the distribution function to observables in numerous X+Y scenarios). In practice, marginalizing over all possible distribution functions rather than just fitting to the velocity moments with the Jeans equations presents significant new technical challenges but by utilizing Jeans's theorem (i.e. using integrals of motion such as specific energy E and angular momentum L rather than \mathbf{x} and \mathbf{v} as coordinates for f) the distribution functions are implicitly guaranteed to be in dynamic equilibrium which is particularly convenient when generalizing to non-spherical systems. For a more detailed discussion of state-of-the-art discrete modelling techniques, we refer the reader to Magorrian (2014).

Clearly, if we had all 6D phase space information for stars in a galaxy, we could directly measure the velocity anisotropy $\beta(r)$ with a simple coordinate transform. The degeneracy is broken and an inference of the density profile is only limited by statistical and experimental noise. Short of having all 6D information, as astronomical experiments become ever more sophisticated one might ask what information we need to tackle this degeneracy head on. We summarize the possible observational scenarios in Table 1 and use the notation X+Y to indicate the scenario in which the observer can measure X components of each stars position and Y components of each stars velocity. Estimators for the (global) anisotropy (Leonard & Merritt 1989; Genzel et al. 2000) and the mass (An et al. 2012) have been developed for 2+3 data sets with projected radii and the full 3D velocity information. Indeed, it has been shown (Wilkinson et al. 2002; Strigari et al. 2007) that the proper motions enable a precise measurement of the density slope (not just the mass) at the half-light radius. Dwarf galaxies are unfortunately sufficiently far away that it will be challenging to get proper motion of stars within a dwarf galaxy in the near future. Proper motions for stars in Galactic globular clusters however may soon be within reach with the *Gaia* satellite (see Table 1 for details) and this 2+2 scenario is explored in An et al. (2012).

Alternatively, if we know the depth of stars in the galaxy and LOS motion, we can break the degeneracy between β and the slope of the DM profile. Depending upon where a star sits in a galaxy, the motion will primarily be either in the tangential or radial direction. Heuristically, from the stars moving in the radial direction, we can compute the dispersion in the radial direction, and similarly, from stars moving in the tangential direction, we can determine the tangential dispersion. In the case of the Andromeda galaxy, Watkins, Evans & An (2010) used 500 satellite galaxies to infer β given positions and LOS velocities. We expand upon the technique and apply the more general method to dwarf galaxies.

In the rest of the paper, we focus on the 3+1 method. In Section 2, we exploit the 3D information to give an intuitive understanding of how β is revealed in the LOS velocity data. As a proof of concept, we then explicitly show how we can isolate β from the density

parameters and thus break the mass-anisotropy degeneracy in a likelihood analysis of dwarf spheroidal data. At the end of the section, we present very simple estimators for β in the form of sample variances.

In Section 3, the performance of these estimators is tested on mock dwarf spheroidal data from the *Gaia* challenge. To simulate a population of variable stars in real dwarf spheroidal galaxies we limit our sample size to 500 stars. We show that a sample of 500 stars is not sufficient to precisely determine the local value of $\beta(r)$ but may be used rule out large regions of the parameter space. We also discuss the sensitivity of the estimator to biases from the experimental errors and the assumption of spherical symmetry. Finally, we turn to the feasibility of our technique. We find that our technique can likely work for Sagittarius. If observational techniques can be improved by a factor of a few, our technique may also be applicable to other nearby dwarfs.

2 DYNAMICS WITH 3D POSITIONAL INFORMATION

In this section, we show how access to each stars depth along the LOS reveals β which parametrizes the anisotropy between radial and tangential velocity dispersions. We show how the LOS depth measurements can be used to tackle the mass-anisotropy degeneracy and derive estimators of β that can be applied to dwarf data sets with 3D positions and LOS velocities.

2.1 The imprint of velocity anisotropy

Let us define the origin of our spherical coordinate system to be at the galaxy's centre such that the observer is located a distance d_c in the negative z -direction. Assuming that d_c is very large relative to the system's scale (as is the case in dwarf spheroidal galaxies), then we may approximate the distance from the observer to any given star to be $d \simeq d_c + z$. The projected 2D radius $R^2 = x^2 + y^2$ that is measured on the sky can be combined with z to give the de-projected 3D radius $r^2 = x^2 + y^2 + z^2$. The LOS velocity of each star is

$$v_z = v_r \cos \theta - v_\theta \sin \theta, \quad (1)$$

where θ is the angle between the star's position vector and the z -axis. Under the assumption of spherical symmetry there is a uniform probability in solid angle $P(\Omega) = 1/4\pi$ and the density of stars $\nu(r)$ depends only on the radius. If the stars belong to an anisotropic distribution function $f(r, \mathbf{v})$, then the LOS velocity dispersion at each (deprojected 3D) radius is

$$\sigma_z^2(r) = \frac{1}{v} \int \frac{d\Omega}{4\pi} \int d^3\mathbf{v} (v_r \cos \theta - v_\theta \sin \theta)^2 f(r, \mathbf{v}). \quad (2)$$

Performing first the integration over velocity space, we use the standard definition,

$$\nu \sigma_k^2(r) = \int d^3\mathbf{v} v_k^2 f(r, \mathbf{v}), \quad (3)$$

for the velocity dispersions of $f(r, \mathbf{v})$, where $k = r, \theta, \phi$. Subsequently integrating over the azimuthal angle ϕ and introducing the anisotropy parameter, $\beta = 1 - \sigma_\theta^2/\sigma_r^2$ we find

$$\sigma_z^2(r, \theta) = \sigma_r^2(\cos^2 \theta + (1 - \beta) \sin^2 \theta). \quad (4)$$

This result has been derived previously in Leonard & Merritt (1989).

Equation (1) clearly shows that the motion observed along the LOS depends upon both the radial and tangential velocity. The




Figure 1. The visual imprint of velocity anisotropy. For systems with isotropic (upper panel), tangentially biased (middle panel) and radially biased (bottom panel) orbits, the LOS velocities of 5000 stars are plotted as a function of the standard polar angle θ between the position vector of each star and the LOS axis. Dashed red lines show the relative contributions of the radial and tangential dispersions to the LOS dispersion σ_z (solid red) as described by equation (4). Black vertical lines show the boundaries of regions A and B that are described in Section 2.2.

relative contribution from each component will depend upon the angle θ and changes sinusoidally.

We can visualize the effect. Fig. 1 plots the LOS velocity of 5000 stars as a function of θ for a fixed radius r . The radial velocity dispersion has been set to $\sigma_r^2 = 1 \text{ km s}^{-1}$. The velocity anisotropy parametrizes a trigonometric envelope which outlines the dispersion of the stars as a function of angle. The shape of this envelope is shown in Fig. 1 as solid red lines. Dashed red lines show the contribution to the LOS variance from σ_r^2 and σ_θ^2 . At the centre ($\theta = \pi/2$) of Fig. 1 the LOS velocities have no radial velocity

component and $\sigma_z = \sigma_\theta$. At the edge of the figure ($\theta = 0, \pi$) the position vector of each star is aligned with the LOS axis and $\sigma_z = \sigma_r$.

With 5000 stars the impact of β is visible by eye. In the isotropic case ($\beta = 0$), the variance is the same as a function of angle since $\sigma_r = \sigma_\theta$ and the amplitude of the sine and cosine functions are the same. In the tangentially biased case $\beta = -2$, the variance has a bulge at $\theta = \pi/2$ since $\sigma_r < \sigma_\theta$. Conversely, in the radial biased case $\beta = 2/3$, the variance increases at the edges since $\sigma_r > \sigma_\theta$.

It is also clear from Fig. 1 that if we fix σ_r (i.e. we fix the solid red lines at the edges of the figure), then the global average of σ_z^2 over all angles is determined uniquely by β . The global average of σ_z^2 increases if β is decreased. Averaging equation (4) over θ gives

$$\sigma_z^2(r) = \left(1 - \frac{2\beta}{3}\right) \sigma_r^2 \quad (5)$$

as was previously derived by Leonard & Merritt (1989).

2.2 Breaking the mass-anisotropy degeneracy

In studying dwarf galaxies, our primary aim is to constrain the density profile of the dwarf's DM halo. Let us say that we observe a dwarf galaxy and our data set comprises of the 3D positions and LOS velocities of stars. We would like to evaluate the likelihood of the data given a model of the density profile. Typically, this is achieved by invoking the assumption of dynamic equilibrium and using the Jeans equation to solve for the variance of the LOS velocity distribution. Unfortunately, this is not possible unless we specify the anisotropy parameter $\beta(r)$ in addition to the DM density profile and the density of stars $\nu(r)$.

Famously there is a degeneracy between the mass and anisotropy parameter if we use the Jeans equation (and subsequent projection for $\sigma_z^2(R)$) to fit LOS dispersion data that is flat as a function of the projected radius R . A similar degeneracy persists if we have all three positional components. To break the mass-anisotropy degeneracy with the new LOS depth measurement we must do more work than simply binning the LOS velocity data in spherical radial bins (as a function of the 3D radius r) rather than the usual cylindrical ones.

Perhaps the simplest and most intuitive way to disentangle the β and density parameters is to split the data set into two angular sub-regions (A and B) of equal stellar content. In a spherically symmetric gravitational potential, the stars at every angular position have the same contribution from the density parameters. As discussed in Section 2.1, the angular position does however effect the relative contributions of the radial and tangential velocities. We can thus use the angular information to isolate β and break the degeneracy.

From Fig. 1, we see that to maximize the impact of β then we should choose region A to be centred at $\pi/2$ and B to enclose the regions nearest the LOS axis in the positive and negative directions. To ensure an equal number of stars in each region (and noting that the stars are distributed uniformly in $\cos \theta$) we define the boundaries at $|\cos \theta| = 1/2$. Averaging the LOS dispersion (equation 4) over all angles θ in each sub-region, we find

$$\langle \sigma_z^2 \rangle_A = \left(1 - \frac{5\beta}{12}\right) \sigma_r^2, \quad A : |\cos \theta| < \frac{1}{2}, \quad (6)$$

$$\langle \sigma_z^2 \rangle_B = \left(1 - \frac{11\beta}{12}\right) \sigma_r^2, \quad B : |\cos \theta| > \frac{1}{2}. \quad (7)$$

The ratio of the above equations depends only on β . We have removed the dependence on the density parameters. By fitting to the velocity dispersion in two angular regions, A and B, we can

then solve for β and break the degeneracy with the DM profile of the halo.

In order to test our method, we create a mock dwarf galaxy data set. To model galaxies we often assume that the positions $\{\mathbf{x}_i\}$ and velocities $\{v_i\}$ of each tracer star are random samples from a smooth 6D distribution function $f(\mathbf{x}, v)$. We can then parametrize a galaxy by defining the distribution of tracer stars f and the DM density profile ρ_{DM} .

For our purposes, here we are only interested in measuring the LOS velocity dispersions. First, we assume spherical symmetry. Secondly, we also assume for simplicity that the LOS velocity distribution is Gaussian (this assumption will be relaxed in Section 3). With these assumptions, we need only consider the spatial density of stars $\nu(r)$ and the variances of the radial and tangential velocity distributions. As discussed previously these can be determined from the Jeans equation if we specify the DM density profile and the anisotropy parameter β . The LOS dispersion then follows from equation (5). We therefore define¹ a ‘model’ for a dwarf galaxy by its spatial density of stars $\nu(r)$, its DM density profile ρ_{DM} and its anisotropy parameter β . Every ‘mock galaxy’ discussed in this section is a Monte Carlo sampling of 5000 stars from a parent model.

We generated a mock galaxy of 5000 stars, galaxy O (for cOre) from parent model O. To generate each LOS velocity, we solved the Jeans equation and used equation (4) as the variance of a Gaussian LOS velocity distribution. The density of stars $\nu(r)$ in model O is parametrized by a Plummer profile and has a characteristic radius of 250 pc. The DM distribution of model O is (as the name suggests) a cored (inner density slope = 0) Hernquist density profile. The anisotropy parameter of model O rises from $\beta = 0$ at the galactic centre to $\beta = 1/2$ at large radii.

Let us say that we are presented with the mock data set galaxy O but do not know that the parent model is the anisotropic and cored model O. We want to show the power of the 3+1 method to discriminate between models that are degenerate in the standard 2+1 (no LOS depth measurement) Jeans equation analysis. We therefore introduce model U (for cUsp). Model U has an identical Plummer profile $\nu(r)$ to model O. The DM profile of model U is a cusped Navarro-Frenk-White (NFW; Navarro, Frenk & White 1997) profile and the anisotropy parameter is $\beta(r) = 0$.

In the top panel of Fig. 2, we assess the standard (2+1) method on galaxy O and demonstrate the mass-anisotropy degeneracy. In this case, we only have access the projected radii R and LOS velocities of the 5000 stars in galaxy O. The data points in the upper panel show the LOS velocity dispersion (v_z^2) of galaxy O's stars in 10 cylindrical radial bins. The solid red and dashed black lines show the variance of the LOS velocity distribution σ_z^2 in models O and U, respectively. Despite the very different DM density profiles of models O and U we see by eye that by suitable tuning of the anisotropy parameter β the LOS variances are very similar.

To assess the statistical significance of this claim for galaxy O, we must compute the likelihood. We can perform a frequentist estimate for the likelihood of a model by generating many mock galaxies from the model parameters and determining the regions where the median and central 67 and 95 per cent of our estimators (in this case $\langle v_z^2 \rangle$) lay. The blue region shows the resulting likelihood contours for model U. Clearly with only 5000 stars the width of the contours is significantly larger than the difference between the two models.

¹ To be clear our use of ‘model’ is not a Bayesian model that defines a parameter space but rather one parameter set of a Bayesian model.




Figure 2. For a mock dwarf galaxy of 5000 stars, galaxy O, we compare an isotropic ($\beta = 0$) model with a cusped density profile (model U) and an anisotropic model with a cored density profile (model O). Galaxy O is in fact one Monte Carlo sampling from model O (solid and dotted red lines). We show a fit to the data with model U (dashed black lines). Blue lines and shaded regions show the median and 67 and 95 percent contours for the likelihood of the galaxy O data given the parameters of model U. See text for details. Upper panel: here, we assume that the observer only has access to the 2D projected radius R and the LOS velocities v_z . Data points show the sample variance $\langle v_z^2 \rangle$ of galaxy O’s LOS velocities in 10 cylindrical radial bins. Solid red and black dashed lines show the Jeans equation prediction for the LOS dispersion after marginalizing over the LOS depth (see equations 4–57 in Binney & Tremaine 1987). Lower panel: now, we assume that the observer additionally has the LOS depth measurements of the mock dwarf data. The data may now be divided into two angular sub-regions A and B according to the condition $|\cos \theta| < 1/2$. Each half of the galaxy O data set is then split into 10 spherical radial bins. Green circular and magenta triangular data points show the sample variance of LOS velocities in regions A and B. Red solid and dotted lines show the expected values of the LOS dispersion in each angular region according to equations 6 and 7. We note that that as model U is isotropic then $\langle \sigma^2 \rangle_A = \langle \sigma^2 \rangle_B$. For clarity, we therefore only show the likelihood contour for the $\langle v_z^2 \rangle_A$ data.

Though we do not explicitly show the likelihood given the cored model O, it is evident that the likelihood of the data given our cusped isotropic model is very similar to that of our cored model with tuned anisotropy. We are thus unable to reject either hypothesis.

With the 3+1 method we can break the degeneracy. The lower panel of Fig. 2 shows what we see if we additionally measure the LOS depth of each star. By splitting galaxy O into angular sub-regions A and B, we see that the LOS velocity dispersions in each region (region A – green circular and region B – magenta triangular) are visibly divided at large radii. This splitting of dispersions in region A and B is a smoking gun for anisotropy $\beta \neq 0$ as can be seen in Fig. 1. In particular, we see from the bottom panel of Fig. 1 that if the LOS dispersion in region A is larger than in region B, then this indicates radial anisotropy $\beta > 0$. The solid and dotted red lines show the variance of the LOS velocity distribution in each region for the anisotropic model O. Conversely, the model U will

have the same LOS velocity distribution in both angular regions which is shown with a black dashed line.

Hence, model U cannot simultaneously fit the LOS dispersion data of galaxy O in angular regions A and B. The degeneracy between the two models is broken. The blue likelihood contours (in this case the estimator is $\langle v_z^2 \rangle_A$) for model U show that this is a statistically significant result for our mock galaxy of 5000 stars.

Of course 5000 stars is an unrealistically large sample size that is chosen for illustrative purposes. We note in Fig. 2 that we may also have ruled out tangentially anisotropic models (where $\langle \sigma_z^2 \rangle_A$ is instead smaller than $\langle \sigma_z^2 \rangle_B$) with an even greater significance. Even for modest sample sizes where a precise measurement of the splitting is obscured by noise, it should still be possible to place a strong upper or lower bound.

2.3 Estimators for β

In the previous section, we described how one could directly tackle the degeneracy problem. For a real galaxy, it may often be the case that only a small sub-set of available stars are amenable to a precise measurement of the LOS depth. Additionally, this measurement will be subject to a myriad of potential uncertainties that introduce complicated systematic biases. In either case, it may therefore be preferable to use the limited 3D data set to make a more robust estimate β and to use this as a prior in existing 2+1 methods that treat larger and more precisely measured 2D data sets. In this section, we introduce simple estimators for β .

First let us adapt the method applied in Section 2.2. Eliminating the σ_r^2 term we derive the estimator for β ,

$$\hat{\beta}_0 = 12 \frac{\langle v_z^2 \rangle_B - \langle v_z^2 \rangle_A}{11 \langle v_z^2 \rangle_B - 5 \langle v_z^2 \rangle_A} \quad (8)$$

that approximates β for a radial bin of stars. Though useful as an illustrative example this method requires a splitting of the data that increases the statistical noise by a factor of approximately $\sqrt{2}$. We can instead obtain a new constraint by reweighting the LOS velocities by $\cos^2 \theta$ and again performing the average over solid angles,

$$\overline{\sigma_z^2 \cos^2 \theta} = \int \frac{d\Omega}{4\pi} (\sigma_r^2 \cos^4 \theta + \sigma_\theta^2 \sin^2 \theta \cos^2 \theta) \quad (9)$$

$$= \frac{1}{3} \left(1 - \frac{2\beta}{5} \right) \sigma_r^2. \quad (10)$$

Again we may cancel out σ_r^2 to get a new estimator,

$$\hat{\beta}_1 = \frac{3}{2} - \frac{1}{5\omega - 1}, \quad \omega \equiv \frac{\langle v_z^2 \cos^2 \theta \rangle}{\langle v_z^2 \rangle} \quad (11)$$

that uses the full data set.

As an aside, one could also perform a maximum likelihood analysis on the individual data points to derive β but we found that introducing the angular dependence strongly altered the simple Gaussian velocity distributions. Numerical tests indicated that the maximum likelihood method is more susceptible to the complicated biases that are discussed in the next section.

3 MONTE CARLO PERFORMANCE OF β ESTIMATORS FOR MOCK DWARF DATA

In this section, we use mock dwarf spheroidal data from the *Gaia* challenge workshop (accessible on the wiki; Read & Kawata 2013)

to test the variance and bias of our anisotropy parameter estimators (equations 8 and 11). We draw stars from a more realistic distribution function $f(r, v)$ rather than the Gaussian approximation used in the preceding sections. As a point of comparison, the fake data sets have been drawn from two different but realistic models. Both models have identical Plummer stellar density profiles with a characteristic radius of 250 pc embedded in a DM halo with a scale radius of 1 kpc.

Model 1 has a cusped NFW density profile and an isotropic velocity distribution at all radii ($\beta = 0$). Model 2 has a cored DM halo and Osipkov–Merritt anisotropy where β rises from zero at the galactic centre to purely radial orbits ($\beta = 1$) at large radii. For more details on these models, we refer the reader to the *Gaia* Challenge data suite described in Walker & Penarrubia (2013). These two models were selected as they have very different anisotropy parameters but similar flat dispersion profiles.

Dwarf galaxies may have of the order of 10 thousand plus stars. In the feasibility section, we will focus on variable stars as our tracer population. Dwarf galaxies typically only have several hundred variable stars so we limit our sample size to $N = 500$.

A rigorous treatment of uncertainties in the distance measurement is a complex and method-dependent issue that is beyond the scope of this work. Indeed, studies [see Binney et al. (2014) for a treatment of stars in the RAVE survey (Steinmetz et al. 2006)] of how to quantify the error distribution of distances to stars are ongoing. As such we consider the performance of our estimator for a simplified model in which the errors for the distance measurements are assumed to be Gaussian before commenting on the impact of a more realistic error distribution.

To evaluate the variance and bias of our anisotropy estimators, we drew a large number of independent samples of $N = 500$ stars from the *Gaia* Challenge data set. To these positions and velocities, we added Gaussian distributed noise to the LOS depth z and velocity v_z measurement with standard deviations δ_z and δ_{v_z} and centred at the true coordinates. For each sample, we split the data into five radial bins of 100 stars, calculated the anisotropy estimator for each bin and added it to an array. In this way, we were able to construct likelihood regions for $\hat{\beta}$ in a similar fashion to those for $\langle v_z^2 \rangle$ in Fig. 2.

3.1 Sources of bias and an improved estimator $\hat{\beta}_{1b}$

For a robust inference of β and in particular to claim a stringent upper or lower bound, we must consider how experimental errors and other systematics bias our estimators. The experimental errors will introduce biases in our estimate for β .

Our numerical tests indicate that only substantially large LOS velocity errors $\delta_{v_z} \geq 8 \text{ km s}^{-1}$ can bias our estimators $\hat{\beta}$. Adding Gaussian distributed noise to the LOS velocities adds a systematic error to the LOS dispersion that is independent of the position of the star. In Fig. 1, we see that this would simply shift the red lines vertically. Because the estimators $\hat{\beta}$ are defined as ratios of the LOS dispersion measurements at different positions, this systematic effectively cancels out. In practice, velocity measurements should be sufficiently precise that we can neglect the bias due to velocity.

Instead, we focus upon the experimental errors in the LOS depth measurement, which will alter the distribution of stars first morphologically and secondly in terms of ordering. These two effects will introduce two separate biases.

Morphologically, experimental distance errors may artificially elongate the galaxy along the LOS. In our Gaussian example with width δ_z , the square average of the LOS depth measurement z

is transformed via $\langle z^2 \rangle \rightarrow \langle z^2 \rangle + \delta_z^2$. Measurements of x and y in plane perpendicular to the LOS are left unchanged however so we have $\langle z^2 \rangle > \langle x^2 \rangle = \langle y^2 \rangle$. In other words, our spherically symmetric halo becomes ellipsoidal. As our estimators are derived under the assumption of spherical symmetry (and particularly uniformity in solid angle) the elongation will bias the inferred value of β . For the interested reader, the bias is similar to the Lutz (Lutz & Kelker 1973) bias. In the Lutz case however, Gaussian errors in the parallax measurement ($p \propto 1/d$) have the effect of squashing the galaxy.

The depth measurement errors will introduce a second bias when we group the stars into radial bins. For sufficiently large scatter, the ordering of stars will be corrupted by moving stars from one bin to another, which introduces a separate ordering bias effecting $\hat{\beta}$. Though the two aforementioned biases can have a dramatic effect on $\hat{\beta}$, we now demonstrate that there are hints from the data that allow us to identify them and in some cases clean them.

First, we introduce a new estimator to remove the morphological bias for our estimate of β . We can identify the departure from spherical symmetry in our data set by measuring

$$b = \frac{1}{N} \frac{\sum_i \cos^2 \theta_i}{\langle \cos^2 \theta \rangle_{\text{true}}} = \frac{3}{N} \sum_i \cos^2 \theta_i. \quad (12)$$

A spherical system with uniform probability in solid angle has $\langle \cos^2 \theta \rangle = 1/3$. The parameter b indicates the fractional deviation from this prediction. By design, the parameter b enables us to formulate a simple estimator for the anisotropy parameter,

$$\hat{\beta}_{1b} = \frac{3}{2} - \frac{b}{5\omega - b} \quad (13)$$

for which we can correct for the morphological bias. In Fig. 3, we have plotted our original estimator $\hat{\beta}_1$ without the correction. We adopt Gaussian errors of $\delta_z = 100 \text{ pc}$ and ($\delta_{v_z} = 3 \text{ km s}^{-1}$) which are shown in Fig. 3. These errors are significant relative to the half-light radius $R_{1/2} = 250 \text{ pc}$ and the LOS dispersion $\sigma_z \approx 11 \text{ km s}^{-1}$.




Figure 3. 500 stars were sampled from a mock dwarf galaxy with an isotropic distribution function ($\beta = 0$ solid red line). Gaussian errors were added to the LOS depth and velocity to simulate experimental errors. Data points show the anisotropy parameter estimator $\hat{\beta}_{1b}$ (bias corrected) for five spherical radial bins of 100 stars. Shaded regions show the likelihood of the data points given the true isotropic model (blue) and a model with Osipkov–Merritt anisotropy (green) where β undergoes a transition from zero at the galactic centre to one at large radii above transition radius $r_a = 250 \text{ pc}$. Central lines show the median and shaded regions show the central 67 and 95 per cent intervals.

Clearly, the original estimator is offset from the true value of $\beta = 0$. Conversely, our morphologically corrected estimator $\hat{\beta}_{1b}$ now aligns with the true value of β . The solid blue line in Fig. 3 that represents the median value of $\hat{\beta}_{1b}$ for a large sample of isotropic mock data sets is approximately centred at zero. We found that if we increased the distance errors to much above $\delta_z = 150$ pc then even our improved estimator $\hat{\beta}_{1b}$ began to show significant bias.

Our corrected estimator assumes that all of the deviation from spherical symmetry is a result of the experimental errors. If the system is not spherically symmetric, we would need to introduce a different estimator with an alternative prediction for $\langle \cos^2 \theta \rangle$. This could be guided by evaluating the deviations from spherical symmetry that are present in the x and y positional data in the plane perpendicular to the LOS. In other words, we can check to see that the galaxy looks circular in projection as is the case in Draco. Any such analysis would vary from dwarf to dwarf and we leave this more complicated case for the future.

We can also predict the effects of the ordering bias, which mixes the stars from different radial bins. Though we measure the position of a star at a point with observed radius r_o , the true position of the star is actually at some other point with a true radius r_t . Given our limited knowledge, we will never know r_t . If we know the density of stars at each radius $\nu(r)$ and the distribution of errors in the depth measurement then we can use a likelihood analysis to determine the probable location, which will allow us mitigate the ordering bias. The 3D density of stars can be estimated by fitting to the surface density of stars that we observe on the sky and performing an Abel inversion.

Let us denote the error in the radial measurement α_r such that $r_o = r_t + \alpha_r$. If we know the distribution of LOS depth errors α_z , then we may use

$$\alpha_r = \sqrt{r_t^2 + r_t \cos \theta \alpha_z + \alpha_z^2} - r_t \quad (14)$$

to determine the distribution of radial measurement errors $P(\alpha_r|r_t)$. In our example where the LOS depth errors α_z are normally distributed we can easily derive the displacement error distribution function $P(\alpha_r|r_t)$ numerically by drawing α_z from the normal distribution and $\cos \theta$ uniformly from -1 to 1 . Even though we have a fixed error distribution for α_z we note that $P(\alpha_r|r_t)$ depends on the true radius r_t . We are now in a position to evaluate the expectation value of the true radius $\langle r_t \rangle$ at an observed radius r_o . This is simply a weighted combination of all r_t and α_r that combine to make the observed radius r_o , namely,

$$\begin{aligned} \langle r_t \rangle(r_o) &= \int_0^\infty dr N(r)r \int_{-\infty}^\infty d\alpha_r P(\alpha_r|r) \delta(r_o - r - \alpha_r) \\ &= \int_0^\infty dr N(r)r P(\alpha_r = r_o - r|r), \end{aligned} \quad (15)$$

where the delta function ensures that $r_o = r + \alpha_r$. Our β estimators depend on sample variances of the LOS velocities. With errors in the LOS depth we must update our estimates of the LOS dispersion to reflect the fact that the majority of stars at r_o originated from radii centred at $\langle r_t \rangle$. The impact of this bias depends on gradients in the LOS dispersion. If the gradients are large, then $\sigma_z^2(r_o)$ could be considerably different to $\sigma_z^2(\langle r_t \rangle)$. Fortunately, real dwarf spheroidals have approximately flat LOS dispersion profiles which minimizes this effect.

In summary, the estimators are prone to bias for any realistic account of the distance errors. Though this bias is not trivial to calculate, we have shown with our simple model that if we have an understanding of the error distributions and a good estimate

of the stellar density profile then there is sufficient information from the data to identify the bias and in some cases to clean it. We found that the estimators are insensitive to errors in the LOS velocity measurement but that even our improved estimator $\hat{\beta}_{1b}$ cannot account for bias if the error in the LOS depth measurement climbs above $\delta_z = 150$ pc.

In reality, the biggest issue that the method faces is that the estimator is sensitive to the assumption of spherical symmetry which is already in tension with observations of real dwarf spheroidals. Sadly, the only motivation for this assumption is simplicity. Nonetheless, the spherical model provides a useful test example to answer a more general question. If we can only observe the LOS component of each velocity, then how well can we utilize the full 3D positional information to determine the velocity anisotropy? For any configuration of stellar positions, the key to answering this question lies in determining the average contribution of each velocity coordinate to the LOS.

3.2 How well could we determine β in practice?

The primary concern of this section is the fundamental limitation of sample size. In practice, distance measurements of stars in dwarf spheroidal galaxies will be limited to variable stars. This will be discussed in greater detail in the feasibility section. We therefore consider small samples of 500 stars which represents the upper limit of variable star populations in dwarf spheroidals. We again adopt the benchmark errors of $\delta_{vz} = 3$ km s⁻¹ and $\delta_z = 100$ pc in the LOS velocity and depth. In the previous section, we showed that these errors are not sufficient to bias the estimator $\hat{\beta}_{1b}$.

From Fig. 3, we note that with 500 stars the blue likelihood contours corresponding to the isotropic galaxy are broad at all radii offering little distinction between data points in areas of mild radial ($\hat{\beta} > 0$) and tangential ($\hat{\beta} < 0$) anisotropy. As with the estimator $\hat{\beta}_0$, this reflects the fact that isotropy does not leave a strong signal on the angular distribution. By contrast, we see from the green contours that if the galaxy is strongly anisotropic, as is the case for the Osipkov–Merritt model at large radii, then the observed data points will almost always be clustered tightly about the true value. This is a key result. With a small sample of 500 tracers the strength of the 3+1 method depends on the nature of the galaxy that we observe.

If the galaxy that we observe is approximately isotropic, then clearly the statistical noise limits an inference of β . The scatter of data points will be consistent with a wide range of mildly anisotropic models and an inference is limited to setting upper and lower bounds that rule out more extreme models. Such information would still be of interest. Mass estimates at the half-light radius (Wolf et al. 2010) are less robust if the anisotropy parameter has large gradients. The mass slope method (Walker & Penarrubia 2011) in particular could be sensitive to such a correction if individual sub-populations have contrasting gradients. Though intuition suggests that large gradients in β are unlikely it would be useful to validate this assumption with observation. Any prior information on β could also be useful for methods that use higher moments (i.e. that measure the shape as well as the width) of the velocity distribution. As discussed in the introduction, these methods are much more sensitive to β than the classic Jeans analysis of velocity dispersions.

If the galaxy that we observe is instead highly anisotropic, then we can make even stronger assertions. Had we sampled our mock galaxy from the anisotropic Osipkov–Merritt model, then we see from the green contours in Fig. 3 that 95 per cent of measurements of $\hat{\beta}_{1b}$ at large radii ($r > 1000$ pc) will be constrained above $\hat{\beta}_{1b} = 0.5$.

The blue contours show that an observation of $\hat{\beta}_{1b} = 0.5$ is in tension with the isotropic model at the 2σ level. Of course, with one data point this is not highly significant but it can be used to place much more stringent constraints on tangential models with $\beta < 0$. If the model was highly anisotropic at all radii then five observations of $\hat{\beta}_{1b} = 0.5$ would be sufficient to exclude the isotropic model.

In this favourable scenario then, limiting the parameter space to positive β with the 3+1 method can break the degeneracy if combined with existing (two projected positions + one LOS velocity) methods. The dwarf that we observe will have a much larger (we no longer require that the stars are variable) complementary data set of 2D projected radii plus LOS velocities. The left-hand panel of fig. 1 in Wolf et al. (2010) shows the degenerate solutions of the Jeans equation that fit the (typically) flat LOS velocity dispersion of the Carina dSph. The mass at the half-light radius is fixed but the mass at the centre is masked by the mass-anisotropy degeneracy. In particular, we see that (in the case of constant anisotropy) radially anisotropic models with smaller interior masses (i.e. cores) are degenerate with tangentially anisotropic models with large interior masses (i.e. cusps). The prior of $\beta > 0$ set by our 3+1 method can thus break the degeneracy between cusps and cores.

4 FEASIBILITY

There are several techniques to determine the distance of individual stars in dwarf galaxies ranging from parallax to variable stars. We discuss how the different distance measurement techniques could be useful. See Table 1 for definitions of the different methods. We find that variable stars offer the best opportunity to apply the 3+1 method. First, we consider parallax, which turns out to be more important for methods other than the 3+1 method. With sufficiently good angular position and long enough baseline, we could get all 6D phase space information for a galaxy. Hence, one would use all the data and not be limited to a 3+1 method. Furthermore, even without sufficiently good parallax data to determine the depth of a star along the LOS, we could still get the proper motion of the stars inside the galaxy. The proper motion of stars in a dwarf plus the LOS velocity would allow one to use the 2+3 method discussed in the introduction. Hence, parallax would be more useful for other techniques.

In the near term, proper motion studies will be limited. To measure the internal motion of stars inside a dwarf will require an observe to determine the proper motion to the per cent level. *Gaia* in the next five years will map out the proper motion and position for over a billion stars and will significantly alter our understanding of the Milky Way. Regardless, *Gaia* will have difficulty to achieve the necessary precision (per cent level) to observe the internal motion of stars inside of dwarf galaxies, with the possible exception of Sagittarius.

Ground-based missions and pointing telescopes may be able to do better than *Gaia*. In the early 1950s, Baade & Swope (1961) made a detailed study of the variable stars in Draco. Over an entire season Baade took many long and deep exposures (100 plates plus) to determine the period of nearly 300 RR Lyrae. Subsequently, over the last 20 years, many telescopes including (CFHT, HST, KPNO, USNO, etc.) have imaged Draco and UMiI with CCDs. Following a technique discussed and used by Sohn, Anderson & van der Marel (2012) and Bouy et al. (2013), one should be able to use at least a 20 year baseline to determine the proper motion and position of the stars in Draco to the few per cent level. A longer base line (60 years) could reduce the error to the sub-per cent level. Hence, Draco and

Sagittarius are potentially interesting targets for the 2+3 method and may even allow for a full 6D analysis. Regardless, parallax data will be sparse and hard to come by in the near term.

Given the potential challenges of parallax, we look to other methods to determine the depth of a star along the LOS. Variable stars are very promising. Unlike proper motion studies, one does not need a long baseline to infer a distance. In addition, dwarf galaxies typically have several hundred variable stars such as RR Lyrae. Fornax has at least 500 RR Lyrae (Bersier & Wood 2002), Draco has nearly 300 RR Lyrae (Kinemuchi et al. 2008) if not more, and the Sagittarius dwarf has at least 400 stars (Soszynski et al. 2011). In the case of RR Lyrae stars, one can certainly determine the absolute distance of RR Lyrae stars to the per cent level using a period–luminosity relationship (PLR; Madore et al. 2013). Much of the error in the absolute distance is due to the constant offset term of the PLR. We are only interested in the relative depth of each star in the dwarf galaxy. Error in the offset term will not alter the relative depth of a star with respect to the centre of the galaxy. Hence, the 3+1 method will not be affected by errors related to the absolute distance relationship.

The fundamental limitation of the PLR for a dwarf spheroidals would be due to any intrinsic scatter. In the *V* band, this is certainly true (Sandage 1990). In the *K* band, theoretical models of the PLR (Bono et al. 2001; Catelan, Pritzl & Smith 2004) appear to have an intrinsic scatter of about 0.032 mag, which corresponds to about a 2 per cent $\sim \delta d/d$ uncertainty in relative distance (where d is the distance to the star and δd is the error in the distance). Bono et al. (2001) assumed an intrinsic mass scatter of about 4 per cent which accounts for most of the intrinsic scatter of the PLR by contributing 0.03 mag. If one can account for the mass, one could reduce any intrinsic scatter.

The *K*-band PLR depends upon metallicity. For simplicity, we have neglected any uncertainties in metallicity in the intrinsic scatter. First, we can directly measure the metallicity of RR Lyrae stars. Secondly, dwarfs such as Draco do not appear to not have a large range of metallicities. On the positive side, *Kepler* has brought new life into the field with new exquisite light curves for RR Lyrae (Kolenberg 2013). *Kepler*'s new insight has improved modelling of RR Lyrae, which may also have the side benefit of improving the PLR.

Regardless, we would like to observe directly any intrinsic scatter in the PLR. Unfortunately at present, observational errors mask any intrinsic scatter. Buckley, Longmore & Dixon (1992) looked at stars in M3 and ω Cen. The authors found a scatter of about 0.03 mag, which appears to be consistent with limited number of observations. If the scatter found by Catelan et al. (2004) and Bono et al. (2001) is real, then the mass and metallicity of stars in M3 and ω Cen are almost identical. Conversely, the PLR relationship may actually be less sensitive to the mass and metallicity of stars than has been found by Catelan et al. (2004) and Bono et al. (2001). In sum at present, the PLR relationship appears limited to the per cent error level in determining distances, but the relationship might still be improved in the future.

Beyond the PLR, the Baade–Wesselink (BW) method (Baade 1926; Wesselink 1946, 1969) may also be used to directly measure the distance to RR Lyrae in dwarf galaxies. The BW method measures the change in a star's luminosity as a function of the star's surface velocity and temperature, from which one can directly infer the intrinsic luminosity. The BW method is limited by how well we can measure the temperature of the star and the star's actual surface velocity.

Even without recent improvements in stellar modelling and various means to determine the temperature of a star such as with widths

of metal lines, the temperature of RR Lyrae stars have previously been determined down to 20 K with only *K*- and *V*-band photometry (see Cacciari, Clementini & Fernley 1992, for further references). Barcza (2010) and Barcza & Benkő (2012) have formulated a new way to directly model the size mass, radius and temperature of the star by exploiting multiple bands – no spectral information. The authors could determine the temperature to within 20 K and limit the mass to better than 4 per cent. We emphasize that the conclusions drawn by Barcza (2010) and Barcza & Benkő (2012) were made without the benefit of deep infrared band coverage (*H*, *J* and *K*) and without the use of any spectral information. In terms of surface velocity, we note that with the addition of spectral information, we can measure the expansion of a star down to a fraction of a km s^{-1} . Apogee can measure the velocity of stars down to 50 m s^{-1} (Deshpande et al. 2013). The stars in the dwarfs will be much fainter than the stars observed with apogee (mag 20 versus mag 13), but we can use a 10 metre telescope versus a 1 metre telescope, which will give a much faster integration time. Regardless, we probably can still measure the surface velocity with a precision on the order of few tenths of a km s^{-1} . With more probes (multiple bands and spectroscopy), we may well be able to improve our determination of the surface temperature, mass, and radius of RR Lyrae. More complimentary data cannot hurt.

As noted before, our numerical tests on mock dwarf spheroidal data indicate that the 3+1 method requires that we know the relative distance to better than 150 pc. We can translate that into a relative distance error of $\delta d/d$, where d is the distance to the star. We take $\delta d = 150 \text{ pc}$. For a fixed δd , we can then infer that the maximum error allowed us to apply the 3+1 method. For example, the maximum error allowed for Sagittarius is 0.7 per cent, 0.3 per cent for Ursa Minor, 0.2 per cent for Draco and 0.1 per cent for Fornax. In summary, we will need distances to a few tenths of a per cent.

A factor of a few improvement in the PLR and the BW method would be sufficient to apply the 3+1 method to the above-mentioned galaxies. As noted previously, PLR appears to only determine the relative distance down to the per cent level (multiple bands and measurement of the stars metallicity could improve the situation). As noted previously, any inherent dispersion appears is either limited to taking insufficient data or due to mass dispersion. If one can use multiple bands to better constrain the mass of RR Lyrae, then PLR maybe sufficient to apply the 3+1 method.

How well can the BW method work? The temperature of an RR Lyrae star is $T \sim 7000 \text{ K}$. The star has a surface velocity of $V \sim 150 \text{ km s}^{-1}$. As a base line, we will assume an error δV of 2 km s^{-1} and δT of 20 K, which gives a relative error on the distance $\delta d/d$ to around 0.7 per cent with $\delta d/d \simeq 2\delta T/T + \delta V/V$ (we have neglected uncertainties on the period, which is small). Most of the error comes from the temperature. Remarkably even without an improvement of BW method we could already successfully apply the 3+1 method to Sagittarius. A factor of a few improvement in the determination of temperature would allow us to also apply the 3+1 method to Ursa Minor, Draco and possibly Fornax. Finally, we emphasize that even with the present limitations on the BW method, the 3+1 method could be used to help constrain β in Sagittarius.

5 DISCUSSION

Without knowledge of the anisotropy parameter β , the standard methods in galactic dynamics are not able to unambiguously determine the DM density profile at the centre of the halo. By adjusting β , one can accommodate a cored or a cusped profile. For systems as distant as dwarf spheroidal galaxies this degeneracy cannot be

broken at present because the available LOS velocities and projected radii offer virtually no information on the anisotropy. With new sophisticated astronomical techniques this could be subject to change in the near future.

The scenario in which data sets are bolstered by the proper motions has been examined thoroughly in the literature. For observational reasons however, less attention has been paid to the case where we first gain access to the LOS depth of each star. In this work, we illustrate clearly how velocity anisotropy can be identified with this information and show for spherically symmetric systems how one could in theory break the mass-anisotropy degeneracy completely. For an application to small samples, we developed simple estimators for the anisotropy parameter $\hat{\beta}$ that could be used to inform priors on β in existing 2D methods. This would be particularly effective for methods that employ higher moments of the velocity distribution which are sensitive to the anisotropy parameter.

To gauge how our estimators might perform in practice we tested them on realistic mock dwarf data from the *Gaia* Challenge. Samples of $N = 500$ stars were given Gaussian experimental errors of $\delta_z = 100 \text{ pc}$ in the LOS depth and $\delta_{v_z} = 3 \text{ km s}^{-1}$ in the LOS velocity measurement. With so few stars we found that a precise measurement of $\hat{\beta}$ is only possible for highly anisotropic galaxies that leave a strong imprint on the data. The performance of the method is thus dependent on the nature of the galaxy.

If the galaxy of interest is approximately isotropic ($\beta = 0$), then the statistical noise limits an inference of β to setting upper and lower bounds. Given that we effectively have no a priori intuition for β this is still useful information and excluding models with radical anisotropy would verify the implicit assumptions of widely used mass estimators.

The results are much more dramatic if the galaxy is in fact highly anisotropic. In this case, the estimators $\hat{\beta}$ are clustered tightly about the true value. Our numerical tests with 500 stars show that an observation of $\hat{\beta} = 0.5$ is in tension with tangential models $\beta < 0$ at the 2σ level. If such an observation were combined with the large 2D data sets available for dwarf spheroidals, then the degeneracy can be broken.

In numerical tests with simple Gaussian error distributions, we found that the estimators $\hat{\beta}$ are robust to errors in the LOS velocity measurements but prone to bias from uncertainties in the distance measurements. For our test model dwarfs with a half-light radius of 250 pc, the bias became highly significant as distance errors climbed above 150 pc. This is only a guide and it is beyond the scope of this work to fully quantify the bias as in practice the relevant error distributions will depend on the observational method. Where possible, we have briefly shown how one can use the data to identify and clean elements of this bias.

An additional issue is the assumption of spherical symmetry which is in tension with measurements of real Milky Way dwarf spheroidals. Numerical tests must be conducted to quantify how well the estimators fare on non-spherical haloes but the spherical model nevertheless gives us an interesting insight into the opportunities that the 3D positional information provides. The general idea holds for non-spherical stellar distributions; with 3D positional data we can see exactly how much each velocity component contributes to a stars LOS. This varies locally and by splitting or reweighting the data according to its position we can isolate the velocity anisotropy and break the degeneracy.

Finally, we argue that our method could successfully be applied to Sagittarius. Our numerical tests on mock dwarf spheroidal data indicate that we must know the relative depth of stars in a galaxy to within 150 pc. As noted in the feasibility section, dwarf galaxies

typically have a population of several hundred RR Lyrae. Via the BW method we can measure the distance to the star directly. The BW method is sensitive to accurate measurements of the surface temperature and the radial velocity of the stars surface. With a highly accurate spectrograph and coverage with multiple bands (especially with the addition of the K band), it may well be possible to apply our method directly to Sagittarius. If we can improve the inferred temperature of an RR Lyrae by a factor of 3 or 4, we could also apply the method to Draco, Ursa Minor, and potentially other nearby dwarf galaxies.

ACKNOWLEDGEMENTS

TR thanks the KCL Graduate School and Malcolm Fairbairn for support and Paolo Gondolo and Celine Boehm for useful discussions. He would also like to thank Jorge Penarrubia and Matthew Walker for making their mock dSph data publicly available. The research of DS has been supported at IAP by the ERC project 267117 (DARK) hosted by Université Pierre et Marie Curie – Paris 6. We would like to acknowledge the Mainz Institute for Theoretical Physics (MITP) for enabling us to complete a significant portion of this work.

REFERENCES

- Ackermann M. et al., 2011, *Phys. Rev. Lett.*, 107, 241302
 Agnello A., Evans N. W., 2012, *ApJ*, 754, L39
 An J., Evans N. W., Deason A. J., 2012, *MNRAS*, 420, 2562
 Baade W., 1926, *Astron. Nachr.*, 228, 359
 Baade W., Swope H. H., 1961, *AJ*, 66, 300
 Barcza S., 2010, *MNRAS*, 406, 486
 Barcza S., Benkő J. M., 2012, *MNRAS*, 419, 2774
 Battaglia G., Helmi A., Breddels M., 2013, *Astron. Rev.*, 57, 52
 Bersier D., Wood P. R., 2002, *AJ*, 123, 840
 Binney J., Tremaine S., 1987, *Galactic Dynamics*. Princeton Univ. Press, Princeton, NJ
 Binney J. et al., 2014, *MNRAS*, 437, 351
 Bono G., Caputo F., Castellani V., Marconi M., Storm J., 2001, *MNRAS*, 326, 1183
 Bouy H., Bertin E., Moraux E., Cuillandre J.-C., Bouvier J., Barrado D., Solano E., Bayo A., 2013, *A&A*, 554, A101
 Buckley D. R. V., Longmore A. J., Dixon R. I., 1992, *Mem. Soc. Astron. Ital.*, 63, 433
 Cacciari C., Clementini G., Fernley J. A., 1992, *ApJ*, 396, 219
 Catelan M., Pritzl B. J., Smith H. A., 2004, *ApJS*, 154, 633
 Chakrabarty D., Portegies Zwart S., 2004, *AJ*, 128, 1046
 Charbonnier A. et al., 2011, *MNRAS*, 418, 1526
 Dejonghe H., Merritt D., 1992, *ApJ*, 391, 531
 Deshpande R. et al., 2013, *AJ*, 146, 156
 Gentile G., Salucci P., Klein U., Vergani D., Kalberla P., 2004, *MNRAS*, 351, 903
 Genzel R., Pichon C., Eckart A., Gerhard O. E., Ott T., 2000, *MNRAS*, 317, 348
 Governato F. et al., 2012, *MNRAS*, 422, 1231
 Kinemuchi K., Harris H. C., Smith H. A., Silbermann N. A., Snyder L. A., La Cluyzé A. P., Clark C. L., 2008, *AJ*, 136, 1921
 Kolenberg K., 2013, in Suárez J. C., Garrido R., Balona L. A., Christensen-Dalsgaard J., eds, *Astrophys. Space Sci. Proc.*, Vol. 31, *Stellar Pulsations: Impact of New Instrumentation and New Insights*. Springer-Verlag, Berlin, p. 109
 Leonard P. J. T., Merritt D., 1989, *ApJ*, 339, 195
 Łokas E. L., 2002, *MNRAS*, 333, 697
 Łokas E. L., Mamon G. A., Prada F., 2005, *MNRAS*, 363, 918
 Lutz T. E., Kelker D. H., 1973, *PASP*, 85, 573
 Madore B. F. et al., 2013, *ApJ*, 776, 135
 Magorrian J., 2014, *MNRAS*, 437, 2230
 Navarro J. F., Frenk C. S., White S. D. M., 1996, *ApJ*, 462, 563
 Navarro J. F., Frenk C. S., White S. D. M., 1997, *ApJ*, 490, 493
 Read J. I., Gilmore G., 2005, *MNRAS*, 356, 107
 Read J., Gieles M., Kawata D., 2013, *First Gaia Challenge Workshop*, available at: <http://astrowiki.ph.surrey.ac.uk/dokuwiki/doku.php?>
 Richardson T., Fairbairn M., 2013, *MNRAS*, 432, 3361
 Sandage A., 1990, *ApJ*, 350, 603
 Sohn S. T., Anderson J., van der Marel R. P., 2012, *ApJ*, 753, 7
 Soszynski I. et al., 2011, *Acta Astron.*, 61, 1
 Spergel D. N., Steinhardt P. J., 2000, *Phys. Rev. Lett.*, 84, 3760
 Steinmetz M. et al., 2006, *AJ*, 132, 1645
 Strigari L. E., Bullock J. S., Kaplinghat M., 2007, *ApJ*, 657, L1
 Tonini C., Lapi A., Salucci P., 2006, *ApJ*, 649, 591
 Walker M. G., Penarrubia J., 2011, *ApJ*, 742, 20
 Walker M., Penarrubia J., 2013, *Spherical Models for the Gaia Challenge*, available at: http://astrowiki.ph.surrey.ac.uk/dokuwiki/lib/exe/fetch.php?media=data:gaiachallenge_spherical.pdf
 Walker M. G., Mateo M., Olszewski E. W., 2009, *AJ*, 137, 3100
 Watkins L. L., Evans N. W., An J. H., 2010, *MNRAS*, 406, 264
 Wesselink A. J., 1946, *Bull. Astron. Inst. Neth.*, 10, 91
 Wesselink A. J., 1969, *MNRAS*, 144, 297
 Wilkinson M. I., Kleyna J., Evans N. W., Gilmore G., 2002, *MNRAS*, 330, 778
 Wolf J., Martinez G. D., Bullock J. S., Kaplinghat M., Geha M., Muñoz R. R., Simon J. D., Avedo F. F., 2010, *MNRAS*, 406, 1220
 Zavala J., Vogelsberger M., Walker M. G., 2013, *MNRAS*, 431, L20

This paper has been typeset from a $\text{\TeX}/\text{\LaTeX}$ file prepared by the author.