



HAL
open science

Thermal Structure and Water Climatology of the Martian Middle-Upper Atmosphere from ACS/TGO Spectroscopy

D. A. Belyaev, J. Alday, Franck Montmessin, Franck Lefèvre, A. Fedorova,
Oleg Korablev, A. Trokhimovskiy, A. Patrakeev, K. S. Olsen

► **To cite this version:**

D. A. Belyaev, J. Alday, Franck Montmessin, Franck Lefèvre, A. Fedorova, et al.. Thermal Structure and Water Climatology of the Martian Middle-Upper Atmosphere from ACS/TGO Spectroscopy. Seventh international workshop on the Mars atmosphere: Modelling and observations, Jun 2022, Paris, France. pp.id.2102. insu-03691046

HAL Id: insu-03691046

<https://insu.hal.science/insu-03691046>

Submitted on 8 Jun 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

THERMAL STRUCTURE AND WATER CLIMATOLOGY OF THE MARTIAN MIDDLE-UPPER ATMOSPHERE FROM ACS/TGO SPECTROSCOPY

D. A. Belyaev, *Space Research Institute of the Russian Academy of Sciences (IKI), Moscow, Russia (bdenya.iki@gmail.com)*, **J. Alday**, *School of Physical Sciences, The Open University, UK*, **F. Montmessin**, **F. Lefèvre**, *LATMOS/CNRS, Paris, France*, **A. Fedorova**, **O. Korablev**, **A. Trokhimovskiy**, **A. Patrakeevev**, *Space Research Institute of the Russian Academy of Sciences (IKI), Moscow, Russia*, **K. S. Olsen**, *Department of Physics, University of Oxford, UK*.

Introduction:

The thermal structure of the middle (50-100 km) and upper (above 100 km) atmosphere of Mars holds key information relevant for climate modeling, atmospheric escape, understanding the impact of solar activity, as well as planning spacecraft maneuvers. The altitude range of 80-130 km hosts the layers of mesopause, the coldest atmospheric zone, and homopause, where the atmosphere is no longer uniformly mixed [1]. Moreover, water vapour molecules above ~80 km, when rises so highly [2-4], are effectively exposed by photolysis from the solar radiation that leads to H and O atoms escape.

In this paper, we report highly sensitive measurements of the vertical distribution of temperature, CO₂ and H₂O densities as measured by the Atmospheric Chemistry Suite (ACS) on board ExoMars Trace Gas Orbiter (TGO) in the regime of solar occultation [5]. The middle-IR channel (ACS-MIR) has been performing the experiment since April 2018 in the spectral range from 2.3 to 4.2 μm with the resolving power exceeding 25 000. In the occultation mode, the instrument senses CO₂ absorption band around 2.7 μm in an extremely broad altitude range, from 20 to 180 km, covering the troposphere, the mesosphere and the thermosphere of Mars. In parallel, the strong H₂O band at 2.66-2.67 μm is also observed for the first time up to 120 km.

In the paper, we discuss our scheme of the temperature and CO₂, H₂O density retrievals [4, 6] validating with atmospheric models and with simultaneous measurements by ACS-NIR below 100 km [2]. The climatology is provided by the seasonal and latitude coverage of observations that includes about 600 vertical profiles spreading over 1.5 Martian Years (MY), from the middle of MY 34 to the end of MY 35. The dataset allows observing seasonal and latitudinal variations of the thermal structure at the dawn and dusk terminators. Temperature and pressure conditions for H₂O and CO₂ saturation are analyzed as well.

Results of observations:

Our measurements cover several Martian seasons containing the second half of MY 34 and the entire MY 35, which correspond to ACS MIR observations from May 2018 to January 2021. The selected data

set comprises 308 occultation sessions in the Northern Hemisphere and 301 sessions in the Southern Hemisphere, encompassing seasonal periods from $L_S = 180^\circ$ to $L_S = 356^\circ$ in MY 35. We highlight several key points of our observations:

- Seasonal variability of the Martian temperature and CO₂ density is observed at altitudes from 20 to 180 km [6].
- The mesopause altitude rises from 70-90 km in the high latitudes of winter to 130-150 km in the summer season for both hemispheres [6].
- The homopause altitude varies from 80 km at aphelion to 110 km at perihelion in the MY 34 and 35, and it depends on dust activity [6].
- For the first time, water relative abundances are reported in a previously unexplored altitude range: from 100 to 120 km [4].
- Both the global dust storm in MY 34 and the two perihelion seasons (MY34, 35) reveal 10–50 parts of H₂O per million by volume (ppmv) at 100–120 km. At the same time, it does not exceed 2 ppmv during aphelion of MY 35 [4].
- High altitude H₂O super saturation conditions (>10) occur at 80-110 km every perihelion season. For CO₂, the condensation episodically occurs at aphelion Polar Regions in troposphere altitudes.

Acknowledgements:

The ACS experiment is led by the Space Research Institute (IKI) in Moscow, assisted by LATMOS in France. Retrievals and analysis of temperature and water distributions, performed by the authors at IKI, are funded by grant #20-42-09035 of the Russian Science Foundation.

Bibliography:

- [1] Bougher et al. (2017). Chapter 14: Upper Atmosphere and Ionosphere, in *The Atmosphere and Climate of Mars*, ed. B. Haberle, M. Smith, T. Clancy, F. Forget, R. Zurek, Cambridge University Press, <https://doi.org/10.1017/9781107016187>.
- [2] Fedorova et al. (2020). Stormy water on Mars: The distribution and saturation of atmospheric water during the dusty season. *Science*, 367(6475), 297–300. <https://doi.org/10.1126/science.aay9522>.
- [3] Fedorova et al. (2021). Multi-annual monitor-

ing of the water vapor vertical distribution on Mars by SPICAM on Mars Express. *Journal of Geophysical Research: Planets*, 126, 2020JE006616. doi:10.1029/2020JE006616.

[4] Belyaev et al. (2021). Revealing a high water abundance in the upper mesosphere of Mars with ACS onboard TGO. *Geophysical Research Letters*, 48, e2021GL093411. <https://doi.org/10.1029/2021GL093411>.

[5] Korablev et al. (2018). The Atmospheric Chemistry Suite (ACS) of three spectrometers for the ExoMars 2016 trace gas orbiter. *Space Science Reviews*, 214(1). <https://doi.org/10.1007/s11214-017-0437-6>.

[6] Belyaev D. et al. (2022). Thermal Structure of the Middle and Upper Atmosphere of Mars from ACS/TGO CO₂ Spectroscopy. *Submitted to Journal of Geophysical Research: Planets*. Under Review.