



HAL
open science

Potential impacts of climate, land use and land cover changes on hydropower generation in West Africa: a review

Salomon Obahoundje, Arona Diedhiou

► **To cite this version:**

Salomon Obahoundje, Arona Diedhiou. Potential impacts of climate, land use and land cover changes on hydropower generation in West Africa: a review. *Environmental Research Letters*, 2022, 17, 10.1088/1748-9326/ac5b3b . insu-03706380

HAL Id: insu-03706380

<https://insu.hal.science/insu-03706380>

Submitted on 28 Jun 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

TOPICAL REVIEW • OPEN ACCESS

Potential impacts of climate, land use and land cover changes on hydropower generation in West Africa: a review

To cite this article: Salomon Obahoundje and Arona Diedhiou 2022 *Environ. Res. Lett.* **17** 043005

View the [article online](#) for updates and enhancements.

You may also like

- [Impacts of changing snowfall on seasonal complementarity of hydroelectric and solar power](#)
Adrienne M Marshall and Jie M Chen
- [Simulation of hydropower at subcontinental to global scales: a state-of-the-art review](#)
Sean W D Turner and Nathalie Voisin
- [Impacts of climate change on hydropower development and sustainability: a review](#)
J Shu, J J Qu, R Motha et al.

ENVIRONMENTAL RESEARCH
LETTERS

TOPICAL REVIEW

OPEN ACCESS

RECEIVED
20 September 2021REVISED
10 February 2022ACCEPTED FOR PUBLICATION
7 March 2022PUBLISHED
21 March 2022

Original content from
this work may be used
under the terms of the
[Creative Commons
Attribution 4.0 licence](#).

Any further distribution
of this work must
maintain attribution to
the author(s) and the title
of the work, journal
citation and DOI.

Potential impacts of climate, land use and land cover changes on
hydropower generation in West Africa: a reviewSalomon Obahoundje¹ and Arona Diedhiou^{1,2,*} ¹ Laboratoire des Sciences de la Matière, de l'Environnement et de l'Energie Solaire (LASMES)—African Centre of Excellence on Climate Change, Biodiversity and Sustainable Development/Université Félix Houphouët Boigny, 22 BP 582 Abidjan 22, Abidjan, Ivory Coast² University Grenoble Alpes, IRD, CNRS, Grenoble INP, IGE, F-38000 Grenoble, France

* Author to whom any correspondence should be addressed.

E-mail: aronadiedhiou@ird.fr**Keywords:** West Africa, climate change, land use, land cover, hydropower, nexus, sustainability**Abstract**


This study aims to review the existing literature on the past and future effects of climate, land use, and land cover changes on hydropower generation in West Africa (WA), based on listings in the Scopus and Google Scholar databases. This review shows that several African hydropower plants have experienced repeated power disruptions over the last three decades due to climate change and variability but it is less documented how increasing land use and land cover changes around the major dams have impacted the hydrological system and the hydropower generation. In the future, the risks of hydropower in WA may not be equally distributed within a country or region. Despite uncertainties in precipitation and on impacts on streamflow and water level in major basins, climate change is likely to reduce the available water over the range of 10%–20% (15%–40%) for the RCP4.5 (RCP8.5) scenario by 2050, which may considerably affect the water demand across all sectors, including hydropower. However, in the Kainji dam (Niger River basin), models project an increase in rainfall favorable to hydropower production for both RCP4.5 and RCP8.5. In contrast, within the Black Volta sub-basin, the intensification of land use is predicted to favor runoff and, consequently, an increase in the generation of Bui hydropower in the near future, even though models predict a rainfall decrease. This increase in land use for agriculture to feed a growing population has other adverse effects that need to be assessed, namely sedimentation and siltation, which are harmful to hydropower plants. Finally, the combined impact of climate and land use changes on the efficiency of hydroelectric infrastructure in WA is not well documented, while sustainable planning and investments in the hydropower sector require consideration of the nexus between climate, land use changes, and water.

1. Introduction

The production of energy from fossil fuels is harmful to the environment in the long term and is considered to be the main source of anthropogenic greenhouse gas (GHG) emissions that contribute to climate change (IPCC 2011). According to the International Energy Agency (IEA), fossil fuels remain the largest source of energy in Africa, with an 80% share of total electricity generation, while 15.52% comes from hydropower, and the remainder from other sources (nuclear, waste, wind, and solar) (IEA 2019). In West Africa (WA), hydropower is the main source of renewable energy. In 2017, the total electrical

energy production was respectively 32 249 GWh in Nigeria (17.14% from hydropower), 14 068 GWh in Ghana (39.92% from hydropower), 8575 GWh in Ivory Coast (17.83% from hydropower), 232 GWh in Togo (88.4%), and 4777 GWh (7.05%) in Senegal (IEA 2019). The hydropower plants already built in Ghana, Nigeria, and Ivory Coast represent more than 80% of the total installed capacity of WA.

To fulfill the Paris Agreement, measures have been taken to promote clean, renewable energy, such as hydropower, to mitigate climate change. Indeed, WA has several transboundary rivers suitable for hydropower. This is confirmed by the figure 1 we plotted in overlaying data from Dams Atlas and from



Global Runoff Data Centre. Figure 1 shows that 22 dams are already operational with total installed capacity of 5401.6 MW, while a larger number are in the planning stages (82 dams with total installed capacity of 18 679.60 MW), confirming the commitment of West African countries to increase the share of renewable energy to comply with the Paris Agreement in addressing the future energy demand. Paradoxically, climate change itself may alter the availability of this natural resource (water for hydropower), adversely affecting the financial viability of both existing and planned infrastructures (Shu *et al* 2018). A similar conclusion was found by Almeida *et al* (2021), who demonstrated that climate change may undermine power generation as well as the economic viability of future Amazon hydropower.

Climate change is predicted to create high rainfall variability and increase in temperature, making water availability uncertain (IPCC 2011). It may exacerbate the water stress currently faced by some countries, lengthening the list of countries at risk of water stress. In all West African basins, the relationships between precipitation, runoff, and discharge are complex due to land use/cover change (LULCC). The Senegal, Niger, and Volta basins, where the major hydropower plants are built, are projected to be severely affected by climate change and variability (Oyebande and Odunuga 2010, Oguntunde and Abiodun 2013). However, studies on the specific influences of climate change and the magnitude of these possible effects on

hydropower in the region are nascent. Owing to the high risk on water availability and variability, future hydropower projects may be significantly affected by changes in climatic conditions (IPCC 2011).

Because of the importance of hydropower in the development of African countries and for the fulfillment of the Paris Agreement, scientific interest in hydropower in Africa is growing, but existing studies have focused mainly on the southern part of the continent (Hamududu and Killingtveit 2016). According to the West African Regional Center on Renewable Energy and Energy Efficiency (ECREEE) (ECREEE 2017), the impact of climate change on West African water resources is well known, but the effects on hydropower generation are not well documented, probably because of the lack of historical data.

In addition to climate change, there are other factors such as population growth, LULCC, water abstraction, and socio-economic development, which can affect hydropower generation. Freshwater ecosystems in Africa are threatened by anthropogenic land use changes, overexploitation of water, diversion of rivers and lakes, increased pollution, and sediment loading in water bodies (IPCC 2011). Indeed, the West African population is increasing rapidly, inducing faster land use/cover changes and impacting the dynamic behavior of hydrological systems (Descroix *et al* 2009). While most West African basins are transboundary with different land use practices and water abstraction, few studies have investigated the impacts

of LULCC on hydropower generation in the region. Additionally, the combined effects of land use/cover and climate change are not well understood. This is a challenge for the sustainability of hydropower in the WA region.

According to ECREEE (2017), most hydropower management systems in the region and local actors have not integrated climate change and/or LULCC into their operations and planning. The main barrier to integrating climate change and LULCC into decision-making for sustainable development is that, on the one hand, the impact of climate on hydropower is uncertain and not evenly distributed across the region, and on the other hand, the few studies on LULCC change in sub-Saharan Africa are not designed to address the issue of the nexus between climate, land, water and energy. More research leading to climate service development is needed to enable dam managers and local political authorities to make decisions under these uncertainties and to mainstream climate change and LULCC in their management plans.

This review aims to synthesize what is known on past and future impacts of climate change and LULCC on hydropower generation in WA. The review methodology based on publications available in the Scopus and Google Scholar databases is presented in section 2. Hydroclimatic data and satellite products used to make the figures supporting the main results are also presented in section 2. Results of the literature review are presented in section 3: the impacts of climate change on water resources and hydropower are presented in section 3.1. Then, the impacts of land use and land cover dynamics on water resources and hydropower generation are presented in section 3.2. Finally, what is known about the combined effects of climate and land use/cover changes on hydropower generation in the main dams of the WA region is presented in section 3.3, with an overall synthesis on the climate—land use/cover—hydropower nexus. The conclusions are presented in section 4, with research perspectives.

2. Method and data

2.1. Methodology

The review protocol was adapted from the Preferred Reporting Items for Systematic Reviews and Meta-Analyses reporting guidelines for scoping reviews (PRISMA-ScR; (Chang 2018, Munn *et al* 2018, Tricco *et al* 2018)). After the identification of the research questions and keywords, a selection of relevant peer-review papers in Scopus and Google Scholar is done. Then, selected articles are classified into five thematic clusters: ‘climate change and water resources’ (CC_WR), ‘climate change and hydropower’ (CC_HP), ‘land use/cover and water resources’ (LULCC_WR), ‘climate change, land use/cover and water resources’ (CC_LULCC_WR),

and ‘climate change, land use/cover and hydropower’ (CC_LULCC_HP).


After the analysis of papers in each cluster, a synthesis of the state of knowledge was performed according to the following three steps, representing the subsequent sections of this manuscript: (a) impacts of climate change on water resources and hydropower, (b) impacts of land use/cover changes on water resources and hydropower, and (c) the nexus between climate, land use/cover, and hydropower with a summary table of existing studies on major hydropower plants of WA.

The following question guided this review: what is known in the existing published research about the past and future impacts of climate change and LULCC on water resources and hydropower in WA? Within this broad question, this review aimed to answer the following questions: how do climate change and LULCC affect hydropower generation in WA? What are the main factors of the nexus between climate and land use/cover that drive hydropower generation in WA?

We first searched the Scopus database for publications with the set of keywords ‘West Africa’, ‘climate’, ‘land use and land cover’, ‘hydropower’ in the abstract, title, and keywords. We found a few peer-reviewed papers discussing the nexus in WA, and we noted 46 available and up-to-date papers. We then reoriented our investigation, filtering by ‘West Africa’, ‘climate’, and ‘hydropower’ keywords, and around 198 papers were found. Next, the keywords ‘West Africa’, ‘land use’, and ‘hydropower’ were used, and only 119 peer-reviewed papers were found to be available. Papers published before the year 2000 and cited by recent studies on the same West African basins were removed from the list of selected articles, as well as several articles dealing with southern or eastern Africa and citing results from West African rivers or basins (Whipple and Viers 2019). Finally, 66 papers published after 2000 were found to be relevant, 78.78% (52) of them were issued after 2010.

Google Scholar was used to perform the bibliometrics because its algorithm searches for keywords in the main text (not only in the abstract) and because it can suggest gray literature (Haddaway *et al* 2015, Moed *et al* 2016). The first list of 400 papers was found, among which 300 papers were also listed in Scopus. By filtering the 100 remaining papers manually and removing academic reports such as PhDs and Master’s theses and conference abstracts, we selected 30 papers. Among these papers, 26 were published after 2010, 18 were research articles, and the others were relevant reports published by national or transboundary water basin agencies, national or regional agencies of electricity, or by international agencies of the United Nations system with quantitative values on water resources and hydropower in WA.

Figure 2 shows the percentage of papers per thematic cluster. Conclusively, 48.96% and 16.67%



of the 96 selected papers were focused on CC_WR and CC_HP, respectively. Only 11.46%, 17.71%, and 5.21% dealt thematically with LULCC_WR, CC_LULCC_WR, and CC_LULCC_HP, respectively. It is worth noting that we did not find any studies specifically focused on the impact of LULCC on hydropower generation in WA, even though the areas around the existing hydropower plants or multipurpose dams are under pressure from multiple socio-economic activities such as human settlement, fisheries, vegetable farming, and agribusiness (Mahe *et al* 2013).

Finally, using the approach taken by Whipple and Viers (2019), the main results from the clusters above are merged and presented in the subsequent subsections. We created flowcharts at the end of each section to synthesize the impacts of climate change on water resources and hydropower (section 3.1), the impacts of land use/cover changes on water resources and hydropower (section 3.2), and the nexus between climate, land, and hydropower (section 4), with a summary table on what is known about the main drivers of past and future changes in hydropower generation in the major WA plants (table 1).

2.2. Hydropower information, hydroclimatic data, and satellite products

Information on hydropower plants (existing and planned) was extracted from Dams Atlas (<http://globaldamwatch.org/data/>) (Zarfl *et al* 2015) and was completed with measured data retrieved directly from

hydropower plant managers (national companies of electricity and river basin authorities) in the region. This was a big gap: there was no place where those data were centralized or made public, and such efforts to gather multidisciplinary data around the dams and information from the different dam managers on how they operate revealed the challenges in the assessment of the impacts, in the sustainability of those infrastructures, and the barriers to performing interdisciplinary studies on the nexus between climate, land, hydropower, and water.

We plotted maps of the spatial distribution of hydropower plants with areas of significant change in precipitation, runoff, and normalized difference vegetation index (NDVI) to determine how each plant is exposed to past and future changes in climate and land cover in WA (figures 3 and 5, respectively). Specifically, we created a map of areas of change in precipitation (figure 3(a)) using the Climate Hazard Group Infrared Precipitation with Stations (CHIRPS) product (1981–2018). CHIRPS is a 30 year quasi-global rainfall dataset, stretching over 50° S–50° N (and all longitudes), covering the period 1981 to near-present and incorporating 0.05° resolution satellite imagery with *in situ* station data to create gridded rainfall time series for trend analysis and seasonal drought monitoring (Funk *et al* 2015). Figure 3(b) was adapted from Stanzel *et al* (2018) study to determine which hydropower plant is (or will be) located in an area with a significant projected decrease/increase in runoff. For this

purpose, we adapted the map of the projected change in runoff over 2046–2065 relative to 1998–2014 from an ensemble of 15 regional climate models (RCMs) of coordinated regional climate downscaling experiment (CORDEX)-Africa and we added the locations of existing and planned dams.

We then computed a map of areas of change in vegetation cover (figure 5) using the NDVI-3rd generation from the Global Inventory Monitoring and Modeling System (National Center for Atmospheric Research Staff (eds) 2018). This NDVI product is a monthly dataset covering the second semester of the 1981 year to the end of 2015 with a very high resolution ($1/12^\circ$). A student test (t -test) was performed to detect areas of significant change between two periods in both precipitation (2000–2018 relative to 1981–1999) and the NDVI (2006–2015 relative to 1982–1991) dataset. Existing and planned dams are added in this figure to illustrate the potential linkages between land cover changes and hydropower.

Finally, in each section, we created flowcharts (figures 4 and 6) based on the literature review to summarize how climate change (precipitation and temperature) or LULCC may affect hydropower production. The last flowchart in figure 7 illustrates the main features of the climate, land, water, and energy nexus based on the literature.

3. Results

3.1. Impact of climate change on water resources and hydropower


The climate of the WA depends on the West African monsoon variability, the meridional movement of the inter-tropical convergence zone (ITCZ), as well as the position and intensity of the African easterly jet and the tropical easterly jet (Biasutti 2019). The ITCZ is the region where the hot and dry harmattan air mass from the Sahara in the north (northeasterly trade wind) meets the cool, moist monsoon air (southwesterly wind) from the South Atlantic. Between December and February, the ITCZ moves toward the Gulf of Guinea, and from March to November, it moves from the Gulf to the higher latitudes, even crossing some areas twice. Therefore, in the southern part of WA (i.e. the Guinea Coast), there are two rainy seasons (March–June and August–November) and two dry seasons (July and December–February). The northern part of the WA (i.e. the Sahel band) has one rainy season from July to September and a long dry season from October to June.

In the 1970s and 1980s, WA experienced the most severe drought recorded in the 20th century, with a 15%–40% drop in rainfall associated with a decrease in the number of rainy events (Le Barbé *et al* 2002, Nicholson 2013). Later, the inter-decadal variability in West African rainfall was attributed to large-scale changes in the Atlantic sea surface temperature, El

Niño Southern Oscillation, and La Nina (Nicholson 2013), as well as regional and local-scale changes in land use and vegetation cover (Biasutti 2019).

The last 30 years have been characterized by greater interannual variability than the previous 40 years over the WA region in general, and in the Sahel subregion in particular (Taylor *et al* 2017). Since the 1990s, the rainfall intensity has increased over the entire WA domain, causing the north-eastern part (central Sahel) of the WA region to become wetter (Panthou *et al* 2013, Taylor *et al* 2017). Although the Sahel has seen a ‘recovery’ in rainfall since the 1980s, cumulative precipitation has not returned to pre-1960s levels and certain characteristics have changed (Biasutti 2019): rainfall events appear to be less frequent and have a shorter duration but greater intensity (Nicholson 2013). Indeed, the West African Sahel became wetter during the period 1981–2014 due to an increase in the number of rainy days, longer wet spells, and shorter dry spells (Bichet and Diedhiou 2018b, Didi *et al* 2020). Along the coast of the Gulf of Guinea, the change in rainfall between 1981 and 2014 was associated with less frequent but more intense rainfall events over both rainy seasons (Bichet and Diedhiou 2018a, Didi *et al* 2020). However, between 1970 and 1990, decreased rainfall in the southern region of WA affected the entire regional hydrological system (Mahe *et al* 2005a), as well as water-related sectors within the basins, namely, irrigation, domestic, livestock, and hydropower generation (Tossou *et al* 2017). Using the CHIRPS precipitation product, and after computing a Student’s t -test at a 95% confidence level to highlight regions with significant changes, we confirm in figure 3(a) that the region above 11° N became significantly wetter during the last two decades (2000–2018), relative to the 1981–1999 period, whereas some parts of Sierra Leone, Guinea, Liberia, and Nigeria exhibited a significant decrease. Among existing dams, only two are located in the Guinea-Conakry Republic (Garafiri and Kaléta) and two major dams in Ghana (Akosombo and Kpong) are located in areas that have experienced a significant decrease in precipitation. It has been reported that the Garafiri dam experienced issues during this period (World Bank 2017) and that the 1998 drought resulted in low water levels in the Akosombo reservoir, leading to an energy crisis in Ghana (Kabo-Bah *et al* 2016, 2018).

Projections of WA precipitation by the Coupled Model Inter-comparison project CMIP3 and CMIP5 archives showed an inter-model variation in both the amplitude and sign of change (Rowell 2012, Rowell *et al* 2016) which is partially attributed to the inability of global climate models (GCMs) to resolve convective rainfall. Projections indicate a decrease and increase in precipitation in the Western Sahel (Senegal) and the Central Sahel respectively in 2100, more or less intense depending on the scenario (Biasutti 2019). The Western Sahel is projected by



80% of the models to experience the strongest drying at 1.5 °C global warming, with a significant increase in the length of dry spells and a decrease in the standardized precipitation evapotranspiration index

(Diedhiou *et al* 2018). In the regions of the Guinea Coast and Central Africa, most CMIP5 GCMs project a weak change in total precipitation associated with a decrease in the length of wet spells and an increase

in heavy rainfall under a global warming of 1.5 °C (Diedhiou *et al* 2018).

RCMs involved in the CORDEX have shown acceptable performance in simulating the spatial and temporal distribution of the main precipitation and temperature features (Gbobaniyi *et al* 2014, Nikulin *et al* 2018). Despite the spread in the magnitude from one model to another (Nikulin *et al* 2018), the ensemble mean considerably improves the performance of most of the individual RCMs (Gbobaniyi *et al* 2014). Combining 19 RCMs of CORDEX, Kumi and Abiodun (2018) showed that the RCMs projected a delay of rainfall onset dates and shorter rainy seasons in the western and eastern Sahel under 1.5 °C and 2 °C global warming for the RCP4.5 and RCP8.5, respectively. Using 25 RCMs of CORDEX Africa, more than 80% of ensemble members agreed that the number of consecutive dry days will increase over the Guinea Coast, with a projected decrease in consecutive wet days at both 1.5 °C and 2 °C global warming levels (Klutse *et al* 2018).

The surface temperature over West Africa has increased over the last 50 years, in the range of 0.5 °C–0.8 °C between 1970 and 2010 (Riede *et al* 2016), with the increase in the temperature being greater in magnitude within the last 20 years than in the preceding decades. It was also found that the number of cold days and nights decreased, while the number of warm days and nights increased over the 1961–2000 period (Riede *et al* 2016). Since the 1950s, diurnal and nocturnal heatwaves over the Gulf of Guinea coastal regions intensified, becoming more pronounced after the 1980s (Ringard *et al* 2016). The rise in temperature has impacted all sectors, including water resources, energy demand, and production (Riede *et al* 2016). Temperature increases within the West African domain are projected to be higher than the global mean temperature increase, and heat waves are expected to be more frequent and lasting longer (Diedhiou *et al* 2018). This will lead to both an increase in the energy demand and evaporation of water in the reservoir needed for hydropower production, and will affect the performance of the plants if no appropriate investments are made to integrate hydro-climate services in the management system.


Regarding the discharge, a study with an ensemble means of 30 regional climate simulations of CORDEX models based on two scenarios (RCP4.5 and RCP8.5) showed that stronger decreases are projected for regions in the northwestern (–40%) and southeastern (–10%) of WA, and pronounced increases are expected mainly for the southwest (+10%) over 2046–2065, relative to 1998–2014 (Stanzel *et al* 2018). Apart from the southwestern regions (Liberia, Sierra Leone, Guinea) where more than 75% of the 30 examined models project increases in runoff, the remaining regions where decreases or unknown trends in runoff are projected could make the future of existing and planned hydropower plants uncertain

(see figure 3(b)). These decreases in runoff could affect all water consumption sectors, including existing and planned hydropower plants, and may lead to conflicts between users if no informed integrated water management system is used.

River flows are expected to decline by 15%–20% by 2020 and by 20%–40% by 2050 (Sylla *et al* 2018a). In transboundary basins such as Niger, Senegal, and Volta, river flows are expected to decrease in proportions varying between 5% and 34%, depending on the time horizon and location (Oyebande and Odunuga 2010, Sylla *et al* 2018a). Specifically, a multi-model GCM ensemble predicted a decrease of 8% (RCP4.5) or 16% (RCP8.5) in annual streamflow as soon as 2050 in the Senegal River Basin, and of 22% (RCP4.5) or 26% (RCP8.5) in the Gambia River Basin (Bodian *et al* 2018), with the Gambia River Basin being more affected in the near future. In the Guiers Lake in Senegal, it was found that climate change and population growth rate will pressurize water resource availability, leading to greater competition between agriculture and municipal demands (Djiby *et al* 2018). In the Volta Basin, all water demands (municipal, hydropower, and agriculture) may not be simultaneously met under any of the projected scenarios, including wet scenarios (Amisigo *et al* 2014). All listed countries rely on hydropower for electricity production.

Extreme climate events in past decades have severely affected hydropower systems in African countries such as Ghana, Kenya, Zimbabwe, and Tanzania, sometimes reducing plants to half of their output capacity (Cole *et al* 2014). In Ghana, energy production failures were frequently observed during the rainfall deficit period from 1990 to 2011, caused by low Akosombo dam levels, especially during the 1998 drought year (Kabo-Bah *et al* 2016). A decrease in mean rainfall with an increase in minimum and maximum annual temperatures was recorded at all weather stations, leading to increases in evaporation. In Togo, it was noted that the hydropower generation is more modulated by four main variables namely inflow to reservoir, water level, rainfall of the actual and the previous year (Obahoundje *et al* 2021a) which are function of climate system. The Kainji Dam in Nigeria has also faced the same challenges as consequences of changes in climate variables (Olofintoye and Adeyemo 2011, Salami *et al* 2015). The decline in hydropower generation at the Kainji Dam was associated with decreases in runoff (Salami *et al* 2015) and a drop in reservoir storage volume (Olofintoye and Adeyemo 2011).

Frequent flooding events in recent decades have affected the hydropower system in WA. The high occurrence of heavy rainfall events in 1999 over the Niger basin has led to exceedances of the Kainji, Jebba, and Shiroro dams' floodgates in Nigeria, causing deaths, socio-economic infrastructure losses, and several displacements (Oyebande and Odunuga 2010). The same issues were experienced in the Senegal Basin



in the same year and the Ghanaian part of the Volta Basin in 2002. It has also occurred in the Yobe Valley in Nigeria in 1998 and 2001 (Oyebande and Odunuga 2010).

Existing peer review papers have shown that globally, hydropower generation (for plants operating today in Africa) is predicted to change very little by the year 2050 (Hamududu and Killingtveit 2012) but it may vary from one country to another. For instance, Ghana, Guinea, and Ivory Coast are projected in 2050 to observe a decrease by -1.6% , -2.9% , and -6.2% in hydropower generation, respectively, while Nigeria and Sierra Leone may experience an increase of $+0.4\%$ and $+6.1\%$ respectively over the same period (Hamududu and Killingtveit 2012). Conversely, Turner *et al* (2017) predicted a reduction in most of the hydropower generation in WA countries (A2 and B1 SRES scenarios and different models) over 2040–2069 relative to the 1965–2000 period. They (Turner *et al* 2017) found that the following countries will experience the strongest negative change in hydropower generation: Guinea (-12.9%), Mali (-13.17%), Togo (-14.4%), Ghana (-14.5%), Burkina Faso (-15.3%), Côte d'Ivoire (-15.7%), and Nigeria (-15.8%).

Figure 4 summarizes the processes associated with climate impacts on surface hydrology and hydropower generation. An increase in precipitation may lead to a decrease in evaporation due to the loading of relative humidity in the atmosphere and an increase in river discharges and water levels, which then results in improved hydropower generation. Conversely, a decrease in precipitation will lead to a decrease in river discharge and an increase in evaporation due to a drier atmosphere, and therefore, will reduce the

water available in the reservoir for hydropower generation. The temperature rise will increase evaporation, which may contribute to a decrease in the water level in the reservoir and negatively affect hydropower production. The change in precipitation should be considered from two perspectives: firstly, by looking at the basin scale (increase in precipitation will increase reservoir inflows); then by looking at the reservoir area (e.g. increase in temperature will lead to higher reservoir evaporation). However, this increase in evaporation may also contribute to an increase in atmospheric moisture (feedback), and therefore, to an increase in precipitation which can counteracts the negative effect. The relationship between precipitation and evaporation has been documented in WA but this requires further investigation. Projected increases in the variability of temperature and precipitation extremes could enhance the risk of flood and drought events, which may be detrimental to hydropower plant management and planning (Blacksher *et al* 2011). Additionally, the change in extreme compound events may also affect hydropower generation, but the key driver in these compound cases is precipitation, which will tend to raise the water level, and therefore favor hydropower generation, while the cold/dry or warm/dry events will negatively affect hydropower generation. In arid and semi-arid areas, evaporation losses due to an increase in temperature could represent a significant fraction of drops in water levels (Saggai and Bachi 2018). In these drylands, the temperature may be a key driver. However, in both regions, wind, sunshine duration, and humidity gradient can modulate evaporation from reservoirs (Condie and Webster 1997). The issue is that the tropics, in general, are expected to experience

increases in temperature and precipitation, while in WA, several uncertainties remain on future precipitation changes.

In addition to precipitation and temperature, there are additional factors that can directly or indirectly affect hydrological systems and thus hydropower generation. Oyebande and Odunuga (2010) and USAID (2013) concluded that the uncertainties associated with the impacts of future climate change on water resources in WA are compounded by many other factors. These include regional demographic factors, non-existing or inadequate water policies, poor infrastructure, inefficient water management systems, and inadequate joint management of basin resources. The current freshwater demand in WA for energy, agriculture, industry, and domestic use is expected to triple by 2025 (USAID 2013). This means that the above factors should be carefully considered when modeling water resources within a basin. It is also very important to carefully consider the effect of upstream socio-economic development on reservoirs while assessing the impacts of climate change on hydropower systems (de Souza Dias *et al* 2018).

3.2. Impact of land use and land cover changes on water resources and hydropower

In WA, the regions above 16° N are mainly covered by barren land, with a NDVI between 0 and 0.1 ($0 < \text{NDVI} \leq 0.1$). The Sahelian zone between 11° N and 14° N is covered by sparse vegetation (shrubs and grasslands or senescing crops) ($0.2 < \text{NDVI} \leq 0.5$). The Sudanian and Guinean zones have NDVI values of approximately 0.6–0.9, corresponding to dense vegetation. We computed and plotted in figure 5, the changes in NDVI over 2006–2015 relative to 1982–1992, and the significance test at a 99% confidence level. The central and eastern Sahel and Guinean (coastline) zones showed a statistically significant decrease in NDVI, while the Sudanian zone presented an increase in NDVI (see figure 5). Overall, this change is validated by existing studies in the region. For instance, between 1975 and 2013, total vegetation cover decreased by 1.6%, associated with massive net gains in croplands (107.8%) and settlements (140%) at the expense of natural vegetation (Barnieh *et al* 2020). It was discovered that human-managed LULC types have largely replaced natural LULC types. It is important to note that, most existing and planned hydropower plants are located in areas with a significant decrease of NDVI, which is not without impact on the hydrological system. Indeed, it has been reported that runoff extremes are increasing in most regions of the world at rates higher than precipitation extremes, due to LULCC (Yin *et al* 2018). As shown in the figure 4, this change in the runoff due to LULCC will influence the streamflow and the inflow and consequently, the hydropower generation.


Indeed, runoff extremes have been reported to increase in most regions of the world at rates higher

than precipitation extremes due to LULCC (Yin *et al* 2018). Basins in WA are facing rapid changes in land use and land cover and have lost, or are losing large areas of their natural landscapes, which are being replaced by human-influenced extents dominated by agriculture, bush fires, and timber extraction (Vitteck *et al* 2013, Akpoti *et al* 2016, Cotillon 2017, Obahoundje *et al* 2018). Between 1975 and 2013, the forest area decreased by 24.6%, while settlement and agricultural land coverage increased by 140% and 11.7% respectively in WA (Cotillon 2017). The size of vegetated areas continued to decline with decreasing rainfall and increasing temperatures, especially in dry regions.

The hydrological responses to LULCC vary with the region (Descroix *et al* 2018, Osei *et al* 2019). In the Black Volta sub-basin, LULCC between 2000 and 2013 contributed to an increase in the surface runoff and lateral flow by 27% and 19%, respectively, while the contribution of groundwater to streamflow has decreased by 6% due to land use change, along with a 4.6% increment in evapotranspiration due to rising temperatures (Akpoti *et al* 2016, Obahoundje *et al* 2017). At all scales in the Sahelian areas, runoff coefficients have generally increased along with river discharges, stimulated by a decrease in vegetation cover (Lienou 2013, Descroix *et al* 2018). Runoff has increased in Sahelian basins despite a 20%–25% decrease in observed rainfall during the period of 1968–1997 (Descroix *et al* 2018, Galle *et al* 2018) and an increase in dams constructed in the basins (Mahe *et al* 2005b). This increase in runoff and discharge despite the decrease in rainfall is called the ‘Sahel paradox’ and was first observed in small basins of Burkina Faso by Albergel (1987). Contrarily, a 15% reduction in rainfall in Sudan has led to a more intuitive reduction in runoff and annual discharges (Descroix *et al* 2018). In the latter region, the decrease of discharges has been two to three times greater than that of rainfall (Descroix *et al* 2018).

Over Sahel, in endorheic areas, the increase in runoff has led to an increase in the volume and duration of ponds (Moussa *et al* 2009, Gardelle *et al* 2010) and in some sites, this has led to an increase through the ponds of the underground water table; this was first found by (Leduc *et al* 2001) and documented by (Cappelaere *et al* 2009) and (Favreau *et al* 2009). In exoreic areas, there is a linear relationship between rainfall, runoff and streamflow but there is not a connection between the rainfall, the runoff or the streamflow with the groundwater recharge (Mahe *et al* 2005b).

Over the southern region of WA, the decrease of vegetation cover may also contribute to decrease the soil water holding capacity and infiltration rate, leading to a decrease in the water table (Mahe *et al* 2005b). The reduction of the underground water recharge capacity or water table can cause a drastically dry season streamflow decrease, which in addition to the




increase of evaporation due to rising in temperature, may severely affect the reservoir water availability for power generation in the warmer season when more energy is needed for cooling (e.g: increase of demand for air conditioning).

Increases in streamflow may lead to floods, which in turn may threaten hydropower infrastructure and electricity generation, as well as human security. The increase in floods observed in several WA rivers is associated with an increase in runoff, leading to higher rates of soil loss and erosion, contributing to silting in river beds and increasing the size of the rivers and reservoirs (Mahe *et al* 2005b, Sighomnou *et al* 2013, Mamadou *et al* 2015). According to Welde and Gebremariam (2017), the increase in bare land and agricultural areas in the Tekeze dam watershed in Ethiopia have resulted in increased annual and seasonal streamflow and sediment yield volumes. An increase in sediment yield volumes associated with increasing land use has also been observed in China (Du Watershed and Loess Plateau) (Yan *et al* 2013, Zuo *et al* 2016). Bissadu *et al* (2017) demonstrated that changes in rainfall intensity and land cover types are the core drivers of soil erosion, and Alkharabsheh *et al* (2013) found similar results in Jordan. Over Asia, Yang *et al* (2003) found that 60% of soil erosion was anthropogenically induced. LULCC for agriculture and urbanization within a basin will expose more land to soil erosion (Steger *et al* 2019). Silting due

to soil erosion may affect water quality through sedimentation (Yan *et al* 2013, Zuo *et al* 2016), and may in turn damage turbines by blocking power intakes and through the abrasion of turbine blades, thus shortening turbine life. This increased sedimentation will subject hydropower projects to additional costs because dredging the rivers to ensure a suitable intake flow will become necessary to maintain some form of equilibrium (Blacksher *et al* 2011).

The potential impacts of land use and land cover changes on the Bui hydropower scheme in the Black Volta Basin under climatic change were investigated using the WEAP model from 2000 to 2040 (Obahoundje *et al* 2017). Evaluation using the WEAP model under wet and dry climate conditions showed that the combined effects of changes in land use/cover and climate will reduce the water available to all sectors of water demand, including the Bui hydropower facilities. The wet condition is considered as a 14% increase in rainfall and the dry condition as a 15% decrease in rainfall between 2012 and 2040. The results showed that under wet conditions and without LULCC, Bui hydropower energy production will decrease by 0.15% and will increase by 40.74% with LULCC. Under current conditions (from 2000 to 2012), a decrease of 2.49% and 23.2% was noted in simulations with and without LULCC, respectively. Under dry conditions, the LULCC and no LULCC scenarios will decrease Bui hydropower generation



by 46.34% and 54%, respectively (Obahoundje *et al* 2017). The increase in land use within the Black Volta Basin is predicted to favor Bui hydropower generation in the near future. However, despite the positive effect of LULCC on the increase of runoff and streamflow and on hydropower generation, LULCC could also threaten hydropower plants with silting and sedimentation in reservoirs. Moreover, the effect of combined changes on the Bui scheme in the far future, as well as on other existing and future planned hydropower plants, is not well known. The consequences of anthropogenic land-use changes on hydropower have received little attention from policy makers and dam managers in WA, while integrating this land-change information is crucial for planning strategies, for efficient national and regional adaptation policies, and for sustainable development (Lienou 2013). Based on the literature review, the processes associated with the links between LULCC and hydropower are shown in figure 6.

3.3. Nexus between climate, land and hydropower in WA

Based on the literature review, table 1 summarizes the existing studies on the potential impact of climate change and land use/cover dynamics on the main hydropower plants of WA. Apart from the hydropower plants listed in table 1, there are other dams that have not been assessed from the perspective of this study (Férou and Sélingué in Mali, Taabo in Côte d'Ivoire, Kaléta in Guinea, Shiroro, and Jebba in Nigeria); consequently, there are no relevant references identified during this review. Most of the hydropower plants constructed in the WA region are multi-purpose dams used for domestic water supply, irrigation, recreation, and tourism. It is worth noting that there are a few studies on future hydropower

changes under climate change and LULCC, while this sector and the related infrastructure are very important for the sustainable development of African countries. Table 1 reveals that a decrease in production is projected for most hydropower plants (only the main dams are shown here). All the hydropower plants located in the Volta Basin (Bagre and Kompenga in Burkina Faso; Bui, Akosombo, and Kpong in Ghana), Sassandra (Buyo), and Bandama (Kossou) may not be able to fully operate to satisfy the rising energy demand. This may negatively affect all associated socioeconomic sectors. In the Mono basin (Togo-Benin), uncertainties in the projected rainfall did not allow for a robust assessment of the future change in hydropower production of the Nangbeto plant. Nevertheless, the hydropower plants located in the Niger River basin (Kandaji in Niger and Kainji in Nigeria) are likely to observe increases in their future power generation. However, plant managers have to prepare themselves to cope with the growing risk of flooding, which may affect the dam and downstream dwellers, as observed over the last 30 years in Niger and Nigeria, where frequent river overflows regularly flooded homes and infrastructure.

Figure 7 illustrates and summarizes the interactions between climate, land, water, and energy. Below, we describe how, in the existing literature, different interactions are documented in WA:

Land use/cover effects on climate change (1): LULCC are major factors contributing to climate change through GHG emission volumes and carbon sequestration (Devaraju *et al* 2018). LULCC can affect rainfall (Pielke *et al* 2002) and hydroclimate systems (Oki and Blyth 2012). Bamba *et al* (2015) showed a linear relationship between precipitation and NDVI in the Sudanian savannah region of WA over the last three decades. Additionally, it has


Table 1. Impacts of projected climate change and land use/cover dynamics on West African hydropower schemes. Near future refers to the period 2030–2060 while far future stands for the period 2070–2100.

HP plant and installed capacity country	Basin	Operation year	Thematic	Impacts on hydropower generation		Authors
				Near future	Far future	
Nangbeto (65.6 MW) Togo	Mono	1987	CC&WR	Projected trends for cumulated precipitation are null or very moderate and diverge among models by 2050 (AIB and RCP8.5).		(Amoussou <i>et al</i> 2020)
				Change in precipitation in the range of –32.4% and +12% with late rainy seasons (RCP8.5).		(Lamboni <i>et al</i> 2019)
				Change in Nangbeto HPP by –5.73% (RCP4.5) and –8.73% (RCP8.5) due to CC by 2050	Change in Nangbeto HPP by –12.71% (RCP4.5) and –24.45% (RCP8.5) due to CC by 2090	(Obahoundje <i>et al</i> 2021b)
Akosombo (1038 MW) Ghana	Main Volta	1965	CC&HP	Change in Nangbeto HPP by +4.51% (RCP4.5) and –1.31% (RCP8.5) due to CC&IULCC by 2050	Change in Nangbeto HPP by +14.66% (RCP4.5) and –1.01% (RCP8.5) due to CC&IULCC by 2090	
				Only 77% of hydroelectricity is generated under current development conditions.	Only, 30% and 24% of the potential average annual hydroelectricity will be generated by the end of 2100 under intermediate and full development scenarios, respectively.	(McCartney <i>et al</i> 2012)
				Only, 53% and 48% of the potential average annual hydroelectricity will be generated by the end of 2050 under intermediate and full development scenarios, respectively.	Hydropower demand coverage varies from 80% to 85% depending on the scenario (2011–2050).	(Amisigo <i>et al</i> 2014)
Bui (400 MW) Ghana	Black Volta	2014	CC&IULCC&HP	Change in energy generation in the range of –54% to 40% depending on the scenarios.		(Obahoundje <i>et al</i> 2017)
				Hydropower demand coverage varies from 85% to 98% depending on the scenario (2014–2050).		(Amisigo <i>et al</i> 2014)

(Continued.)

Table 1. (Continued.)

HP plant and installed capacity country	Basin	Operation year	Thematic	Impacts on hydropower generation		Authors
				Near future	Far future	
Kompianga (14 MW) Burkina Faso	White Volta		CC&HP	Hydropower demand coverage varies from 75% to 100% depending on the scenario (2011–2050).		(Amisigo <i>et al</i> 2014)
Bagre (16 MW) Burkina Faso	Oti		CC&HP	Hydropower demand coverage varies from 78% to 95% depending on the scenario (2011–2050).		(Amisigo <i>et al</i> 2014)
Kossou (174 MW) Côte d'Ivoire	Bandama	1972	CC&WR	Change in annual flows: –3.33% by 2030s (RCP4.5). –13.33% by 2030s (RCP8.5).		(Soro <i>et al</i> 2019)
Buyo (165 MW) Côte d'Ivoire	Sassandra	1980	CC&WR	Decrease in annual runoff by 27.7% and in ground water recharge 34.2% for the 2050s (A1B).	Decrease in annual runoff by 40.0% and in ground water recharge 45.8% for 2080s (A1B).	(Koua <i>et al</i> 2014)
Manantali (200 MW) Mali	Sénégal	1988	CC&WR	Change in average annual flows: [–8.1%; –33.4%] by 2050s (RCP4.5). [–1.3%; –56.8%] by 2050s (RCP8.5). Change in Senegal River flow: –8% by 2050s (RCP4.5) –16% by 2050s (RCP8.5)		(Bodian <i>et al</i> 2015) (Bodian <i>et al</i> 2018)
Kainji (800 MW) Nigeria	Niger	1968	CC&HP	Change in annual energy generation: +8.72% by 2035 (RCP4.5). +12.81% by 2035 (RCP8.5).	Change in annual energy generation: +8.63% by 2085 (RCP4.5). +24% by 2085 (RCP8.5).	(Oyerinde <i>et al</i> 2016)



been demonstrated that the presence of forest, tall grass, and short grass savanna at the Sahel–Sahara interface tends to decrease the mean summer surface temperature, while afforestation options tend to increase the precipitation substantially for the whole Sahel region of WA (Diba *et al* 2018). The predicted decrease in temperature is associated with strengthening and weakening of the latent and sensible heat flux, respectively, whereas the increase in precipitation is accompanied by an increase in atmospheric moisture and strong interannual variability over the whole WA region (Sylla *et al* 2016, Diba *et al* 2018). Additionally, LULCC may impact regional and global climate through the surface energy budget (Pielke *et al* 2002). Decreases in vegetation cover may increase the land surface temperature because of reduced evapotranspiration. Conversely, forests and natural wetlands tend to decrease the temperatures in Africa. Although LULCC can lead to climate change, it is strongly sensitive to climate.

Climate change effects on land use (2): Voortman (1998) demonstrated that the WA ecological patterns (spatio-temporal distribution of land cover) have drastically changed due to climatic conditions with a latitudinal shift of isohyets towards the south associated with a general reduction in average precipitation. Scheiter and Savadogo (2016) confirmed that climate change is expected to displace large areas of savannah in WA by promoting transitions to wood-dominated vegetation, due to the fertilizing effects of carbon dioxide. Additionally, the uncertainties of rainfed agricultural productivity may lead to an increasingly irrigated agriculture around the reservoirs of dams,

which will consequently add pressure to the water stored for energy generation.

Climate change effects on hydropower energy (3): The energy demand may be affected by climate change (Wilbanks *et al* 2008), particularly by changes in extreme precipitation and temperature, and compound extreme climate events. Hydropower is particularly vulnerable to changes in precipitation and temperature patterns (Oyebande and Odunuga 2010, IPCC 2011, Hamududu and Killingtveit 2012). This review study showed in the previous section that, in the past, WA hydropower plants have experienced the effects of climate change on their stored water levels as well as on their power generation (Olofintoye and Adeyemo 2011, Salami *et al* 2015).

Hydropower energy generation effects on climate (4): Barros *et al* (2011) showed that depending on the size of the dam, it may be a significant source of GHG emissions due to the long-term decomposition of devastated forests and the geographical location that can be inundated by the water dammed up at hydropower plants. It is worth noting that the emissions from hydropower sources in WA countries are ignored in the GHG emissions inventory from the Intergovernmental Panel on Climate Change (IPCC), while this emission source is very important and significant in tropical regions (Fearnside 2015). We did not find any study assessing the impact of hydropower on WA climate, but we found two papers that contributed to documenting how hydropower plant construction can modify the mesoscale hydroclimatic patterns of a given watershed in WA (Amoussou *et al* 2012, Faye *et al* 2015).

Hydropower energy effects on land use/cover (5): Land cover is degraded by the installation of hydropower infrastructure and by infrastructure for energy supply and distribution (Dale *et al* 2011). Deforestation or vegetation devastation for hydropower plant buildings can first contribute to modifying the land surface albedo, which influences the atmospheric circulation system and rainfall, leading to changes in vegetation dynamics (Charney 1975). We did not find any scientific papers assessing the impact of hydropower on land cover and land use in WA, but we found several reports on environmental and social impact assessment of hydropower projects raising this issue, highlighting the changes in the landscape and the extent of flooded areas caused by dam construction (Burgeap Environmental Consultancy 2006, Government of Ghana 2007).

Land use effects on hydropower energy (6): The characteristics of a landscape determine whether hydropower infrastructure can be established (Dale *et al* 2011). Any changes in land cover in the basin may result in changes in precipitation and discharge, which may affect the efficiency of the hydropower plant. For instance, Obahoundje *et al* (2018) showed how LULCC may influence the hydrology of three transboundary basins in WA, and Kouame *et al* (2019) illustrated how LULCC impacted the Kossou hydropower generation.

4. Conclusion

Hydropower is the main renewable energy source for electricity generation in WA and is essential for the mitigation of climate change. The effects of climate change and land use/cover dynamics on hydropower generation in the WA region were assessed and synthesized. It is worth noting that, among the available literature on this specific topic, several papers led by African institutions are not indexed in Scopus or Web of Science. There are also several reports from national/regional/international institutions that are also of interest to this study, but they are only indexed in Google Scholar. To describe the different interactions between climate, land use/cover, water, and hydropower generation in WA, we considered the main findings from papers in both Scopus and Google Scholar. This approach helped to draw the synthesized flowcharts at the end of each section. To achieve the specific aim of this paper, we selected 300 papers from Scopus and 400 papers from Google Scholar only 96 papers issued after the year 2000, with 66 papers both found in Scopus and Google Scholar and only 30 from Google Scholar.

The analysis of existing peer review papers reveals that the effects of climate change on hydropower systems affect almost all existing plants in WA due to the change and variability of precipitation over large areas of the region, and due to the increase in temperature over the entire region. Precipitation variability

and extreme events such as droughts and floods have caused many hydropower plant disruptions over the last 30 years. Future projections agree on the rise in mean temperatures, but precipitation changes are uncertain, and the influence of this change on streamflow in major WA river basins is very uncertain. The change in both variables may affect the runoff and river discharges, depending on the area and land use (ECREEE 2017), and climate change may considerably affect water demand sectors, including hydropower plants within river basins. West African rivers are likely to face severe freshwater shortages, leading to competition among water-use sectors and riparian residents. An increase in temperature will considerably increase crop water and irrigation demand (Sylla *et al* 2018b). Many hydropower plants will have to adjust their operations in response to the extreme water conditions associated with the projected longer-lasting dry spells and heat waves. This will impact not only energy production but also electricity demand, transmission, and distribution (van Vliet *et al* 2016).

In this review, we found that most planned dams are located over regions (south-west of WA) where a significant increase in runoff is projected to be favorable for hydropower production. However, it is not clear how LULCC will evolve in the future, and there are uncertainties in future climate variability. This suggests that there is a need to assess and understand the combined effects of climate change and variability and LULCC in different basins to reduce the risks of local floods and for sustainable management and planning of hydropower production.

It was also shown that apart from climate change, there are additional factors such as water abstraction for food and LULCC, which can affect the amount of water available for hydropower generation. Indeed, WA countries are exposed to LULCC with increases in settlement and agricultural areas and reductions in vegetated areas. These changes are projected to intensify in the future owing to population and socioeconomic growth rates. The literature review showed that LULCC in WA may increase streamflow (in reducing the infiltration rate), which may improve water availability and favor hydropower generation. However, this increase in LULCC may lead to soil erosion and siltation, with the potential to reduce reservoir capacity, increase sediment yields, and raise the riverbed. Silting associated with increases in streamflow can block hydropower water intakes and cause turbine abrasion, and may intensify the risk of floods and threaten hydropower infrastructure. Unfortunately, such risks have not yet been well documented in the WA region.

Due to the Paris Agreement, major investments in hydropower and dam construction projects are planned for the WA region. However, it is crucial to highlight that dam construction can also contribute to changes in land use and land cover and

modify the surrounding climate and precipitation patterns, as well as the hydrological dynamics of the river (Degu *et al* 2011). This can have a significant impact on water availability (Amoussou *et al* 2012, Faye *et al* 2015). The construction of dams, coupled with rising temperatures, may also favor the development of pest species such as water hyacinth, which is harmful to hydropower generation systems. This has already been observed in the Volta River basin, where the construction of multipurpose dams for hydropower and irrigation has reduced the flow of the river system, triggering an increase in the abundance of water hyacinth (Honlah *et al* 2019). Indeed, the water hyacinth can disrupt water abstraction facilities, obstruct hydroelectric turbine intakes, irrigation channels, and watercourses, causing siltation and flooding as well as high water loss through evapotranspiration.

Considering the uncertainties on the future climate and LULCC, improving hydropower modeling and understanding and modeling the complex nature of interlinkages between climate land water, and energy are essential for the efficient management of existing dams, for sustainable use of each resource, and for the viability of future hydropower projects (Hermann *et al* 2012, Howells *et al* 2013). In response to this, research for the development of climate services integrating the climate-land-water-energy and food nexus is crucial to achieving sustainable development goals. In addition to extensive scientific research on the nexus between climate, land, water, and hydropower at the local level, we also suggest that riparian states in the various catchments encourage anti-erosion farming practices that can impact the inflow to reservoirs to mitigate the negative effects of LULCC.

Funding

The research leading to this publication is co-funded by the NERC/DFID 'Future Climate for Africa' programme under the AMMA-2050 project, Grant No. NE/M019969/1 and by IRD (Institut de Recherche pour le Développement; France) Grant Number 'UMR IGE Imputation 252RA5'. The support for the investigation with new multidisciplinary data was possible thanks to the SUSTAINDAM project co-funded by Agence Nationale de la Recherche (ANR, France; Contract n° 400914/00) implemented in the frame of the Belmont Forum collaborative actions on transdisciplinary research for pathways to sustainability.

Data availability statement

The data that support the findings of this study are openly available at the following URL/DOI: https://data.chc.ucsb.edu/products/CHIRPS-2.0/global_daily/netcdf/p05/.

Acknowledgments

The authors thank the Institute of Research for Development (IRD, France) and Institute of Geosciences for Environment (IGE, University Grenoble Alpes) for providing the facility (the RCM Platform) to perform this study at the University Felix Houphouët Boigny (Abidjan, Côte d'Ivoire) and the IT support funded by IRD/PRPT contract. This study was implemented at the national high performance computing center of Côte d'Ivoire (Centre National de Calcul de Côte d'Ivoire, CNCCI). The support for the investigation with new multidisciplinary data was possible thanks to the Belmont Forum collaborative actions on transdisciplinary research for pathways to sustainability.

Conflict of interest

The authors declare no conflict of interest.

ORCID iD

Arona Diedhiou  <https://orcid.org/0000-0003-3841-1027>

References

- Akpoti K, Antwi E and Kabo-Bah A 2016 Impacts of rainfall variability, land use and land cover change on stream flow of the black Volta Basin, West Africa *Hydrology* **3** 26
- Albergel J 1987 Drought, desertification and surface water resources—application to small basins in Burkina Faso/Sécheresse, désertification et ressources en eau de surface—application aux petits bassins du Burkina Faso *IAHS-AISH Publ.* **168** 355–65
- Alkharabsheh M M, Alexandridis T K, Bilas G, Misopolinos N and Silleos N 2013 Impact of land cover change on soil erosion hazard in Northern Jordan using remote sensing and GIS *Proc. Environ. Sci.* **19** 912–21
- Almeida R M *et al* 2021 Climate change may impair electricity generation and economic viability of future Amazon hydropower *Glob. Environ. Change* **71** 102383
- Amisigo B A, McCluskey A and Swanson R 2014 WIDER working paper 2014/033 modeling impact of climate change on water resources and agriculture demand in the Volta Basin and other basin systems in Ghana
- Amoussou E, Awoye H, Vodounon H S T, Obahoundje S, Camberlin P, Diedhiou A, Kouadio K, Mahé G, Houndénou C and Boko M 2020 Climate and extreme rainfall events in the mono river basin (West Africa): investigating future changes with regional climate models *Water* **12** 1–27
- Amoussou E, Camberlin P and Mahé G 2012 Impact de la variabilité climatique et du barrage nambéto sur l'hydrologie du système Mono-Couffo (Afrique de l'ouest) *Hydrol. Sci. J.* **57** 805–17
- Bamba A, Dieppois B, Konaré A, Pellarin T, Balogun A, Dessay N, Kamagaté B, Savané I and Diédhiou A 2015 Changes in vegetation and rainfall over West Africa during the last three decades (1981–2010) *Atmos. Clim. Sci.* **5** 367–79
- Barnieh B A, Jia L, Menenti M, Zhou J and Zeng Y 2020 Mapping land use land cover transitions at different spatiotemporal scales in West Africa *Sustainability* **12** 1–52
- Barros N, Cole J J, Tranvik L J, Prairie Y T, Bastviken D, Huszar V L M, del Giorgio P and Roland F 2011 Carbon

- emission from hydroelectric reservoirs linked to reservoir age and latitude *Nat. Geosci.* **4** 593–6
- Biasutti M 2019 Rainfall trends in the African Sahel: characteristics, processes, and causes *Wiley Interdiscip. Rev. Clim. Change* **10** 1–22
- Bichet A and Diedhiou A 2018a Less frequent and more intense rainfall along the coast of the Gulf of Guinea in West and Central Africa (1981–2014) *Clim. Res.* **76** 191–201
- Bichet A and Diedhiou A 2018b West African Sahel has become wetter during the last 30 years, but dry spells are shorter and more frequent *Clim. Res.* **75** 155–62
- Bissadu K D, Koglo Y S, Johnson D B and Akpoti K 2017 Coarse scale remote sensing and GIS evaluation of rainfall and anthropogenic land use changes on soil erosion in Nasarawa State, Nigeria, West Africa *J. Geosci. Geomatics* **5** 259–66
- Blacksher B, Crocker T, Drucker E and Filoon J 2011 Hydropower vulnerability and climate change *A Framework for Modeling the Future of Global Hydroelectric Resources, Middlebury College Environmental Studies Senior Seminar, Fall* (available at: https://www.academia.edu/download/31038070/globalhydro_final_dm.pdf) (Accessed 10 January 2018)
- Bodian A, Dezetter A and Diop L 2015 Evolution des apports en eau au barrage de Manantali à l'horizon 2050 *Int. J. Water Resour. Dev.* **32** 1–13
- Bodian A, Dezetter A, Diop L, Deme A, Djaman K and Diop A 2018 Future climate change impacts on streamflows of two main West Africa River Basins: Senegal and Gambia *Hydrology* **5** 21
- Burgeap Environmental Consultancy 2006 *Environmental and Social Management Framework & Environmental Audit* vol 1 (available at: <http://documents1.worldbank.org/curated/en/300391468034175129/pdf/E20640VOL1010P1BOX0334127B01PUBLIC1.pdf>)
- Cappelaere B *et al* 2009 The AMMA-CATCH experiment in the cultivated Sahelian area of south-west Niger—investigating water cycle response to a fluctuating climate and changing environment *J. Hydrol.* **375** 34–51
- Chang S 2018 Scoping reviews and systematic reviews: is it an either/or question? *Ann. Intern. Med.* **169** 502–3
- Charney J G 1975 Dynamics of deserts and drought in the Sahel *Q. J. R. Meteorol. Soc.* **101** 193–202
- Cole M A, Elliott R J R and Strobl E 2014 Climate change, hydro-dependency, and the african dam boom *World Dev.* **60** 84–98
- Condie S A and Webster I T 1997 Evaporation from reservoirs *Water Resour. Manage.* **33** 2813–22
- Cotillon S E 2017 *West Africa Land Use and Land Cover Time Series (No. 2017-3004, pp 1–4). US Geological Survey* (available at: <https://eros.usgs>)
- Dale V H, Efroymson R A and Kline K L 2011 The land use-climate change-energy nexus *Landsc. Ecol.* **26** 755–73
- de Souza Dias V, Pereira da Luz M, Medero G and Tarley Ferreira Nascimento D 2018 An overview of hydropower reservoirs in Brazil: current situation, future perspectives and impacts of climate change *Water* **10** 592
- Degu A M, Hossain F, Niyogi D, Pielke R, Shepherd J M, Voisin N and Chronis T 2011 The influence of large dams on surrounding climate and precipitation patterns *Geophys. Res. Lett.* **38** 1–7
- Descroix L *et al* 2009 Spatio-temporal variability of hydrological regimes around the boundaries between Sahelian and Sudanian areas of West Africa: a synthesis *J. Hydrol.* **375** 90–102
- Descroix L *et al* 2018 Evolution of surface hydrology in the Sahelo-Sudanian strip : an updated review *Water* **10** 1–37
- Devaraju N, de Noblet-Ducoudré N, Quesada B and Bala G 2018 Quantifying the relative importance of direct and indirect biophysical effects of deforestation on surface temperature and teleconnections *J. Clim.* **31** 3811–29
- Diba I, Camara M, Sarr A and Diedhiou A 2018 Potential impacts of land cover change on the interannual variability of rainfall and surface temperature over West Africa *Atmosphere* **9** 376
- Didi S R M, Mouhamed L, Kouakou K, Adeline B, Arona D, Houebagnon J C S, Kouadio A K C, Coulibaly H T J, Obahoundje S and Issiaka S 2020 Using the CHIRPS dataset to investigate historical changes in precipitation extremes in West Africa *Climata* **8** 1–26
- Diedhiou A *et al* 2018 Changes in climate extremes over West and Central Africa at 1.5 °C and 2 °C global warming *Environ. Res. Lett.* **13** 1–11
- Djiby S, Bernd D, Dina W and Verah H 2018 WLC17-09-22—water availability and demand under climate change and population growth, in Lake Guiers, Senegal *17th World Lake Conf. (Lake Kasumigaura, Ibaraki, Japan, 2018)* pp 877–9
- ECREEE 2017 *GIS Hydropower Resource Mapping and Climate Change Scenarios for the ECOWAS Region—Technical Report on Methodology and Lessons Learnt for ECOWAS Countries* (ECOWAS Centre for Renewable Energy and Energy Efficiency (ECREEE), Praia, Cabo Verde. All results and maps are published on the ECOWAS Observatory for Renewable Energy and Energy Efficiency—www.ecowrex.org/smallhydro (available at: www.ecowrex.org/smallhydro)
- Favreau G, Cappelaere B, Massuel S, Leblanc M, Boucher M, Boulain N and Leduc C 2009 Land clearing, climate variability, and water resources increase in semiarid southwest Niger: a review *Water Resour. Res.* **45** 1–18
- Faye C, Diop E H S and Mbaye I 2015 Impacts des changements de climat et des aménagements sur les ressources en eau du fleuve Sénégal: caractérisation et évolution des régimes hydrologiques de sous-bassins versants naturels et aménagés *Belgeo. Rev. belge géographie* **4** 1–25
- Fearnside P M 2015 Emissions from tropical hydropower and the IPCC *Environ. Sci. Policy* **50** 225–39
- Funk C *et al* 2015 The climate hazards infrared precipitation with stations—a new environmental record for monitoring extremes *Sci. Data* **2** 1–21
- Galle S *et al* 2018 AMMA-CATCH, a critical zone observatory in West Africa monitoring a region in transition *Vadose Zo. J.* **17** 1–24
- Gardelle J, Hiernaux P, Kergoat L and Grippa M 2010 Less rain, more water in ponds: a remote sensing study of the dynamics of surface waters from 1950 to present in pastoral Sahel (Gourma region, Mali) *Hydrol. Earth Syst. Sci.* **14** 309–24
- Gbobaniyi E *et al* 2014 Climatology, annual cycle and interannual variability of precipitation and temperature in CORDEX simulations over West Africa *Int. J. Climatol.* **34** 2241–57
- Government of Ghana 2007 *Environmental and Social Impact Assessment of the Bui Hydropower Project* (London: Environmental Resource Management (ERM))
- Haddaway N R, Collins A M, Coughlin D and Kirk S 2015 The role of google scholar in evidence reviews and its applicability to grey literature searching *PLoS One* **10** 1–17
- Hamududu B H and Killingtveit Å 2016 Hydropower production in future climate scenarios: the case for Kwanza River, Angola *Energies* **9** 1–13
- Hamududu B and Killingtveit A 2012 Assessing climate change impacts on global hydropower *Energies* **5** 305–22
- Hermann S, Welsch M, Segerstrom R E, Howells M I, Young C, Alfstad T, Rogner H H and Steduto P 2012 Climate, land, energy and water (CLEW) interlinkages in Burkina Faso: an analysis of agricultural intensification and bioenergy production *Nat. Resour. Forum* **36** 245–62
- Honlah E, Yao Segbefia A, Odame Appiah D, Mensah M and Atakora P O 2019 Effects of water hyacinth invasion on the health of the communities, and the education of children along River Tano and Abby-Tano Lagoon in Ghana *Cogent Soc. Sci.* **5** 1–18
- Howells M *et al* 2013 Integrated analysis of climate change, land-use, energy and water strategies *Nat. Clim. Change* **3** 621–6
- IEA 2019 Statistics: global energy data at your fingertips (available at: www.iea.org/statistics/)

- IPCC 2011 2011: summary for policymakers *IPCC Special Report on Renewable Energy Sources and Climate Change Mitigation* ed O Edenhofer *et al* (Cambridge: Cambridge University Press) pp 1–24
- Kabo-Bah A T, Diji C J and Yeboah K A 2018 *Hydropower Generation in West Africa—The Working Solution Manual* (Amsterdam: Academic Press) (<https://doi.org/10.1016/B978-0-12-813016-2.00013-7>)
- Kabo-Bah A, Diji C, Nokoe K, Mulugetta Y, Obeng-Ofori D and Akpoti K 2016 Multiyear rainfall and temperature trends in the Volta River Basin and their potential impact on hydropower generation in Ghana *Climate* **4** 49
- Klutse N A B *et al* 2018 Potential impact of 1.5 °C and 2 °C global warming on consecutive dry and wet days over West Africa *Environ. Res. Lett.* **13** 055013
- Koua T J *et al* 2014 Potential climate change impacts on water resources in the Buyo Lake Basin (Southwest of Ivory Coast) *Int. J. Innov. Appl. Stud.* **8** 1094–111
- Kouame Y M *et al* 2019 Climate, land use and land cover changes in the Bandama Basin (Côte d'Ivoire, West Africa) and incidences on hydropower production of the Kossou Dam *Land* **8** 103
- Kumi N and Abiodun B J 2018 Potential impacts of 1.5 °C and 2 °C global warming on rainfall onset, cessation and length of rainy season in West Africa *Environ. Res. Lett.* **13** 055009
- Lamboni B, Emmanuel L A, Manirakiza C and Djibib Z M 2019 Variability of future rainfall over the Mono River Basin of West-Africa *Am. J. Clim. Change* **08** 137–55
- Le Barbé L, Lebel T and Tabsoa D 2002 Rainfall variability in West Africa during the Years 1950–90 *J. Clim.* **15** 187–202
- Leduc C, Favreau G and Schroeter P 2001 Long-term rise in a Sahelian water-table: the continental terminal in South-West Niger *J. Hydrol.* **243** 43–54
- Lienou G 2013 Integrated future needs and climate change on the River Niger Water availability *J. Water Resour. Prot.* **2013** 887–93
- Mahe G, Lienou G and Descroix L 2013 The rivers of Africa: witness of climate change and human impact on the environment *Hydrol. Process.* **27** 2105–14
- Mahe G, Olivry J C and Servat E 2005a Sensitivity of West-African rivers to climatic and environmental changes: extremes and paradoxes *IAHS-AISH Publ.* 169–77
- Mahe G, Paturel J, Servat E, Conway D and Dezetter A 2005b The impact of land use change on soil water holding capacity and river flow modelling in the Nakambe River, Burkina-Faso *J. Hydrol.* **300** 33–43
- Mamadou I, Gautier E, Descroix L, Noma I, Bouzou Moussa I, Faran Maiga O, Genthon P, Amogu O, Malam Abdou M and Vandervaere J P 2015 Exorheism growth as an explanation of increasing flooding in the Sahel *Catena* **131** 130–9
- McCartney M, Forkuor G, Sood A, Amisigo B, Hattermann F and Muthuwatta L 2012 *The Water Resource Implications of Changing Climate in the Volta River Basin* International Water Management Institute (IWMI Research Report 146) (P O Box 2075, Colombo, Sri Lanka i) (available at: https://books.google.ci/books?hl=fr%26id=K2AVAgAAQBAJ%26oi=fnd%26pg=PR6%26dq=The+Water+Resource+Implications+of+Changing+Climate+in+the+Volta+River+Basin.+%26ots=SCO1M-Uq4Y%26sig=7gogVdHjvRelUL94Kj9A6mEKJTw%26redir_esc=y#v=onepage%26q=TheWaterResourceImplications)
- Moed H F, Bar-Ilan J and Halevi G 2016 A new methodology for comparing Google Scholar and Scopus *J. Informetr.* **10** 533–51
- Moussa I, Faran O, Ambouta J, Sarr B and Descroix L 2009 Les conséquences géomorphologiques de l'occupation du sol et des changements climatiques dans un bassin-versant rural sahélien/the geomorphological consequences of land use and climate change in a rural Sahelian watershed *Secheresse* **20** 145–52
- Munn Z, Peters M D J, Stern C, Tufanaru C, McArthur A and Aromataris E 2018 Systematic review or scoping review? Guidance for authors when choosing between a systematic or scoping review approach *BMC Med. Res. Methodol.* **18** 1–7
- National Center for Atmospheric Research Staff (eds) 2018 The climate data guide: NDVI: normalized difference vegetation index-3rd generation: NASA/GFSC GIMMS *Last Modif. 14 March 2018* (available at: <https://climatedataguide.ucar.edu/climate-data/ndvi-normalized-difference-vegetation-index-3rd-generation-nasagfsc-gimms>)
- Nicholson S 2013 The West African Sahel: a review of recent studies on the rainfall regime and its interannual variability *ISRN Meteorol.* **2013** 1–32
- Nikulin G *et al* 2018 The effects of 1.5 and 2 degrees of global warming on Africa in the CORDEX ensemble *Environ. Res. Lett.* **13** 065003
- Obahoundje S *et al* 2018 Assessment of spatio-temporal changes of land use and land cover over South-Western African Basins and their relations with variations of discharges *Hydrology* **5** 56
- Obahoundje S, Amoussou E, Youan T M, Kouassi L K and Diedhiou A 2021a Multiyear rainfall variability in the Mono river basin and its impacts on Nangbeto hydropower scheme *Proc. Int. Assoc. Hydrol. Sci.* **384** 343–7
- Obahoundje S, Ofosu E, Akpoti K and Kabo-Bah A 2017 Land use and land cover changes under climate uncertainty: modelling the impacts on hydropower production in Western Africa *Hydrology* **4** 2
- Obahoundje S, Youan T M, Diedhiou A, Amoussou E and Kouadio K 2021b Sensitivity of hydropower generation to changes in climate and land use in the Mono Basin (West Africa) using CORDEX dataset and WEAP model *Environ. Process.* **8** 1073–97 (available at: <https://link.springer.com/article/10.1007/s40710-021-00516-0>)
- Oguntunde P and Abiodun B 2013 The impact of climate change on the Niger River Basin hydroclimatology, West Africa *Clim. Dyn.* **40** 81–94
- Oki T and Blyth E 2012 *Land Cover and Land Use Changes and their Impacts on Hydroclimate, Ecosystems and Society* (Plenary paper for the WCRP open science conference, Denver) (available at: ftp://ftp.cgd.ucar.edu/archive/jhurrell/OSC_papers/Oki/WCRP_Land_RevisedDraft03.docx)
- Olofintoye O and Adeyemo J 2011 The role of global warming in the reservoir storage drop at Kainji dam in Nigeria *Int. J. Phys. Sci.* **6** 4614–20
- Osei M A, Ko L, Wemegah D D, Preko K, Gyawu E S and Obiri-danso K 2019 The impact of climate and land-use changes on the hydrological processes of Owabi catchment from SWAT analysis *J. Hydrol. Reg. Stud.* **25** 100620
- Oyebande L and Odunuga S 2010 Climate change impact on water resources at the transboundary level in West Africa: the cases of the Senegal, Niger and Volta basins *Open Hydrol. J.* **4** 163–72
- Oyerinde G, Wissner D, Hountondji F, Odofin A, Lawin A, Afouda A and Diekkrüger B 2016 Quantifying uncertainties in modeling climate change impacts on hydropower production *Climate* **4** 34
- Panthou G, Vischel T, Lebel T, Quantin G, Pugin F, Blanchet J and Ali A 2013 From pointwise testing to a regional vision: an integrated statistical approach to detect nonstationarity in extreme daily rainfall. Application to the Sahelian region *J. Geophys. Res. Atmos.* **118** 8222–37
- Pielke R A, Marland G, Betts R A, Chase T N, Eastman J L, Niles J O, Niyogi D D S and Running S W 2002 The influence of land-use change and landscape dynamics on the climate system: relevance to climate-change policy beyond the radiative effect of greenhouse gases *Philos. Trans. Math. Phys. Dn Eng. Sci.* **360** 1705–19
- Riede J O, Posada R, Fink A H and Kaspar F 2016 What's on the 5th IPCC report for West Africa? *Adaptation to Climate Change and Variability in Rural West Africa* ed Yaro and J Hesselberg (Cham: Springer) pp 7–23
- Ringard J *et al* 2016 The intensification of thermal extremes in west Africa *Glob. Planet. Change* **139** 66–77

- Rowell D P 2012 Sources of uncertainty in future changes in local precipitation *Clim. Dyn.* **39** 1929–50
- Rowell D P, Senior C A, Vellinga M and Graham R J 2016 Can climate projection uncertainty be constrained over Africa using metrics of contemporary performance? *Clim. Change* **134** 621–33
- Saggai S and Bachi O E K 2018 Evaporation reduction from water reservoirs in arid lands using monolayers: Algerian experience *Water Resour.* **45** 280–8
- Salami A W, Mohammed A A, Adeyemo J A and Olanlokun O K 2015 Assessment of impact of climate change on runoff in the Kainji lake basin using statistical methods *Int. J. Water Resour. Environ. Eng.* **7** 7–16
- Scheiter S and Savadogo P 2016 Ecosystem management can mitigate vegetation shifts induced by climate change in West Africa *Ecol. Modell.* **332** 19–27
- Shu J, Qu J J, Motha R, Xu J C and Dong D F 2018 Impacts of climate change on hydropower development and sustainability: a review *IOP Conf. Ser. Earth Environ. Sci.* **163** 0–11
- Sighomnou D *et al* 2013 La crue de 2012 à Niamey: un paroxysme du paradoxe du Sahel ?/the 2012 flood in Niamey: a paroxysm of the Sahel paradox? *Secheresse* **24** 3–13
- Soro G E, Amichiatchi N J M C, Kouassi W F and Goula B T A 2019 Assessment of climate change impacts on Kossou power dam inflows (White Bandama River, Côte d'Ivoire) *Clim. Change* **5** 82–89
- Stanzel P, Kling H and Bauer H 2018 Climate change impact on West African rivers under an ensemble of CORDEX climate projections *Clim. Serv.* **11** 36–48
- Steger K, Fiener P, Marvin-Dipasquale M, Viers J H and Smart D R 2019 Human-induced and natural carbon storage in floodplains of the Central Valley of California *Sci. Total Environ.* **651** 851–8
- Sylla M B, Faye A, Klutse N A B and Dimobe K 2018a Projected increased risk of water deficit over major West African river basins under future climates *Clim. Change* **151** 247–58
- Sylla M B, Pal J S, Faye A, Dimobe K and Kunstmann H 2018b Climate change to severely impact West African basin scale irrigation in 2 °C and 1.5 °C global warming scenarios *Sci. Rep.* **8** 1–9
- Sylla M B, Pal J S, Wang G L and Lawrence P J 2016 Impact of land cover characterization on regional climate modeling over West Africa *Clim. Dyn.* **46** 637–50
- Taylor C M, Belusic D, Guichard F, Parker D, J, Vischel T, Bock O, Harris P P, Janicot S, Klein C and Panthou G 2017 Frequency of extreme Sahelian storms tripled since 1982 in satellite observations *Nature* **544** 475–8
- Tossou E M, Ndiaye M L, Traore V B, Sambou H, Kelome N C, Sy B A and Diaw A T 2017 Characterisation and analysis of rainfall variability in the Mono-Couffo River Watershed Complex, Benin (West Africa) *Resour. Environ.* **7** 13–29
- Tricco A C *et al* 2018 PRISMA extension for scoping reviews (PRISMA-ScR): checklist and explanation *Ann. Intern. Med.* **169** 467–73
- Turner S W D, Ng J Y and Galelli S 2017 Examining global electricity supply vulnerability to climate change using a high-fidelity hydropower dam model *Sci. Total Environ.* **590–591** 663–75
- USAID 2013 *Climate Change and Water Resources in West Africa: Transboundary River Basins* (United States Agency for International Development by Tetra Tech ARD, through a Task Order under the Prosperity, Livelihoods, and Conserving Ecosystems (PLACE) Indefinite Quantity Contract Core Task Order (USAID Contract No. AID-EPP-I-00-06-00008, Order N)
- van Vliet M T H, Sheffield J, Wiberg D and Wood E F 2016 Impacts of recent drought and warm years on water resources and electricity supply worldwide *Environ. Res. Lett.* **11** 124021
- Vitteck M, Brink A, Donnay F, Simonetti D and Desclée B 2013 Land cover change monitoring using landsat MSS/TM satellite image data over west Africa between 1975 and 1990 *Remote Sens.* **6** 658–76
- Voortman R L 1998 Recent historical climate change and its effect on land use in the eastern part of west Africa *Phys. Chem. Earth* **23** 385–91
- Welde K and Gebremariam B 2017 Effect of land use land cover dynamics on hydrological response of watershed: case study of Tekeze Dam watershed, northern Ethiopia *Int. Soil Water Conserv. Res.* **5** 1–16
- Whipple A A and Viers J H 2019 Coupling landscapes and river flows to restore highly modified rivers *Water Resour. Res.* **55** 4512–32
- Wilbanks T J *et al* 2008 *Effects of Climate Change on Energy Production and Use in the United States*
- World Bank 2017 *Water Atlas of Fouta Djallon Highland* (Washington, DC: The World Bank Group)
- Yan B, Fang N F, Zhang P C and Shi Z H 2013 Impacts of land use change on watershed streamflow and sediment yield: an assessment using hydrologic modelling and partial least squares regression *J. Hydrol.* **484** 26–37
- Yang D, Kanae S, Oki T, Koike T and Musiak K 2003 Global potential soil erosion with reference to land use and climate changes *Hydrol. Process.* **17** 2913–28
- Yin J, Gentine P, Zhou S, Sullivan S C, Wang R, Zhang Y and Guo S 2018 Large increase in global storm runoff extremes driven by climate and anthropogenic changes *Nat. Commun.* **9** 4389
- Zarfl C, Lumsdon A E, Berlekamp J, Tydecks L and Tockner K 2015 A global boom in hydropower dam construction *Aquat. Sci.* **77** 161–70
- Zuo D, Xu Z, Yao W, Jin S, Xiao P and Ran D 2016 Assessing the effects of changes in land use and climate on runoff and sediment yields from a watershed in the Loess Plateau of China *Sci. Total Environ.* **544** 238–50