



HAL
open science

Reappraising the Production and Transfer of Hydrogen to the Upper Atmosphere at Times of Elevated Water Vapor

Franck Montmessin, Denis A. Belyaev, Franck Lefèvre, Juan Alday, Anna A. Fedorova, Oleg I. Korablev, Alexander Trokhimovskiy, Mike Chaffin, Nick M. Schneider

► To cite this version:

Franck Montmessin, Denis A. Belyaev, Franck Lefèvre, Juan Alday, Anna A. Fedorova, et al.. Reappraising the Production and Transfer of Hydrogen to the Upper Atmosphere at Times of Elevated Water Vapor. Europlanet Science Congress 2022, Sep 2022, Granada, Spain. pp.EPSC2022-436. insu-03753837

HAL Id: insu-03753837

<https://hal-insu.archives-ouvertes.fr/insu-03753837>

Submitted on 18 Aug 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Reappraising the Production and Transfer of Hydrogen to the Upper Atmosphere at Times of Elevated Water Vapor

Franck Montmessin¹, Denis Belyaev², Franck Lefevre¹, Juan Alday³, Anna Fedorova², Oleg Korablev², Alexander Trokhimovskiy², Mike Chaffin⁴, and Nick Schneider⁴

¹LATMOS, CNRS/IPSL/UVSQ/UPMC, Guyancourt, France (franck.montmessin@latmos.ipsl.fr)

²Space Research Institute of the Russian Academy of Sciences (IKI RAS), Moscow, Russia

³Open University, Milton Keynes, U.K.

⁴LASP, Boulder, Colorado, United States

We have used a 1D hybrid model to represent the ascent of a wet air parcel at times of intense dust and transport activity. This model combines observations of the ACS instrument that measured, for the first time, water vapor abundance from 20 to 120 km. These observations enable the in-depth study of how the water vapor penetration to high altitude contributes to hydrogen production above 80 km. In contrast with other 1D models that have been used to explore Mars' photochemistry, our model represents the vertical transport through advection with a constant velocity of 10 cm/s up to 100 km. Our results imply that, contrary to a common assumption made in models used to study Mars' photochemistry and escape processes, the region between 60 and 80 km cannot be neglected in the production and migration of hydrogen to the upper atmosphere. In particular, these results imply that upper atmosphere photochemistry models intending to capture Southern Summer conditions need to carefully consider the flux boundary condition for H at the lower boundary if it is higher than 80 km. Testing a variety of configurations, from the MY34 GDS to the recent MY35 perihelion period, we have been able to assess how the hydrogen upward flux from above 60 km varies with events. Stochastic events (GDS and A, B, C- storms) have a strong imprint on the escape budget, but our results suggest perihelion remains the dominant escape component on the long term.