



HAL
open science

Sources, sinks, and cycling of dissolved organic copper binding ligands in the ocean

Angel Ruacho, Camille Richon, Hannah Whitby, Randelle M. Bundy

► **To cite this version:**

Angel Ruacho, Camille Richon, Hannah Whitby, Randelle M. Bundy. Sources, sinks, and cycling of dissolved organic copper binding ligands in the ocean. *Communications Earth & Environment*, 2022, 3, 339, p. 12-21. 10.1038/s43247-022-00597-1 . insu-03868827

HAL Id: insu-03868827

<https://insu.hal.science/insu-03868827>

Submitted on 24 Nov 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

Sources, sinks, and cycling of dissolved organic copper binding ligands in the ocean

Angel Ruacho^{1✉}, Camille Richon², Hannah Whitby³ & Randelle M. Bundy^{1✉}

Organic Cu-binding ligands have a fundamental influence on Cu distributions in the global ocean and they complex >99% of the dissolved Cu in seawater. Cu-binding ligands however, represent a large diversity of compounds with distinct sources, sinks and chemical properties. This heterogeneity makes the organic Cu-binding ligand pool difficult to study at the global scale. In this review, we provide an overview of the diversity of compounds that compose the marine Cu-ligand pool, and their dominant sources and sinks. We also summarize the most common analytical methods to measure ligands in marine water column samples. Generally, ligands are classified according to their conditional binding strength to Cu. However, the lack of a common definition for Cu ligand categories has previously complicated data inter-comparison. To address this, we provide a general classification for Cu-binding ligands according to their binding strength and discuss emerging patterns in organic Cu-binding ligand distributions in the ocean according to this classification. To date, there is no global biogeochemical model that explicitly represents Cu ligands. We provide estimates of organic Cu-binding ligand fluxes at key interfaces as first order estimates and a first step for future modeling efforts focused on Cu and Cu-binding ligands.

Copper (Cu) is a widely studied trace metal due to its importance for various biological functions¹. It also plays a role as a toxicant when present in high concentrations of the unbound ion, Cu²⁺²⁻⁴. The cycling of dissolved Cu (Cu) in the oceans, estuaries and rivers has been relatively well-studied over recent years⁵⁻¹³. Recent knowledge acquired largely thanks to the international GEOTRACES program (<https://www.geotraces.org>) and new biogeochemical models has allowed us to gain fresh insights into natural and anthropogenic Cu sources as well as removal processes^{14,15}. However, despite growing datasets on dissolved Cu in seawater and the development of some of the first biogeochemical models for Cu^{9,12,16,17}, many of the internal cycling processes impacting global Cu distributions such as regeneration and scavenging are not well understood. Organic Cu-binding ligands are one aspect of the biogeochemical cycling of Cu that remains to be further explored and likely plays an important role in governing the reactivity of Cu in the ocean. In seawater greater than 99% of dissolved Cu is complexed to organic ligands¹⁸, maintaining Cu in the dissolved phase as well as controlling the concentration of Cu²⁺ and thus playing an important role in Cu bioavailability and toxicity¹⁹. The pool of organic ligands that bind dissolved Cu in seawater consists of a heterogeneous mix of compounds with varying Cu-binding strengths and reactivities^{5,7,11,20-22}. These organic ligands are operationally grouped according to their conditional binding strength ($\log K_{\text{CuL}_i, \text{Cu}^{2+}}$) with the stronger ligands termed “L₁”, followed by L₂, L₃, ..., L_{*n*}. While much work is yet to be done, advances in laboratory experiments and a dramatic increase in Cu-binding ligand measurements in the ocean are beginning to shed light on the sources and sinks of these ligands^{5,7,16,23}.

¹School of Oceanography, University of Washington, Seattle, WA, USA. ²Laboratoire d’Océanographie Physique et Spatiale UBO-CNRS-Ifremer-IRD, IUEM, 29280 Plouzané, France. ³School of Environmental Sciences, University of Liverpool, Liverpool L69 3GP, UK. ✉email: aruacho@uw.edu; rbundy@uw.edu

Despite the recognition of the importance of ligands in dissolved iron (Fe) cycling, relatively little attention has been paid in the literature to Cu-binding ligands and their impact on the oceanic inventory of Cu. For example, it has been established that including a dynamic ligand cycle for dissolved Fe-binding ligands is essential for reproducing the measured in situ datasets of dissolved Fe profiles from GEOTRACES basin-scale sections^{24,25}. However, a similar exercise has not yet been performed for Cu, despite the knowledge that the majority of dissolved Cu in seawater is associated with ligands. This review discusses our current understanding of the biogeochemical cycling of Cu-binding ligands in seawater. We summarize what is known so far about the chemical identity of Cu-binding ligands, including how we measure these ligands using the most commonly employed analytical techniques (voltammetry). Then we provide a general overview of the nature and importance of Cu complexation in marine waters with a focus on how observational work can complement biogeochemical modeling efforts to better understand the role that Cu-binding ligands may play in impacting the reactivity and bioavailability of Cu in the marine environment. Based on a compilation of all water column Cu-binding ligand studies to date ($n = 54$ studies), we find that rivers and sediments are the most important external sources of Cu-binding ligands to the open ocean, while photodegradation and microbial uptake are the dominant sinks in surface waters. Future efforts that characterize fluxes of Cu-binding ligands and their internal cycling are critical for better understanding Cu cycling in the global ocean.

Components of the marine Cu-binding ligand pool

Characterization of Cu-binding ligands is complicated by the broad assortment of chemical compounds contributing to the organic complexation of dissolved Cu in the marine environment. In fact, the ligand pool is most likely a continuum of compounds of varying Cu-binding capacities, with some ligand types themselves exhibiting multiple complexation sites (e.g., humic substances²⁶). As such, in some cases it may not be appropriate to characterize ligands into groups (L_1 , L_2 etc.) defined by their conditional stability constants at all²⁷. However, identifying key ligand groups or key functional groups dominating marine Cu complexation can teach us about the processes that control the distribution and bioavailability of Cu throughout the water column. In oxic seawater, the +II oxidation state of inorganic Cu is thermodynamically favored²⁸, although Cu can be reduced to the +I oxidation state by photochemical and biological processes^{29,30}. Around 10% of the dissolved Cu in ocean surface waters has been shown to be Cu(I)³¹, with potentially as much as 80–90% as Cu(I) in estuarine waters^{11,32}. Ligands containing nitrogen and oxygen-based-binding groups tend to stabilize the +II oxidation state, while ligands with sulfur-binding groups can stabilize the +I state, which acts to reduce Cu^{2+} and relieve Cu toxicity^{33,34}. It is likely that the Cu-binding ligand pool present in seawater is complexing a combination of either Cu(I) or Cu(II), depending on the identity of the ligand.

Several different classes of organic molecules have been identified in seawater that have the capacity to bind Cu, including humic-like substances, exopolysaccharides, thiols and other low molecular weight microbially produced compounds. All of these compounds vary in terms of their overall binding strengths and their impact on Cu cycling. Humic-like substances and exopolysaccharides are components of the dissolved organic matter (DOM) pool that can bind Cu and other metals^{35–37}, likely through carboxylic and nitrogen-containing functional groups³⁸. In the marine environment, these substances are largely derived from marine algae and its decomposition^{39,40}, while terrestrially derived humics contribute to Cu complexation in estuarine and

coastal waters^{11,26,41–43}. Humic-like substances have been found to bind to dissolved Cu with $\log K_{\text{CuL}_i, \text{Cu}^{2+}}$ ranging from 10 to 12^{26,43} while exopolysaccharides bind Cu with $\log K_{\text{CuL}_i, \text{Cu}^{2+}} < 8$ ⁴⁴.

In addition to marine and terrestrial humic-like substances and exopolysaccharides, low molecular weight microbially produced ligands also contribute to the Cu-binding ligand pool in seawater. For example, methanobactins are multidentate ligands that are defined as chalkophores, which have a very high binding affinity for Cu. Chalkophores are thought to be similar to Fe-binding siderophores, in that they have a high specificity for Cu and may be used either to facilitate the uptake of Cu or perhaps to detoxify Cu⁴⁵. Methanobactins have a $\log K_{\text{CuL}_i, \text{Cu}^{2+}} > 14$ ^{46,47}. They are secreted by methanotrophs, which require Cu for various enzymes⁴⁸. Previously thought to be limited to a few key organisms, methanotrophs have now been found in estuaries, open-ocean waters, deep sea sediments, methane seeps and hydrothermal vents^{49–52}. Different species of methanotrophs are also now hypothesized to be ubiquitous within the biosphere, with methanobactin biosynthesis potentially extending beyond methane oxidizers⁵³, suggesting their role in marine Cu complexation may so far be underestimated.

Other microbially produced compounds that are not considered chalkophores, but can also bind Cu with high affinity are likely important Cu-binding ligands in seawater (Table 1). Reduced sulfur substances such as thiols like glutathione, cysteine and thiourea are important for metal detoxification in cell metabolism and have been found to bind Cu in seawater⁵⁴. Phytochelatin, an oligomer of glutathione, is ubiquitously produced by marine algae for metal detoxification, particularly for Cu^{55–57}. Similarly, metallothionein is a family of cysteine-rich, low molecular weight proteins synthesized by marine algae, which also play a role in protection against metal toxicity and oxidative stress⁵⁸. Various thiols and thiol groups have been detected in pore waters, estuaries, coastal and open-ocean waters globally^{11,54,59–65} with $\log K_{\text{CuL}_i, \text{Cu}^{2+}}$ ranging from 11 to 16 depending on the oxidation state of the Cu and the stoichiometry of the ligand complex^{11,34,60,66}.

Other uncharacterized Cu ligands have also been found in marine waters. Recent work on identifying the chemical structures of microbially produced Cu-binding compounds in the southern Pacific Ocean found one major ligand type was a compound of molecular formula $[\text{C}_{20}\text{H}_{21}\text{N}_4\text{O}_8\text{S}_2\text{M}]^+$ ($M =$ metal isotope), containing several azole-like metal-binding groups as well as a sulfur group⁶⁷. Additional ligands were also identified in this study that bound both nickel and Cu. Although the ultimate source of these ligands is unknown, it is likely they are biologically produced⁶⁷.

Hemocyanins are Cu-containing metalloproteins responsible for transporting oxygen in the blood of many marine invertebrates such as cephalopods, mollusks and crustaceans, including copepods and krill. Similar to the role of hemes in the marine Fe cycle^{68,69}, hemocyanins could be potentially important contributors to the marine Cu cycle and Cu-binding ligand pool, but have received little attention in marine Cu studies to date. Various other Cu-complexing compounds have also been identified in marine waters, such as domoic acid⁷⁰. Such specific compounds may be important to Cu bioavailability either on very localized scales or periodically, such as during mass feeding events or during large harmful algal bloom events (e.g., domoic acid). Overall, we have only just begun to unravel the identities of Cu-binding ligands in seawater.

Measuring copper-binding ligands

In this section, we present the most commonly used methods for characterizing Cu-ligand concentrations and binding strengths in

Table 1 Organic copper-binding ligands produced by marine microbes.

Region or microbe	Species	Ligand class	Ligand concentration (nmol L ⁻¹)	logK _{CuI₂,Cu²⁺}	m/z	Molecular formula, functional groups, ID	Method	References
Coccolithophore	<i>Hymenomonas carterae</i>	-	-	-	-	-	ASV	Croot et al. (2000) ¹⁴⁰
	<i>Emiliania huxleyi</i>	L ₁	54	15.6	-	-	CLE-CSV	Leal et al. (1999) ¹⁴²
	<i>Emiliania huxleyi</i>	L ₂	145–160	11.6	-	Arginine-cysteine (Arg-Cys) and glutamine-cysteine (Gln-Cys)	CLE-CSV	Dupont et al. (2004) ¹⁴¹
Cyanobacteria	<i>Synechococcus</i> WH 7803	-	40	-	-	-	ASV	Croot et al. (1999) ¹³⁸
	<i>Synechococcus</i> WH 7803	L ₂	15–110	12.3–13.3	-	-	CLE-CSV	Moffett and Brand (1996) ¹⁹
	<i>Synechococcus</i> PCC 7002	L ₂	139	12.2	-	-	CLE-CSV	Lawrence (1998) ¹³⁹
Dinoflagellate	<i>Amphidinium carterae</i>	-	75	-	-	-	ASV	Croot et al. (1999) ¹³⁸
	<i>Prorocentrum micans</i>	-	-	-	-	-	ASV	Croot et al. (2000) ¹⁴⁰
Diatoms	<i>Skeletonema costatum</i>	-	47	-	-	-	ASV	Croot et al. (1999) ¹³⁸
	<i>Thalassiosira weissflogii</i>	-	1–30	-	-	Glutathione; thiol	HPLC	Tang et al. (2005) ²¹⁸
Heterotrophic bacteria	<i>Vibrio alginolyticus</i>	L ₂	31	11.0	-	-	ASV	Gordon et al. (2000) ⁹⁶
Methane-oxidizing bacteria	-	L ₁	-	-	-	Methanobactins; thioamide groups	ion exchange and hydrophobic interaction chromatography	Zahn and DiSpirito (1996) ²¹⁹
Brown algae	<i>Fucus vesiculosus</i>	L ₂	20–800	11.2	-	-	CLE-CSV	Gledhill et al. (1999) ²²⁰
Algae and Fungi	-	-	-	-	-	Phytochelatins; thiol	HPLC	Ahner et al. (1995) ¹⁴⁵ ; Zenk (1996); Kawakami et al. (2006)
Eukaryotes	-	-	-	-	-	Glutathione; thiol	HPLC, CSV	Dupont et al. (2004); Dupont and Ahner (2005); Kawakami et al. (2006) ²²¹
Southern Equatorial Pacific surface waters	-	-	0.05–0.15	-	572.009	[C ₂₀ H ₂₁ N ₄ O ₈ S ₂ Cu] ⁺ ; azole	LC-ICP/ESI-MS	Boiteau et al. (2016) ⁶⁷

Organic copper-binding ligands that have been observed to be produced by microbes in culture or in natural seawater. The “-” denotes that parameter was not determined.

seawater. Then, we propose new operational definitions of Cu-ligand classes based on measured stability constants and known model ligands, and call for a consensus on defining these ligand pools to facilitate comparisons across datasets and analyses in future studies.

Anodic stripping voltammetry (ASV). The earliest methods to measure the organic complexation of dissolved Cu in seawater used anodic stripping voltammetry (ASV)^{71,72}. This method involves titrating a natural sample with dissolved Cu, and measuring the “ASV-labile” Cu pool over the course of the titration. The ASV-labile Cu pool is predominantly a mixture of inorganic Cu complexes (Cu⁰), which includes any free or hydrated Cu²⁺ and Cu(CO₃)⁰. However, ASV-labile Cu could also include organically complexed Cu (CuL) if the complexes are not kinetically stable. This method is a direct speciation technique, in that the sample is not perturbed beyond the addition of dissolved Cu during the titration, and the labile Cu is measured directly. In this method, the labile Cu⁰ is reduced to Cu⁰ and forms a Cu-mercury amalgam over a certain deposition time. The Cu⁰ is then oxidized (“stripped”) out of the mercury and the current is measured by the instrument. The peak heights are plotted against the total Cu in the sample (Cu_T = dissolved Cu + added Cu) producing a titration curve. Kinetically stable Cu is not electrochemically active, and thus only labile Cu⁰ will adsorb to the mercury drop at the deposition potential (voltage) used in ASV measurements. When the total dissolved Cu in a sample is calculated, the difference between the total dissolved Cu and ASV-labile Cu is said to equal the strongly organically complexed Cu. The ligand concentration (L) and binding strength (logK_{CuL_i,Cu²⁺}) can be calculated based on the analytically measured current, and different linearization techniques^{73–75}. However, any species that dissociate at the applied plating potential will contribute Cu to the labile Cu peak, including known Cu-binding ligands such as thiols, therefore ASV measurements can overestimate labile Cu and thus underestimate ligand concentrations.

An important aspect of ASV is the analytical window, or the detection window of the method. The analytical window is defined by α , or a side reaction coefficient. Since ASV detects only labile Cu, which only includes inorganic Cu, then the α for ASV is approximately equal to the inorganic side reaction coefficient for Cu, or $\alpha_{Cu} = \frac{\sum[CuX_i]}{[Cu^{2+}]}$, where $\sum[CuX_i]$ is the sum of all inorganic Cu complexes. Thus, the α for ASV $\sim \alpha_{Cu}$, which is equal to 10–20 depending on the salinity and pH of the seawater⁷⁶. The analytical window is often expressed as a log value, so for ASV the log α for pH 8.2 seawater is 1.3^{76,77}. Techniques with higher analytical windows will be able to measure stronger organic Cu complexes while lower analytical windows will target weaker Cu complexes. The side reaction coefficient of the natural ligands in a sample is defined by $\alpha_L = K_L \times [L]$ where K_L is the conditional stability constant of the natural ligands and $[L]$ is the concentration of ligands not already bound to Cu. When α_L is within an order of magnitude on either side of the α of the chosen analytical method, those ligands will be detected, while those outside this window will not be detected accurately⁷⁸. The ASV method has a very low analytical window since it only measures labile Cu complexes (inorganic Cu), and thus studies using ASV can generally only measure weak Cu-binding ligands⁷⁷.

Competitive ligand exchange-adsorptive cathodic stripping voltammetry (CLE-ACSV). More recently, the most common methods for understanding the Cu-binding ligand pool in seawater use a variation on the ASV technique and were first

employed in the 1980s^{79–81}. These methods are called competitive ligand exchange-adsorptive cathodic stripping voltammetry (CLE-ACSV) as described for Cu in Campos and van den Berg¹⁸. With this method, aliquots of a filtered seawater sample are titrated with increasing concentrations of dissolved Cu, in the presence of a buffer to maintain a constant pH (usually near 8). A well-characterized artificial ligand is then added to compete with the natural ligands for dissolved Cu complexation. It is recommended that the dissolved Cu additions range from +0 to at least 10x the ambient dissolved Cu concentration with at least two initial +0 nmol L⁻¹ Cu amendments, across 10 or more titration points (ideally 15 points)^{82,83}. After a period of equilibration (15 min to overnight^{5,18,20,84}), each sample aliquot of the titration is analyzed sequentially using ACSV, whereby a hanging mercury drop electrode is used to apply a potential during the deposition step, in which the Cu and added ligand (AL) complex adsorb onto the mercury drop while the solution is stirred. The potential is then scanned in a negative direction and the reduction current of the Cu–AL complex is measured. This is repeated for each aliquot and the height of this peak is recorded and plotted against the concentration of dissolved Cu, ultimately producing the titration curve. This titration data is then interpreted using either linear^{81,85,86} or non-linear⁸⁷ transformations to determine the ambient ligand concentration [L_i] (where *i* denotes ligand class) and binding strengths ($K_{CuL_i,Cu^{2+}}$ or K_{CuL_i,Cu^0}) of the natural Cu-binding ligand pool. Conditional stability constants for these ligand classes are expressed as either $K_{CuL_i,Cu^{2+}}$ or K_{CuL_i,Cu^0} , where $K_{CuL_i,Cu^{2+}} = K_{CuL_i,Cu^0} \times \alpha_{Cu}$. The technique described above is known as a “forward” titration, but Nuester and van den Berg⁸⁸ describe a “reverse” titration in which the AL is spiked in increasing concentrations while the dissolved Cu remains constant. The “reverse” titration technique is employed in circumstances when high dissolved Cu concentrations could result in the [dissolved Cu] > [L] such as around hydrothermal sites or in some coastal environments⁸⁹. Both techniques have their advantages; the “forward” titrations are more commonly employed for determining Cu-binding ligand parameters.

Similar to ASV, the identity of the AL and the AL concentration together determine the analytical window, which in turn controls the range of binding strengths of the ligands that may be detected. The analytical window of the chosen AL is defined by $\alpha_{Cu(AL)_x} = K_{AL} \times [AL]^x$ and $[AL]^x$ is the concentration of the AL not bound to Cu, raised to the power of the stoichiometry of the Cu–AL complex^{78,90}. In CLE-ACSV the [AL] \gg dissolved Cu, so effectively the [AL] = [AL]. The resulting ligand concentrations and conditional stability constants measured with this technique thus represent an average of the different ligands that are detected within the analytical window applied. Depending on the AL, the $\alpha_{Cu(AL)_x}$ can vary several orders of magnitude⁷⁷, generally from log $\alpha_{Cu(AL)_x}$ 2.1–6.5⁷⁷. These methods are useful in that they allow calculation of the ligand parameters and resulting Cu²⁺ concentrations with minimal alteration of the natural sample, similar to ASV. However, aside from the logK_{CuL_i,Cu²⁺}, they do not provide information on the identity of the compounds that might make up the organic Cu-ligand pool in seawater, and are restricted by the detection window. An additional important consideration for these methods is the amount of dissolved Cu that is exchangeable with the added ligand. Findings from Moriyasu and Moffett demonstrate that dissolved Cu in older deep waters is primarily found in an inert form, with implications for speciation calculations⁹¹. When dealing with samples with a high percentage of inert Cu, an overestimation of a ligand’s conditional stability constant will occur unless the inert fraction of dissolved Cu is omitted from the total dissolved Cu used in speciation calculations.

Direct quantification of ligand groups using voltammetry. Complementary voltammetric analyses can provide additional information on the identity of Cu-binding ligands by quantifying the electrochemical response of specific Cu-binding ligand groups within a sample, such as electroactive humic substances and thiols⁹². Here, the voltammetric peaks corresponding to the organic compounds being measured are quantified by additions of commercial standards (e.g., thiols such as glutathione and thioacetamide, and humic or fulvic acid isolates, typically from the Suwannee River (SRHA or SRFA)). These compound concentrations are then converted to a corresponding Cu-binding concentration using a ligand-binding ratio with Cu that has been measured independently for the standard being used. One caveat for this technique is that it is assumed the Cu-binding capacities of the natural ligands are the same as the standards and the method cannot discriminate the target ligand from other similar compounds present in the natural sample. For example, exopolymeric substances can be confused with humic substances, or multiple thiol peaks can coalesce and include sulfide, due to the compounds having similar electrochemical behaviors. These methods are continually being developed and improved⁹³ and they are fast and have low detection limits, allowing for a deeper insight into the key classes of Cu-binding ligands present in seawater than CLE-ACSV alone can provide.

Additional methods for characterizing the Cu-ligand pool. In addition to voltammetric techniques, several additional methods for characterizing the Cu-binding ligand pool in seawater and pore waters have also been used. These methods include liquid chromatography coupled to inductively coupled plasma mass spectrometry and electrospray ionization mass spectrometry (LC-ICP/ESI-MS) or Fourier-transform ion cyclotron resonance mass spectrometry (FT-ICR-MS), and Cu(II)-immobilized metal affinity chromatography (IMAC). Mass spectrometry techniques like LC-ICP/ESI-MS or FT-ICR-MS are used on samples that have first been extracted from seawater onto a column using solid phase extraction, and then these compounds are eluted in methanol or another solvent and the compounds are then separated using reverse phase chromatography coupled to a mass spectrometer^{67,94}. Standards are generally not commercially available for the natural microbially produced Cu-binding ligands that might be found in seawater, so quantifying these compounds via mass spectrometry techniques is a challenge. However, some putative compounds have been quantified using LC-ICP-MS, where Cu-containing compounds are first identified by peaks in ⁶³Cu on the ICP-MS, and then these peaks are quantified and matched to their mass to charge ratio (*m/z*) found using LC-ESI-MS⁶⁷.

In the Cu(II)-IMAC method, the IMAC column is first loaded with Cu²⁺ ions and then the sample is passed over the column to capture organic compounds with an affinity for Cu⁹⁵. These compounds are then eluted and measured either via UV-adsorption or mass spectrometry^{96,97}. This method is promising in that it is able to isolate only Cu-binding ligands while other solid phase extraction techniques capture a myriad of organic compounds. Both the Cu(II)-IMAC method and the mass spectrometry methods that have been employed thus far to identify and characterize the organic Cu-binding ligand pool in seawater have the potential to rapidly accelerate our understanding of marine Cu complexation, but very few published studies exist^{95–99}. Importantly, new work also suggests that a large fraction of dissolved Cu (up to 90%) in deep waters may be inert, which has implications for these analytical methods as well as for dissolved Cu reactivity and bioavailability⁹¹. Future work that aims to characterize Cu-binding ligands in the marine

environment will be particularly insightful to our understanding of the marine Cu cycle.

A call for an operational consensus on copper ligands.

Although many methods have been used to characterize organic Cu speciation, we largely focused our discussion here based on lessons learned from voltammetry Cu speciation studies, which are the methods that have been employed for the bulk of the GEOTRACES studies and many other basin-scale datasets. All of the voltammetric studies that have been completed in the water column to our knowledge are summarized in Table 2 (*n* = 54 studies). These studies are those that only focused on measuring Cu-binding ligands in the water column of estuarine, coastal and open-ocean areas, excluding studies from specific sources such as hydrothermal vent fluids, pore waters, freshwater, or culture studies. Ligands characterized through voltammetric methods are usually categorized as either “strong” (L₁) or “weak” (L₂ or L₃) based on the measured $\log K_{\text{CuL}_i, \text{Cu}^{2+}}$, but the measured stability constants span several orders of magnitude ($\log K_{\text{CuL}_i, \text{Cu}^{2+}} = 7.5\text{--}16.5$; Table 2) and there is no consensus in the literature for Cu-binding ligands on what constitutes a L₁, L₂, or L₃ ligand for Cu. This is complicated by the fact that different analysts use different analytical windows, which influences the range of ligands that can be detected accurately by a given method. For organic Fe-binding ligands, Gledhill and Buck¹⁰⁰ proposed that ligand classes should be consistently operationally defined between studies based on their stability constants, where the strongest L₁ ligands are defined as ligands with $\log K_{\text{FeL}_i, \text{Fe}^{3+}} > 12$. There are known model ligands in seawater that fall within this operational class such as siderophores, which are compounds that bind Fe with a similar or greater binding strength than this operational definition. These operational definitions allow identified compounds to be connected with a particular ligand class, and enable comparisons across studies or distinct analytical windows.

However, much less is known about Cu-binding ligands in seawater and what compounds might comprise both the strong and weak Cu-binding ligand pool, and many studies only define a single ligand class (Table 2). Based on the few model Cu-binding ligands that we know of, we can start to operationally define some ligand classes. For example, the model chalkophore methanobactin binds to Cu with conditional binding strengths of 14–16^{46,101}. Some of the strongest Cu-binding ligands observed in seawater have $\log K_{\text{CuL}_i, \text{Cu}^{2+}} > 14.0$. Kleint et al.¹⁰² report $\log K_{\text{CuL}_i, \text{Cu}^{2+}}$ of 14.1 in shallow hydrothermal vents¹⁰² and $\log K_{\text{CuL}_i, \text{Cu}^{2+}}$ of 14–16 have been reported in coastal waters around Scotland and in the NE Pacific^{5,42}. Several additional studies in the Pacific^{5–7,89,103,104}, North Atlantic^{16,78,80,105,106}, and Southern Ocean^{21,107} also all report stability constants exceeding 14.0 (Table 2). With this available observational data and the knowledge that model chalkophores fit within the range of binding strengths observed in seawater, we propose that L₁ ligands should be classified as ligands with stability constants ≥ 14.0 . The majority of the Cu speciation studies compiled in Table 2 report ligands with $\log K_{\text{CuL}_i, \text{Cu}^{2+}}$ ranging from 9.0 to 15.5 (*n* = 24 studies), which likely represent an average of both strong and weak ligands. Model weaker ligands such as humic substances have $\log K_{\text{CuL}_i, \text{Cu}^{2+}} = 10.0\text{--}12.0$ ²⁶, while thiols have a range of $\log K_{\text{CuL}_i, \text{Cu}^{2+}}$ values from 11.0 to 16.0^{34,66}. Finally, another model weaker ligand domoic acid has a $\log K_{\text{CuL}_i, \text{Cu}^{2+}} = 10.3$ ⁷⁰. Based on the range of observed stability constants of these known model ligands, and the prevalence of weaker ligands detected in voltammetric studies (Table 2) we recommend ligands with $\log K_{\text{CuL}_i, \text{Cu}^{2+}} 10.0\text{--}14.0$ should be operationally defined as L₂

Table 2 All voltammetry studies on organic copper-binding ligands.

Region	Depth (m)	Filter (µm)	[Cu] (nmol L ⁻¹)	Method	[L ₁], [L ₂], [L ₃] (nmol L ⁻¹)	$\frac{\log K_{CuL_1, CuL_2, CuL_3}}{\log K_{CuL_1, CuL_2, CuL_3}}$	References
North Pacific n = 13	Surface	0.45	<31	ASV	<30	-	Srma et al., 1980 ²²²
	Surface	0.45	4.25-23.2	ASV	1.25-22.20	-	Nilsen and Lund, 1982 ^{223,c}
	Surface, deep	0.3	0.62	ASV	4.1-5.13	-	Donat et al., 1986 ²²⁴
	0-1400	0.3	0.57-2.19	ASV	0.62-2.4, L ₁	10.42-12.24	Coale and Bruland, 1988 ⁷¹
	0-500	0.3	0.5-1.5	ASV	0.63-10.20, L ₂	8.24-9.24	Coale and Bruland, 1990 ⁷²
	0-3000	0.2	1.5-4.5	CSV	0.74-2.64, L ₁	10.21-12.24	Moffett and Dupont, 2007 ¹⁰³
	0-3000	0.45	0.62-3.7	CSV	0.20-10.29, L ₂	8.07-9.95	Buck et al., 2012 ¹⁰⁴
	Surface and near bottom	0.2	24.3-43.1	CSV	2.5-6.5	12.7-14.1	Li et al., 2015 ²²⁵
	0-1400	0.2	1.2-3.3	CSV	1.54-6.10	12.84-15.11	Whitby et al., 2018 ⁵
	0-58	0.2	0.6-4.7	CSV	22.7-53.2, L ₁	12.7-13.6	Wong et al., 2019 ⁶
0-2000	0.2	0.6-2.5	CSV	67.6-321, L ₂	15.0-16.5	Wong et al., 2021 ²²⁶	
South Pacific n = 2	Surface	0.2	2.9-124.8	Reverse CSV	1.2-4.0, L ₁	13.0-16.2	Wiwit et al., 2021 ²²⁷
	Surface	0.2	1.14	CSV	1.1-20.1, L ₂	15.6-16.5	
	0-1000	0.2	0.26-1.75	CSV	0.7-6.9, L ₁	14.0-15.7	
	0-5000	0.2	0.22-5.33	CSV	2.9-31, L ₂	12.0-13.3	
	Surface	0.45	3.14-9.44	ASV	0.9-4.5, L ₁	13.2-14.5	
	0-140	0.22	8.18-25.97	ASV	2.8-27, L ₂	11.7-13.2	
	Surface	-	3	ASV	3-121	12.29-14.53	
	Surface	-	2.3	ASV	1.4-2.0, L ₁	12.5-14	
	0-3000	0.45	1.2-5.9	ASV	7.7-11.3, L ₂	13.8-14.0	Jacquot et al., 2013 ⁸
	Surface, deep	0.45	1.3-3.5	ASV	1.55-6.46	15.6-16.5	Ruacho et al., 2020 ⁷
North Atlantic n = 14	Surface	0.45	0.92-3.2	CSV	0.75-8.88	12.29-14.53	Duinker and Kramer, 1977 ²²⁸
	Surface	0.2	0.9-2	CSV	7-50	10.5-15.9	Hasle and Abdullah, 1981 ²²⁹
	Surface	0.2	12.7-27.9	ASV	0.3-9.59	11.9-13.2	Piotrowicz et al., 1982 ²³⁰
	Surface	0.2	40.8	ASV	0.615	>12	van den berg, 1984 ⁷⁹
	Surface	0.2	31.1-92.0	ASV	11, L ₁	Mean 8.8	Buckley and van den berg, 1986 ²³¹
	Surface	0.2	12.6-13.6	ASV	33, L ₂	14.3	Kramer, 1986 ²³²
	Surface	0.2	12.6-13.6	ASV	23-488	7.56-10.3	Donat and van den berg, 1992 ²³³
	Surface	0.2	12.6-13.6	ASV	50-821	10.5-15.9	Moffett, 1995 ²³⁴
	Surface	0.2	12.6-13.6	ASV	1.2-73	11.9-13.2	Kozelka and Bruland, 1998 ²³⁵
	Surface	0.2	12.6-13.6	ASV	1.4-4.5	Mean 7.7	Croot et al., 1999 ^{138,b]}
South Atlantic n = 1	Surface	0.2	12.6-13.6	ASV	16-38, L ₁	14.3	Bruland et al., 2000 ⁷⁷
	Surface	0.2	12.6-13.6	ASV	15-40, L ₂	8.8-9	Bruland et al., 2000 ⁷⁷
	Surface	0.2	12.6-13.6	ASV	54-100, L ₃	11.5-15.6	Bruland et al., 2000 ⁷⁷
	Surface	0.2	12.6-13.6	ASV	11.5, L ₁	10.6-11.5	Oldham et al., 2014 ¹⁰⁵
	Surface	0.4	1.4-19.2	ASV	48.6-104, L ₂	14.0-15.1	Jacquot et al., 2015 ¹⁶
	Surface	0.2	0.43-3.07	ASV	13-16, L ₁	12.6-14.1	Arnone et al., 2022 ¹⁰⁶
	Surface	0.2	0.07-4.03	ASV	13-15, L ₂	13.4-14.4	Zitoun et al., 2021 ²³⁶
	Surface	0.2	0.5-3.5	ASV	1.4-4.5	11.6-14.6	Mellet and Buck, 2020 ¹⁸¹
	Surface	0.2	1.2-6.0	ASV	5.7-23.2, L ₁	10.7-12.1	Donat and van den berg, 1992 ²³³
	Surface	0.45	1.96-3.24	ASV	9-17.6, L ₂	12.1-15.7	van den Berg et al., 1987 ⁸⁰
Indian Ocean n = 1	Surface	0.45	10.7-152	ASV	4.13-17.8	12.1-15.7	
	Surface	0.45	10.7-152	ASV	26-206, L _T (single fit)	11.7-12.78	

Table 2 (continued)

Region	Depth (m)	Filter (µm)	[Cu] (nmol L ⁻¹)	Method	[L ₁], [L ₂], [L ₃] (nmol L ⁻¹)	logK _{CuL₁} , Cu ²⁺ , logK _{CuL₂} , Cu ²⁺ , logK _{CuL₃} , Cu ²⁺	References
n = 17							
	Surface	0.45	45–48	CSV	23–152, L ₁ (2ligand fit) 44–131, L ₂ (2ligand fit)	13.2–14.9 11.5–12.8	Donat et al., 1994 ²³⁷
	Surface	0.2	3.8–117	CSV	13, L ₁ 60–80, L ₂	13.5 9.0–9.6	Moffett et al., 1997 ¹¹⁹
	Surface	0.2	7.4–27.9	ASV	10.5–33.3, L ₁ 39.6–135.7, L ₂	12.57–14.3 9.94–11.72	Wells et al., 1998 ²³⁸
	Surface	0.2	7–23.6	ASV	16–107, L ₁ 40–100, L ₃	Exceeded window 8.2–10.5 8.8–9.0	Laglera and van den berg, 2003 ⁵³
	Surface	0.2	81.9–190.1	CSV	9.6–32.5, L ₁ 13.7–299, L ₂	14.51–16.14 12.7–13.46	Dryden et al., 2004a ²³⁹
	Surface	0.2	2.9–22.7	CSV	66.6–160.7	12.5–12.9	Shank et al., 2004 ¹²⁶
	Surface	0.45	17.9–49.9	CSV	7–229	13.5	Buck and Bruland 2005 ²⁰
	Surface	0.2	5.6–6.4	CSV	22–265, L ₁ 55–135, L ₂	13.1–14.3 12–12.9	Shank et al., 2006 ^{173a}
	Surface	0.2	13.1–24.7	CSV	116–132	13.5	Dryden et al., 2007 ⁶⁰
	0–115	0.2	4.08–6.12	CSV	26–56	11.8–12.3	Jacquot et al., 2014 ²⁴⁰
	Surface	0.2	11.6–54.5	CSV	2.98–7.23	12.9–15.6	Whitby and van den Berg, 2015 ⁴³
	Surface	0.2	2.9–65	CSV	21.9–57.5	12.2–13.8	Whitby et al., 2017 ¹¹
	0–40	0.2	2.03–2.65	CSV	16–51, L ₁ 30–112, L ₂	14.0–15.2 12.1–13.4	Wong et al., 2018 ²⁴¹
	Surface	0.02	3–77	ASV	1.95–4.21, L ₁ 6.3–33, L ₂	13.7–16.0 11.1–12.8	Dulaquais et al., 2020 ⁴¹
	Surface	0.45	5–82	ASV	6–120, L ₁ 8–147, L ₂	9.4–11.4 11.4–13.4	Padjan et al., 2021 ¹⁵
	Surface	0.22	5–50	ASV	3.5–63, L ₁ 17–104, L ₂	9.6–10.8 8.2–9.0	Hollister et al., 2021 ¹⁰
Southern n = 5	Surface	0.015	0.7–283	CSV	1.3–66.3, L ₁ 2.8–83.2, L ₂	13.2–15.3 12.6–14.2	Batley and Florence, 1976 ²⁴²
	Surface	0.45	3.93–8.81	ASV	1.57	-	Buck et al., 2010 ^{107a}
	180	0.4	2.16–2.4	CSV	2.5–4.13, L ₁ 0.3–1.8, L ₂	15.7–16.4 14.6–15.5	Bundy et al., 2013 ²¹
	Surface	0.4	1.33–2.84	CSV	2.26–8.04, L ₁ 2.54–8.32, L ₂	13.57–16.0 12.73–13.33	Thompson and Ellwood, 2014 ¹²
	0–3500	0.2	0.32–3.41	CSV	1.08–5.22	12.00–13.64	
	Surface	0.2	0.47–0.55	CSV	1.5–2.5, L ₁ 7.4–7.76, L ₂	12.64–13.57 11.2–11.72	Heller and Croot, 2015 ⁸⁴
	0–3000	0.2	0.85–2.66	CSV	-	>13	
				CSV	23.6–73.3, L ₂	11.1–12.14	

All published voltammetry (ASV, CLE-ASV) studies to date that characterized organic Cu-binding ligands in the water column.
^aDenotes a study that measured water column samples from photochemical or biological incubation experiments.
^bIncludes studies that measured model Cu-binding ligands.
^cStudies that performed an acidification and a digestion.
^d“.” denotes samples where a particular ligand class was not detected or reported.

ligands. There are no known model ligands with binding strengths less than 10.0, though several studies have identified ligands in seawater with weaker binding constants (Table 2) and freshwater algae have been shown to exude exopolysaccharides with $\log K_{CuL_i, Cu^{2+}} < 10.0^{44}$. We therefore recommend that any ligands that are characterized with a $\log K_{CuL_i, Cu^{2+}} < 10.0$ should be classified as L₃ ligands (Table 3). By operationally defining these ligand classes based on the available observational data and known model Cu-binding ligands, we hope future studies will be able to discern emerging patterns in strong and weak Cu-binding ligand cycling.

Sources, sinks, and internal cycling

Sources of Cu-binding ligands

Rivers and margins. Rivers and estuaries are important margin sources for organic Cu-binding ligands. In estuaries, dissolved Cu concentrations typically show conservative behavior, governed by the mixing of high dissolved Cu river water with lower dissolved Cu seawater (Fig. 1)^{108–115}. Deviations from the conservative trend have also been observed, as a result of sorption to solids, colloidal flocculation and/or biological uptake^{41,108,110,116,117}, or Cu inputs along the estuary²⁰. Anthropogenic activity can enhance the supply of dissolved Cu through the contamination of aerosols, rivers and groundwater, particularly in coastal areas^{15,118–120}.

The dissolved Cu in rivers and estuaries is largely organically complexed (>99%), and therefore organic Cu-binding ligands play an important role in the cycling of dissolved Cu in estuarine systems (Fig. 1). For example, humic substances bind strongly to dissolved Cu, and the flocculation of dissolved Cu can occur when humic substances mix with seawater^{121,122}, although the stabilization of dissolved Cu-humic complexes may also mitigate this removal process¹²³. Flocculation during estuarine mixing removes 20–40% of the dissolved Cu^{59,124}, and up to 97% of the ligands¹⁰, reducing dissolved Cu and Cu-ligand concentrations to the levels typically found in coastal waters^{10,60,125,126}. The Cu-binding ligand concentrations are also generally conservative along the salinity gradient in most studies (Fig. 1). However, few studies have examined the true riverine endmember, so it is possible that non-conservative mixing occurs in the low salinity end of estuaries (Fig. 1), possibly due to flocculation of a colloidal fraction of ligands and dissolved Cu¹⁰. Data from Buck and Bruland (2005)^{20,127,128} for example, found much higher Cu-ligand concentrations at the low salinity end of the estuary compared to dissolved Cu, and the excess Cu-binding ligands (excess ligand = [ligand] - [dissolved Cu]) decreased as salinity increased (Fig. 1). Whether there are more sources of Cu-binding ligands in freshwater, or Cu ligands are preferentially scavenged relative to dissolved Cu, is unknown.

Anthropogenic activities that lead to elevated dissolved Cu in rivers and the margins may also trigger ligand production in the estuary. Cyanobacteria such as *Synechococcus* are particularly sensitive to Cu toxicity even at very low levels of free Cu¹⁹. To mitigate the effects of Cu toxicity, *Synechococcus* have been shown to produce strong Cu-binding ligands and release them into solution to mediate Cu toxicity and alter the bioavailability to the in situ dissolved Cu¹⁹. This process might be an important source of Cu-binding ligands in coastal environments, particularly where total dissolved Cu is high and cyanobacteria are abundant. Anti-fouling paint itself has also been found to be a source of Cu-binding ligands in addition to dissolved Cu¹²⁹.

Operationally defined ligand class	$\log K_{CuL_i, Cu^{2+}}$
L ₁	>14.0
L ₂	10.0–14.0
L ₃	<10.0

Proposed operational definitions of Cu-binding ligand classes in seawater using CLE-ACSV methods.




Fig. 1 Dissolved copper and organic ligands in estuarine systems. Data from select studies showing typical **A** dissolved Cu concentrations (x) and organic Cu-binding ligand concentrations (circles), and along the river to seawater continuum. **B** Concentrations of electroactive humic substances expressed in terms of their potential Cu-binding equivalent concentration, based on values calculated for SRHA in Whitby et al.⁴³. Amazon, Mersey, and Sapelo data use the CuHS technique⁴³; San Francisco Bay, Fundao Dam, and Irish Sea use the FeHS technique^{243–245}, while Loire Estuary uses the MoHS technique⁹²; different methods have been shown to give comparable eHS concentrations^{35,246}.

Sediments. Shelf sediments are another important source of Cu-binding ligands to the ocean^{7,21,84} and can be a source of L₁ ligands^{7,21}. In the Equatorial Pacific, strong L₁ ligands ($\log K_{\text{CuL}_i, \text{Cu}^{2+}} > 14.0$) were detected off the coast of Peru, which are some of the strongest ligands that have been detected from shelf sources. These ligands corresponded with a plume of ²²⁸Radium (²²⁸Ra) from the continental shelf^{7,130}. Bundy et al.²¹ detected intermediate L₂ ligands ($\log K_{\text{CuL}_i, \text{Cu}^{2+}}$ 10.0–14.0), from the Antarctica Peninsula shelf and Heller and Croot⁸⁴ similarly detected intermediate ligands in this region. The variability in binding strengths of detected ligands is likely due, in part, to differences in the analytical window and data processing method used by each study^{127,128}, but could also be due to differences in the type of ligands associated with sediment sources. Studies in estuarine and coastal marine sediments have reported multiple ligand classes, with one study in a shallow lagoon finding two types of ligands, with $\log K_{\text{CuL}_i, \text{Cu}^{2+}}$ 14.2 and 12.5, which were associated with thiol compounds⁵⁹. Another coastal study also found one relatively stronger and one relatively weak ligand, but both with $\log K_{\text{CuL}_i, \text{Cu}^{2+}} < 10.0$, which, with our new operational definition would thus be characterized as L₃ ligands¹³¹. Additional observations are needed to gain a better understanding of the class of ligands coming from sediments. It is important to understand whether dissolved Cu emanating from sediments is likely to be associated with either weaker or stronger Cu-binding ligands, because this would impact the longevity and residence time of sediment-derived dissolved Cu and its export offshore. Weaker ligands have thus far been found in higher concentrations than stronger ligands in continental shelf regions, however even low concentrations of strong ligands will be primarily responsible for binding dissolved Cu from margin sources if they are present in excess of dissolved Cu. Determining the sources of strong ligands in near shore environments is important to constrain in future studies, to uncover the impact of ligands in stabilizing external margin sources of dissolved Cu to the ocean.

Hydrothermal vents. Hydrothermal vents play an important role for trace metal biogeochemistry as vents can provide an influx of trace metals to the overlying waters^{132,133}. Hydrothermal vents may also be a source of Cu-binding ligands^{22,134–136}. Samples originating from within the vent or its immediate surroundings have been observed to have very high concentrations of Cu-binding ligands, ranging from 29 to 4460 nmol L⁻¹, especially when compared to ligand concentrations found within a typical open-ocean profile (typically ranging from less than 1 nmol L⁻¹ to 4–6 nmol L⁻¹). Within waters impacted by hydrothermal venting, thiols and small proteins are thought to be responsible for at least some fraction of the Cu complexation²². A possible source of Cu-binding ligands within vent fluids may come from vent microbes who have been shown to produce ligands in order to buffer against high labile Cu concentrations immediately surrounding the vent field¹³⁵. Though samples within and immediately around a hydrothermal vent have been shown to have Cu-binding ligand concentrations up to 1000-fold of those found within the water column, it is not known what fraction of these ligands make it into the neutrally buoyant plume and potentially impact the global cycling of ligands and dissolved Cu¹³⁶. Current available water column data from GEOTRACES expeditions GA03¹⁶ and GP16⁷ do not show a surplus of Cu-binding ligands near hydrothermal vents in the neutrally buoyant plumes, though the sampling on these expeditions happened tens to hundreds of meters above the ridge. If hydrothermal vents are indeed a source of Cu-binding ligands to the overlying water column this would be important to constrain, as it would impact the transport of dissolved Cu in these systems.

Microbial production of Cu-binding ligands. Similar to Fe-binding ligands, a variety of marine microbes have been shown to produce Cu-binding ligands¹³⁷. Cu-binding ligand production, usually in response to Cu stress, has been observed in cultures of heterotrophic bacteria⁹⁶, cyanobacteria^{19,96,138,139}, dinoflagellates^{138,140}, coccolithophores^{141,142}, and diatoms¹³⁸. Voltammetric measurements of these ligands in culture showed a range in $\log K_{\text{CuL}_i, \text{Cu}^{2+}}$ from 11.0 to 15.6¹³⁷ (Table 1). Cu-binding ligands such as glutathione¹⁴¹, phytochelatins and metallothioneins^{55,57,143–146} are used to detoxify Cu, while compounds such as methanobactins and staphylopinins have been shown to facilitate Cu uptake^{47,101}.

Although many Cu-binding ligands produced by microbes in culture have been characterized, much less is known about the impact of microbial Cu-ligand production in seawater. The similar binding strengths of ligands observed in culture compared to those measured in seawater suggest that at least a portion of the Cu-ligand pool in seawater likely has a microbial source. Despite known ligand production from phytoplankton and bacteria, studies have only rarely identified a correlation with productivity or biomass and Cu-binding ligands⁹⁹. Thiols, for example, are produced by many phytoplankton and bacteria and yet a strong connection between productivity and thiols has not been observed in the marine environment. Unlike siderophores for Fe, thiols can be produced for a variety of reasons, including dealing with free radicals¹⁴⁷, or to detoxify other metals^{143,148–152}. Thus, the lack of a significant relationship of thiols with productivity is perhaps not surprising. It is clear however, that Cu-binding ligand production in the marine environment appears to be distinct from that of Fe ligands, likely related to differences in the role of these metals as micronutrients and toxicants. Further research is necessary to determine what might trigger Cu-binding ligand production by microbes in seawater. It is important to note however, that many of the other sources of Cu-binding ligands that have been observed (e.g., sediments, hydrothermal vents) are likely ultimately from a microbial source. Thus, understanding microbial production of ligands is critical.

A few studies have connected voltammetric measurements of the Cu-binding ligand pool with ligand characterization techniques via mass spectrometry^{23,97} or specialized voltammetric techniques aimed at characterizing specific Cu compounds such as thiols^{61,93,103,153}. Initial work that characterized Cu-binding ligands in seawater used Cu(II)-IMAC columns to capture Cu-containing organic compounds and then elute and measure them via UV-adsorption or mass spectrometry^{96,97}. This work found that the extracted Cu compounds contained nitrogen and thiol-like functional groups, and were similar to some ligands that had been found in algal cultures. Cu-binding ligands identified using Cu(II)-IMAC in the northeast Pacific along Line P also correlated with chlorophyll *a* and phaeopigments across both near shore and offshore stations and in different seasons, providing additional evidence for a biological ligand source⁹⁸. Voltammetric studies have also routinely noted high ligand concentrations and low dissolved Cu concentrations in surface waters and in the chlorophyll maximum^{8,16,103}. In the eastern South Pacific, Boiteau et al.⁶⁷ identified several distinct Cu-binding compounds via LC-ICP/ESI-MS techniques in surface waters with azole-binding groups, some of which also bound nickel (section “Components of the marine Cu-binding ligand pool”). The Cu-binding ligands identified in this study were higher near shore and decreased offshore, along with the dissolved Cu. At present, the sources of these ligands are unknown, but are presumed to be biological in origin. Further studies able to identify Cu-binding ligands in seawater will provide essential next steps in characterizing microbial sources of Cu ligands in seawater.

Microbial production of Cu ligands is likely particularly important in oxygen minimum zones (OMZs), as Cu may be a limiting factor for microbial production in these regions. Microbial production in OMZs is dominated by ammonia oxidizing archaea (AOA) and bacteria (AOB). Recent studies using genome sequencing techniques revealed that AOA and AOB produce a wide range of Cu containing enzymes^{154–156}. Moreover, experiments from Amin et al.¹⁵⁷ showed that AOA ammonia oxidation rates and cell density were strongly decreased in Cu-limiting conditions, and totally inhibited in the absence of Cu. They also observed a strong dependency of AOA-specific growth to Cu^{2+} concentrations, with limiting effects below $10^{-13} \text{ mol L}^{-1}$ and potentially toxic effects above $10^{-11} \text{ mol L}^{-1}$. These observations indicate a potential link between Cu complexation in seawater and AOA growth, since AOA appear to primarily take up free Cu. Experiments by Amin et al.¹⁵⁷ indicated that the AOA *N. maritimus* may be a source of strong Cu-binding ligands when grown under Cu-replete conditions. In these conditions, AOA may produce Cu-binding ligands to regulate Cu^{2+} concentrations at optimal levels for growth. However, Jacquot et al.⁸ argued that the low Cu^{2+} concentrations observed in the Pacific OMZ was the result of high consumption rates of Cu, leading to low Cu^{2+} , rather than high ligand production rates leading to low Cu^{2+} . They suggested that the elevated ligand concentrations found in these regions may have originated from remineralization of organic particles or advected dissolved organic material from the shelf. Direct microbial ligand production cannot be determined from these studies, because the identities of the ligands are unknown and thus any potential biosynthesis genes for ligand production cannot be explored. Whether organisms in OMZs are producing organic Cu-binding ligands to regulate growth is an important future area of dissolved Cu biogeochemistry to constrain.

Sinks of Cu-binding ligands

Microbial Cu-ligand uptake. In addition to Cu-binding ligand production, microbes may also be a sink for Cu-binding ligands. Initial work in culture studies showed that Cu uptake rates were driven primarily by Cu^{2+} concentrations and not the total dissolved Cu concentration in the culture media^{4,158,159}. These observations led to the understanding that uncomplexed Cu (Cu^{2+}) might be the only bioavailable form of Cu. However, additional studies demonstrated that eukaryotic Cu uptake rates exceeded those that would be expected from simply a diffusive supply of Cu^{2+} by 2–1000-fold, suggesting that at least some marine phytoplankton are capable of accessing organically bound Cu^{2,101,160–166}. It was unknown at the time however, whether in situ marine phytoplankton and bacteria were able to take up Cu from the natural Cu ligands present in seawater. Semeniuk et al.¹⁶⁷ provided the first such evidence that this was indeed the case, and showed that dissolved Cu in the northeast Pacific was taken up 5 times faster than would be expected based on simply Cu^{2+} . This was then substantiated with a more detailed study that showed definitively that phytoplankton and bacteria were able to take up naturally present strong and weak Cu-ligand complexes¹⁶⁸. Thus, it appears that Cu may act similarly to dissolved Fe in that Cu likely has an “envelope” of bioavailability^{169,170}, where Cu^{2+} is the most bioavailable, but some microbes can also utilize Cu from a range of ligand complexes. Bioavailability is likely related to the reducibility of Cu(II) to Cu(I) from ligand complexes, as Cu uptake transporters (known as CTR) target Cu(I) and appear to be relatively common uptake systems in some diatoms^{166,171}. Semeniuk et al.¹⁶⁸ also suggested that some phytoplankton may use weaker Cu-binding ligands as a “weak ligand shuttle” to move dissolved Cu from strong complexes to weaker, more reducible Cu complexes




Fig. 2 Copper uptake mechanisms in the marine environment. Known mechanisms of copper uptake associated with organic copper-binding ligands in the marine environment by microbes.

(Fig. 2). For example, cysteine has been shown to enhance Cu bioavailability¹⁷². Regardless, there is still much to explore with respect to how Cu speciation impacts dissolved Cu bioavailability to marine phytoplankton and bacteria, and how microbial uptake impacts Cu-binding ligand concentrations.

Photochemical degradation. Cu-binding ligands are well known to be photochemically degradable by photosynthetically active radiation and UV-light^{31,173–177}. The exact mechanism of the photodegradation is not as well known as it is for some Fe-binding ligands such as siderophores¹⁷⁸. However, both humic substances and thiols are known to be photochemically degradable^{62,177,179}. The photochemical effects on Cu-binding ligand capacity have also been observed in incubation experiments^{107,180} and in seasonal differences in Cu-binding capacities in the Gulf of Mexico¹⁸¹. Although the photochemical degradation of Cu-binding ligands is well known, the bulk of the studies focused on this topic have been in estuaries and less explored in the open ocean. It is very likely that photochemical degradation of Cu-ligand complexes impacts the bioavailability of Cu¹⁸⁰, so further work in this important area will shed important insights on ligand cycling.

Global open-ocean distributions of Cu-binding ligands. While the sources and sinks of dissolved Cu have been explored recently in basin-scale studies and biogeochemical modeling efforts^{7–9,16,182}, the relevant sources and sinks and internal cycling of Cu-binding ligands are not as well understood. Advances in metal speciation data processing have facilitated high sample throughput leading to large-scale datasets of dissolved Cu speciation. Open-ocean data from the Pacific and Atlantic oceans shows patterns in Cu-binding ligand distributions that are similar to distributions of dissolved Cu, with low nanomolar concentrations near the surface and quasi-linear increasing concentrations with depth^{5,7,8,12,16} (Fig. 3). The highest concentrations of Cu-binding ligands have been detected in the deep waters (>3000 m) from full depth ocean profiles, and near bottom sediments (concentrations up to 4–6 nmol L⁻¹). In a comparison of Cu-binding ligand distributions between the Pacific and North Atlantic basins, higher concentrations of both dissolved Cu and Cu-binding ligands were observed on average in the Pacific basin. This observation was interpreted as an accumulation of Cu-binding organic ligands in older Pacific waters (Fig. 3B⁷). The difference in ligand concentrations in deep waters of the Pacific compared to the Atlantic are not as great as the differences in dissolved Cu concentrations however, leading to




Fig. 3 Dissolved copper and ligand data for the North Atlantic and South Pacific Oceans. **A** Averaged dissolved Cu data for the Pacific and North Atlantic oceans. **B** Averaged Cu-binding ligand data for the Pacific and North Atlantic oceans collected from available datasets. **C** Averaged free Cu, $\log[\text{Cu}^{2+}]$, from available datasets for the Pacific and North Atlantic oceans^{5,7,8,12,16,247}.

some higher Cu^{2+} concentrations in the deep Pacific (Fig. 3C)⁷, though not approaching toxicity thresholds for cyanobacteria ($\sim 10^{-11} \text{ mol L}^{-1}$). Organic ligands appear to “buffer” free Cu concentrations to within a relatively narrow range in both the Atlantic and Pacific Ocean basins (Fig. 3), perhaps approaching levels that are even limiting to growth for some organisms. Recent work that suggests that a large fraction of the dissolved Cu in seawater is likely inert to exchange⁹¹, and the observation that a large portion of dissolved Cu is associated with very strong ligand complexes^{7,16} is important for considering open ocean dissolved Cu reactivity. The impact that organic ligands have on the cycling of dissolved Cu on the basin scale also has important implications for understanding whether dissolved Cu is reversibly scavenged throughout the water column, as has been proposed by a modeling study¹⁸² and some work on Cu isotopes^{183,184}.

While some strides have been made in identifying ligands in surface waters, we still do not have a thorough understanding of the identities of Cu ligands. For example, algal cultures release Cu-binding ligands such as thiols^{55,142}, and Boiteau et al.⁶⁷ identified Cu-binding ligands in the surface waters of GP16 where biological utilization of dissolved Cu is occurring, but the ultimate origin of those ligands is unknown. Though Cu-binding ligands are starting to be identified in open-ocean waters⁶⁷, these ligands have not been connected to micro-organisms or have been shown to be produced under specific circumstances. Available data from the NE Pacific suggested up to 32% of Cu-binding ligand pool was made up of humic substances and thiol type ligands⁵. While the study of humic substances and thiols in the open ocean is important for understanding the role of some key contributors to marine Cu speciation, this demonstrates that there is still a large portion of the Cu-binding ligand pool that remains unidentified, particularly in the deep ocean.

Internal cycling. Despite recent ocean-basin-scale datasets on the distributions of Cu-binding ligands, we know very little about their internal cycling. A comparison of data from the North Atlantic¹⁶ and the South Pacific⁷ suggests that both dissolved Cu and Cu-binding ligands accumulate in the oldest waters of the deep Pacific, implying that ligands are likely produced along with dissolved Cu during regeneration of sinking particles^{7,16}. Indeed, Fe and Cu-binding humic-like compounds were recently found to be directly produced during particle degradation¹⁸⁵. Deep profiles of Cu ligands from both the Atlantic and Pacific also suggest that

Cu-binding ligands are scavenged onto sinking particles. Although Cu displays a nearly linear profile with depth indicative of an influence of reversible scavenging⁹, Cu-binding ligands do not have the same profile (Fig. 3). Cu-binding ligands tend to remain in excess of dissolved Cu even near the ocean bottom, but the amount of excess ligand decreases with depth^{7,16}. It is unclear whether this is due to a lack of deep ligand sources, degradation of Cu-binding ligands with depth, or a scavenging of Cu-binding ligands onto particles. However, it is clear that the cycling of organic Cu-binding ligands is not always entirely coupled to that of dissolved Cu. Owing to our lack of knowledge on the sources, sinks and internal cycling of Cu-binding ligands we cannot yet estimate a residence time for these complexes. The only estimate of a residence time for deep ocean Fe-binding ligands was 779–1035 years¹⁸⁶, but no estimates for residence times for ligands in surface waters have been proposed. Many processes acting on Fe-binding ligands also likely overlap for Cu-binding ligands, so it is possible they have similar residence times. However, even the residence time of Cu is up for debate^{9,187} and a lot of work remains to be able to better understand the residence time of both dissolved Cu and Cu-binding ligands.

Modeling the copper-binding ligand pool

The rarity of in situ measurements and lab experiments concerning Cu-binding ligands limits our current understanding of their global cycling and their potential importance for ocean biogeochemistry. In this context, models represent a valuable tool since they enable the testing of different hypotheses regarding the influence of different processes on ligand cycling. In this section, we present some of the modeling work that focus on Cu-binding ligands. These models differ in scope, scale and complexity, but they allow for studying specific aspects of Cu-ligand cycling. We then briefly discuss how information from these different models could be integrated into global ocean models.

Geochemical models. Geochemical models can be used to constrain some external sources of ligands. For instance, Sander and Koschinsky²² and Stüeken¹⁸⁸ studied the impacts of hydrothermal sources on oceanic budgets of Cu-binding ligands. These models represent the thermodynamic mixing effects between hydrothermal fluids and the overlying water column. Both fluids have fixed Cu and ligand concentrations with fixed binding strength ($\log K_{\text{CuL}_i, \text{Cu}^{2+}}$). One fundamental hypothesis of these

models is that some ligands originate from hydrothermal vents, but that the majority are already present in the water column and that both ligand sources have similar complexing capacities. Using box models to integrate the results over the ocean, these studies revealed that 2–20% of total dissolved Cu in the modern ocean is supplied by hydrothermal vents and stabilized by organic ligands in the deep water column. Stüeken¹⁸⁸ also used this geochemical thermodynamic modeling method to represent past hydrothermal vents and showed that rivers became the dominant ocean dissolved Cu source in the Proterozoic.

Models of Cu chemical speciation. Metal ions that bind to natural organic matter such as humics vary according to metal-substrate affinity constants, reaction stoichiometries, competition between different ions, and may be impacted by water pH and salinity. Many thermodynamic chemical models exist to represent metal ion binding with humic substances and oxides in freshwater, seawater and soils^{189–194}. These models provide affinity constants for the binding of different metals (including Cu) with humics as well as descriptions of competition effects between different metals for humics. These models are frequently used in ecotoxicological assessments to identify metal contamination in freshwater, estuarine and soil systems. Furthermore, chemical speciation models are used to predict Cu toxicity to plankton and fish through the use of Biotic Ligand Models (BLM)¹⁹⁵. These models are frequently used by scientists and governments to evaluate Cu contamination and toxic impacts on soils, freshwater and estuarine systems^{196–200}.

The work of Hirose²⁰¹ specifically addressed Cu speciation and binding with organic ligands in marine systems. This author used similar principles of thermodynamic chemistry modeling to study different roles of Cu ligands, including the protecting role of weak ligands for microbes against Cu toxicity in high Cu regions²⁰¹, Cu-Fe competition for strong ligands, which may reduce Fe concentrations and lead to Fe deficiency²⁰², or the buffering effects of the excess ligands, which may shield Cu speciation from the effects of ocean acidification²⁰³.

Towards integrated modeling of Cu ligands in global biogeochemical models (GBCM). Because of the importance of Cu in regulating microbial production and due to competing hypotheses about the shape of its vertical profile, Cu has recently been included in two global models^{9,187}. However, both these models lack a representation of Cu-ligand cycling. Only in Richon and Tagliabue⁹ are Cu ligands explicitly represented with a uniform concentration and complexing capacity. Even though we are far from representing the 3-dimensional complexity of Cu-ligand cycling in global biogeochemical models (GBCMs), results from small scale modeling and increased availability of Cu-binding ligand distributions may provide some future directions for GBCM development.

The geochemical and other types of box models may provide information on the external sources of Cu ligands. Even though they do not provide constraints on the ligand fluxes from hydrothermal or riverine sources, models of thermodynamic equilibrium seem to confirm that the ligands from external sources have similar complexing capacity to those already present in the water column. These modeling efforts may indicate that there is no change in the complexing capacity of ligands during their cycling in seawater or that their ultimate sources (e.g., microbes) are the same.

Chemical speciation models provide useful information on the complexing capacities and the kinetics of Cu-ligand reactions. However, these models showed that there is a range of complexing capacities depending on the chemical nature of the

ligand. Therefore, more studies on the nature of Cu ligands are necessary in order to accurately represent Cu complexing in GBCMs. Results from BLM and other toxicity models can inform GBCMs with Cu concentration thresholds, which can be implemented to represent the toxic impacts of Cu on oceanic species. This type of information has already been used by Prosnier et al.²⁰⁴ to model Cu toxicity on freshwater *Daphnia*. Fe cycling is well represented in GBCMs and results such as Hirose (2007)²⁰² may be used to represent the competition between Cu and Fe for ligands and the potential impacts on global ocean productivity. Finally, future studies that define Cu-ligand distributions in the operationally defined classes stated in this manuscript, will facilitate cross comparisons between regions, analysts and analytical methods.

Synthesis: towards an integrated view of organic Cu-ligand cycling

Based on our review of current knowledge and understanding of Cu-ligand cycling, we summarize Cu-ligand sources, sinks and internal cycling in the global ocean and in the following section attempt to calculate some of the first global fluxes of Cu-binding ligands (Fig. 4).

Constraining the external sources of Cu-binding ligands to the oceans. Several studies indicate that hydrothermal vents may be sources of Cu ligands (section “Hydrothermal vents”). However, the nature of the ligands found near vent sites is up for debate^{22,135} and recent measurements found no surplus of ligands near vent sites^{7,8}. Therefore, based on the current understanding, it is difficult to assess the global contribution of hydrothermal vents to the global Cu-ligands budget and we have not attempted to calculate it here.

Many studies on riverine ligands highlighted a statistically significant relationship between L_1 concentrations and dissolved organic carbon (DOC)^{42,126,205,206}. This relationship is summarized in Supplementary Table 1. Based on the data we compiled, we derive a generic relationship between DOC fluxes and L_1 concentrations: $[L_1] = 0.0002[\text{DOC}] - 0.0367$ ($R^2 = 0.8129$). This equation allows the calculation of a global estimate of riverine sources of strong ligands based on global estimates of DOC riverine fluxes^{207–209}, yielding a global riverine source of ligands between 30 and 1000 $\text{Gmol L}^{-1} \text{yr}^{-1}$.

A few studies have estimated the sediment-water exchange flux of Cu ligands. Shank et al.¹²⁶ measured ligand fluxes between 850 and 870 $\text{nmol m}^{-2} \text{day}^{-1}$ (with standard deviation about ± 600). Santos-Echeandía et al.²¹⁰ calculated Cu and ligand fluxes from tidal pore waters in a salt marsh estuary in Portugal between 2 and 6 $\text{mol m}^{-2} \text{day}^{-1}$. However, ligand concentrations in estuaries are usually high, therefore the local inputs from sediments may have limited impacts on overall ligand budgets. Murray et al.²¹¹ estimated the global tidal flats area to be about $128 \times 10^9 \text{m}^2$, so using Santos-Echeandía's estimate of ligand exchange from tidal flooding gives about 0.2 Gmol L yr^{-1} of strong ligand sourced from tidal exchange. Using a global estimate of oceanic shelf areas (using the ORCA2 grid), we calculate a sediment source of Cu ligands of about 5.7 Gmol L yr^{-1} .

Several studies have measured aerosol ligand to organic carbon ratios^{212–214}. Overall, a higher ligand to carbon ratio is observed in samples taken in a forest throughfall^{212,213} than in coastal plains²¹⁴, indicating that vegetation may be a source of Cu ligands in aerosols. Based on these studies, we found an average ligand/organic carbon ratio around 450 $\text{nmol L}^{-1} \text{mg}^{-1} \text{C}$ in rainwater. Multiplying this ratio by Kanakidou et al.²¹⁵ estimate of the global wet deposition of organic carbon to the oceans




Fig. 4 Current understanding of the cycling of organic copper-binding ligands in the ocean. Schematic of copper-binding ligand cycling in the global ocean. Source fluxes that have been included in Table 4 are shown as red arrows, while sink fluxes that are shown in Table 4 are orange arrows. The flux of ligands to sediments is assumed based on assumptions of steady state. Additional ligand sources and sinks where flux calculations have not been completed are represented by black arrows. The potential for copper limitation or copper toxicity is shown to generally follow total copper concentrations, with the potential for copper toxicity greatest in coastal regions and the potential for copper limitation being greatest in open-ocean regions.

(230 Tg C yr⁻¹) gives an estimate of 0.1 Gmol L yr⁻¹ of Cu ligands from aerosol wet deposition. This estimate comes with a wide uncertainty, both because of the large differences in the ligand/organic carbon ratios in the literature, and the large uncertainties in global aerosol deposition fluxes.

To our knowledge, the only estimates of Cu-ligand concentrations in ice are from Bundy et al.²¹ who reported ligand concentrations measured in sea ice, glacier ice, and algal-influenced sea ice from Admiralty Bay (Antarctica) of 12.5, 2.7, and 26.15 nmol L⁻¹, respectively.

Abernathy et al.²¹⁶ estimated the Southern Ocean water flux from sea ice and glacier water to be, respectively, 15,750 and 1575 Gt yr⁻¹. When multiplied by Bundy et al.²¹ estimates of Cu-ligand concentrations, we obtain a potential Cu-ligand source from sea ice between 0.2 and 0.5 × 10⁻³ Gmol L⁻¹ yr⁻¹ and a potential source from glacier water of around 4 × 10⁻⁶ Gmol L⁻¹ yr⁻¹.

Several observations have shown that different phytoplankton species produce Cu ligands in conditions of Cu limitation or Cu toxicity (section “Microbial production of Cu-binding ligands”). However, quantitative estimates of Cu-ligand production rates are still missing. To our knowledge, Echeveste et al.²¹⁷ provided the only ligand production flux by the coccolithophore *E. huxleyi* (about 12.5 fM cell⁻¹ in high Cu conditions). Unfortunately, this estimate is difficult to generalize to the global ocean and more work on microbial ligand production would be necessary to quantify the magnitude of this source at the ocean scale. While it is likely that microbial ligand production is one of the dominant sources of ligands to the ocean, the current lack of rate measurements makes it impossible to estimate the value of this source given the current datasets.

Constraining the sinks of copper-binding ligands in the ocean.

Based on Semeniuk et al.¹⁶⁸ estimates, the uptake of ligand-bound Cu by phytoplankton is about 50–250 pmol Cu L⁻¹ d⁻¹, which makes 18–6750 mmol Cu m⁻³ yr⁻¹. Generalizing this consumption rate to the global surface ocean (0–100 m) gives a rough estimate of the magnitude of the Cu-ligand sink from phytoplankton uptake of 0.6–270 Gmol L yr⁻¹. This estimate has a large range and uncertainties, as regional differences in microbial abundances and community composition will have large impacts on this flux.

Using samples from Cape Fear, Shank et al.¹²⁶ estimated a degradation rate of Cu-binding ligands by sunlight of about 0.28 d⁻¹, which is much higher than the DOC photooxidation rates estimated at about 0.01 d⁻¹. This degradation rate constant can be used in the following equation to estimate ligand concentrations:

$$L_i = L_{i0}e^{-kt} \quad (1)$$

where L_{i0} is the initial ligand concentrations and at the time of exposure in days, and k is the degradation rate (estimated around 0.28 d⁻¹). Using a global ligand source of about 500 Gmol L yr⁻¹ supplied to the surface ocean, calculated from the previous section’s estimates, we obtain an estimate for global ligand photodegradation of about 123 Gmol L⁻¹ yr⁻¹.

Missing sinks and uncertainties. The fluxes discussed in the previous two sections are summarized in Table 4 and should be considered as first order estimates giving the magnitude of the sources and sinks and their large uncertainties. From this table, we can conclude that rivers are likely the major source of

Table 4 Global estimated fluxes of copper-binding ligands.

Source or sink	Global estimate (Gmol yr ⁻¹)	References
Rivers	30–1000	Muller and Batchelli (2003) ⁴² , Hoffman et al. (2007) ²⁰⁵ , Shank et al. (2004) ¹²⁶
Sediments	5.7 (continental shelf) 0.2 (tidal exchange)	Shank et al. (2004) ¹²⁶ , Santos-Echeandia et al. (2013) ²¹⁰
Aerosols (wet deposition)	0.1	Kanakidou et al. (2012) ²¹⁵ , Hou et al. (2005) ²¹³ , Karavoltos et al. (2017) ²⁴⁸ , Witt et al. (2007) ²¹⁴
Sea ice	0.2–0.5 × 10 ⁻³	Abernathy et al. (2016) ²¹⁶ , Bundy et al. (2013) ²¹
Glacier (estimates for the Southern Ocean)	4 × 10 ⁻⁶	
Phytoplankton uptake	0.6–270	Semeniuk et al. (2015) ¹⁶⁸
Photodegradation	123	Shank et al. (2006) ¹⁷³

Estimates of global oceanic fluxes of organic Cu-binding ligand sources and sinks. We have not included hydrothermal or microbial sources despite both likely being an important source, as the appropriate data was not readily available.

Cu-binding ligands to ocean waters, with sediments and aerosols being second-order sources. Sea ice and glacier melt probably have a minor impact on Cu-ligand budgets, however, sea-ice algae may be a strong source of Cu ligands and subsequent release after sea-ice melt may locally influence Cu cycling. The estimates for Cu-ligand sinks show that both phytoplankton uptake and photodegradation may have comparable magnitude, however, these estimates have large uncertainties that may span several orders of magnitude. If we assume steady state for the global Cu-ligand budget, the global ligand sink to sediments can be estimated on the order of hundreds of Gmol L yr⁻¹, making sedimentation likely the greatest sink for Cu ligands. However, we found no in situ estimate for Cu-ligand loss to sediments. Future research efforts should focus on constraining this sink by measuring ligands associated with sinking organic material.

Conclusions

Organic Cu-binding ligands bind the majority of the dissolved Cu in the ocean and are recognized as being important for dissolved Cu cycling, and yet much remains to be discovered about their sources, sinks and internal cycling, and none of the existing Cu biogeochemical models include a dynamic cycle of Cu-binding ligands. Here, we summarized the current knowledge of Cu-ligand cycling in the global ocean with a focus on large-scale open-ocean processes based largely on basin-scale GEOTRACES datasets and found that margin sediments and rivers are the major sources of Cu-binding ligands in seawater, and sedimentation, microbial uptake and photochemical degradation are the major sinks. Future studies that focus on understanding Cu-binding ligand fluxes and identifying Cu-binding ligands in seawater will be particularly insightful for future modeling efforts, and for understanding the impact of Cu-binding ligands on Cu bioavailability to marine organisms.

Data availability

The data for Fig. 1 is available via figshare (<https://doi.org/10.6084/m9.figshare.21183709>) and data associated with Fig. 3 is available from BCO-DMO (<https://www.bco-dmo.org/dataset/740051>).

Received: 3 May 2022; Accepted: 19 October 2022;

Published online: 02 November 2022

References

- Twining, B. S. & Baines, S. B. The trace metal composition of marine phytoplankton. *Ann. Rev. Mar. Sci.* **5**, 191–215 (2013).

- Annett, A. L., Lapi, S., Ruth, T. J. & Maldonado, M. T. The effects of Cu and Fe availability on the growth and Cu:C ratios of marine diatoms. *Limnol. Oceanogr.* **53**, 2451–2461 (2008).
- Maldonado, M. T. et al. Copper-dependent iron transport in coastal and oceanic diatoms. *Limnol. Oceanogr.* **51**, 1729–1743 (2006).
- Brand, L. E., Sunda, W. G. & Guillard, R. R. L. Reduction of marine phytoplankton reproduction rates by copper and cadmium. *J. Exp. Mar. Bio. Ecol.* **96**, 225–250 (1986).
- Whitby, H., Posacka, A. M., Maldonado, M. T. & van den Berg, C. M. G. Copper-binding ligands in the NE Pacific. *Mar. Chem.* **204**, 36–48 (2018).
- Wong, K. H., Obata, H., Kim, T., Wakuta, Y. & Takeda, S. Distribution and speciation of copper and its relationship with FDOM in the East China Sea. *Mar. Chem.* **212**, 96–107 (2019).
- Ruacho, A. et al. Organic dissolved copper speciation across the U.S. GEOTRACES equatorial Pacific zonal transect GP16. *Mar. Chem.* **225**, 103841 (2020).
- Jacquot, J. E., Kondo, Y., Knapp, A. N. & Moffett, J. W. The speciation of copper across active gradients in nitrogen-cycle processes in the eastern tropical south Pacific. *Limnol. Oceanogr.* **58**, 1387–1394 (2013).
- Richon, C. & Tagliabue, A. Insights Into the major processes driving the global distribution of copper in the ocean from a global model. *Global Biogeochem. Cycles* **33**, 1594–1610 (2019).
- Hollister, A. P. et al. Dissolved concentrations and organic speciation of copper in the Amazon River estuary and mixing plume. *Mar. Chem.* **234**, 104005 (2021).
- Whitby, H., Hollibaugh, J. T. & van den Berg, C. M. G. Chemical speciation of copper in a salt marsh estuary and bioavailability to Thaumarchaeota. *Front. Mar. Sci.* **4**, 178 (2017).
- Thompson, C. M., Ellwood, M. J. & Sander, S. G. Dissolved copper speciation in the Tasman Sea, SW Pacific Ocean. *Mar. Chem.* **164**, 84–94 (2014).
- Posacka, A. M. et al. Dissolved copper (dCu) biogeochemical cycling in the subarctic Northeast Pacific and a call for improving methodologies. *Mar. Chem.* **196**, 47–61 (2017).
- Little, S. H., Vance, D., Walker-Brown, C. & Landing, W. M. The oceanic mass balance of copper and zinc isotopes, investigated by analysis of their inputs, and outputs to ferromanganese oxide sediments. *Geochim. Cosmochim. Acta* **125**, 673–693 (2014).
- Paytan, A. et al. Toxicity of atmospheric aerosols on marine phytoplankton. *Proc. Natl. Acad. Sci. USA* **106**, 4601–4605 (2009).
- Jacquot, J. E. & Moffett, J. W. Copper distribution and speciation across the International GEOTRACES Section GA03. *Deep. Res. Part II Top. Stud. Oceanogr.* **116**, 187–207 (2015).
- Roshan, S. & Wu, J. The distribution of dissolved copper in the tropical-subtropical north Atlantic across the GEOTRACES GA03 transect. *Mar. Chem.* **176**, 189–198 (2015).
- Campos, M. L. A. M. & Van Den Berg, C. M. G. Determination of copper complexation in sea water by cathodic stripping voltammetry and ligand competition with salicylaldehyde. *Analyt. Chim. Acta* **284**, 481–496 (1994).
- Moffett, J. W. & Brand, L. E. Production of strong, extracellular Cu chelators by marine cyanobacteria in response to Cu stress. *Limnol. Oceanogr.* **41**, 388–395 (1996).
- Buck, K. N. & Bruland, K. W. Copper speciation in San Francisco Bay: A novel approach using multiple analytical windows. *Mar. Chem.* **96**, 185–198 (2005).
- Bundy, R. M., Barbeau, K. A. & Buck, K. N. Sources of strong copper-binding ligands in Antarctic Peninsula surface waters. *Deep. Res.* **90**, 134 (2013).
- Sander, S. G. & Koschinsky, A. Metal flux from hydrothermal vents increased by organic complexation. *Nat. Geosci.* **4**, 145–150 (2011).

23. Boiteau, R. M. et al. Siderophore-based microbial adaptations to iron scarcity across the eastern Pacific Ocean. *Proc. Natl. Acad. Sci. USA* **113**, 14237–14242 (2016).
24. Tagliabue, A. et al. The integral role of iron in ocean biogeochemistry. *Nat. Publ. Gr.* **543**, 51–59 (2017).
25. Völker, C. & Tagliabue, A. Modeling organic iron-binding ligands in a three-dimensional biogeochemical ocean model. *Mar. Chem.* **173**, 67–77 (2015).
26. Kogut, M. B. & Voelker, B. M. Strong copper-binding behavior of terrestrial humic substances in seawater. *Environ. Sci. Technol.* **35**, 1149–1156 (2001).
27. Town, R. M. & Filella, M. Dispelling the myths: is the existence of L1 and L2 ligands necessary to explain metal ion speciation in natural waters? *Limnol. Oceanogr.* **45**, 1341–1357 (2000).
28. Turner, D. R., Whitfield, M. & Dickson, A. G. The equilibrium speciation of dissolved components in freshwater and sea water at 25 °C and 1 atm pressure. *Geochim. Cosmochim. Acta* **45**, 855–881 (1981).
29. Moffett, J. W. & Zika, R. G. Oxidation kinetics of Cu(I) in seawater: implications for its existence in the marine environment. *Mar. Chem.* **13**, 239–251 (1983).
30. Wuttig, K., Heller, M. I., & Croot, P. L. Pathways of superoxide (O₂(–)) decay in the eastern tropical North Atlantic. *Environ. Sci. Technol.* <https://doi.org/10.1021/es401658t> (2013).
31. Moffett, J. W. & Zika, R. G. Measurement of copper(I) in surface waters of the subtropical Atlantic and Gulf of Mexico. *Geochim. Cosmochim. Acta* **52**, 1849–1857 (1988).
32. Buerge-Weirich, D. & Sulzberger, B. Formation of Cu(I) in estuarine and marine waters: application of a new solid-phase extraction method to measure Cu(I). *Environ. Sci. Technol.* **38**, 1843–1848 (2004).
33. Michael, J. P. & Pattenden, G. Marine metabolites and metal ion chelation: the facts and the fantasies. *Angew. Chem.* **32**, <https://doi.org/10.1002/anie.199300013> (1993).
34. Walsh, M. J. & Ahner, B. A. Determination of stability constants of Cu(I), Cd(II) & Zn(II) complexes with thiols using fluorescent probes. *J. Inorg. Biochem.* **128**, 112–123 (2013).
35. Abualhajja, M. M., Whitby, H. & van den Berg, C. M. G. Competition between copper and iron for humic ligands in estuarine waters. *Mar. Chem.* **172**, 46–56 (2015).
36. Martin-Pastor, M. et al. Structure, rheology, and copper-complexation of a hyaluronan-like exopolysaccharide from *Vibrio*. *Carbohydr. Polym.* **222**, 114999 (2019).
37. Hassler, C. S., van den Berg, C. M. G. & Boyd, P. W. Toward a regional classification to provide a more inclusive examination of the ocean biogeochemistry of iron-binding ligands. *Front. Mar. Sci.* **4**, 19 (2017).
38. Croue, J. P., Benedetti, M. F., Violleau, D. & Leenheer, J. A. Characterization and copper binding of humic and nonhumic organic matter isolated from the South Platte river: evidence for the presence of nitrogenous binding site. *Environ. Sci. Technol.* **37**, 328–336 (2003).
39. Harvey, G. R., Boran, D. A., Chesal, L. A. & Tokar, J. M. The structure of marine fulvic and humic acids. *Mar. Chem.* **12**, 119–132 (1983).
40. Myklestad, S. M. Release of extracellular products by phytoplankton with special emphasis on polysaccharides. *Sci. Total Environ.* **165**, 155–164 (1995).
41. Dulaquais, G., Waeles, M., Breitenstein, J., Knoery, J. & Riso, R. Links between size fractionation, chemical speciation of dissolved copper and chemical speciation of dissolved organic matter in the Loire estuary. *Environ. Chem.* **17**, 385–399 (2020).
42. Muller, F. L. L. & Batchelli, S. Copper binding by terrestrial versus marine organic ligands in the coastal plume of River Thurso, North Scotland. *Estuar. Coast. Shelf Sci.* **133**, 137–146 (2013).
43. Whitby, H. & van den Berg, C. M. G. Evidence for copper-binding humic substances in seawater. *Mar. Chem.* **173**, 282–290 (2015).
44. Lombardi, A. T., Hidalgo, T. M. R. & Vieira, A. A. H. Copper complexing properties of dissolved organic materials exuded by the freshwater microalgae *Scenedesmus acuminatus* (Chlorophyceae). *Chemosphere* **60**, 453–459 (2005).
45. Kenney, G. E. & Rosenzweig, A. C. Chalkophores. *Annu. Rev. Biochem.* **87**, 645–676 (2018).
46. Choi, D. W. et al. Spectral, kinetic, and thermodynamic properties of Cu(I) and Cu(II) binding by methanobactin from *Methylosinus trichosporium* OB3b †. *Biochemistry* **45**, 1442–1453 (2006).
47. El Ghazouani, A. et al. Variations in methanobactin structure influences copper utilization by methane-oxidizing bacteria. *Proc. Natl. Acad. Sci. USA* **109**, 8400–8404 (2012).
48. Semrau, J. D., Dispirito, A. A. & Yoon, S. Methanotrophs and copper. *FEMS Microbiol. Rev.* **34**, 496–531 (2010).
49. Chen, S. et al. Population dynamics of methanogens and methanotrophs along the salinity gradient in Pearl River Estuary: implications for methane metabolism. *Appl. Microbiol. Biotechnol.* <https://doi.org/10.1007/s00253-019-10221-6> (2020).
50. Elsaied, H. E., Hayashi, T. & Naganuma, T. Molecular analysis of deep-sea hydrothermal vent aerobic methanotrophs by targeting genes of 16S rRNA and particulate methane monooxygenase. *Mar. Biotechnol. (NY)* <https://doi.org/10.1007/s10126-004-3042-0> (2004).
51. Lesniewski, R. A., Jain, S., Anantharaman, K., Schloss, P. D. & Dick, G. J. The metatranscriptome of a deep-sea hydrothermal plume is dominated by water column methanotrophs and lithotrophs. *ISME J.* **6**, 2257–2268 (2012).
52. Tavormina, P. L., Ussler, W., Joye, S. B., Harrison, B. K. & Orphan, V. J. Distributions of putative aerobic methanotrophs in diverse pelagic marine environments. *ISME J.* **4**, 700–710 (2010).
53. Dassama, L. M. K., Kenney, G. E. & Rosenzweig, A. C. Methanobactins: from genome to function. *Metallomics* **9**, 7–20 (2017).
54. Swarr, G. J., Kading, T., Lamborg, C. H., Hammerschmidt, C. R. & Bowman, K. L. Dissolved low-molecular weight thiol concentrations from the U.S. GEOTRACES North Atlantic Ocean zonal transect. *Deep. Res. Part I Oceanogr. Res. Pap.* **116**, 77–87 (2016).
55. Ahner, B. A., Kong, S. & Morel, F. M. M. Phytochelatin production in marine algae. 1. An interspecies comparison. *Limnol. Oceanogr.* **40**, 649–657 (1995).
56. Ahner, B. A., Lee, J. G., Price, N. M. & Morel, F. M. M. Phytochelatin concentrations in the equatorial Pacific. *Deep Sea Res. Part I Oceanogr. Res. Pap.* **45**, 1779–1796 (1998).
57. Ahner, B. A., Morel, F. M. M. & Moffett, J. W. Trace metal control of phytochelatin production in coastal waters. *Limnol. Oceanogr.* **42**, 601–608 (1997).
58. Tripathi, S. & Poluri, K. M. Metallothionein-and phytochelatin-assisted mechanism of heavy metal detoxification in microalgae. approaches to remediate. *Inorg. Pollut.* 323–344 https://doi.org/10.1007/978-981-15-6221-1_16 (2021).
59. Chapman, C. S., Capodaglio, G., Turetta, C. & Van den Berg, C. M. G. Benthic fluxes of copper, complexing ligands and thiol compounds in shallow lagoon waters. *Mar. Environ. Res.* **67**, 17–24 (2009).
60. Dryden, C. L., Gordon, A. S. & Donat, J. R. Seasonal survey of copper-complexing ligands and thiol compounds in a heavily utilized, urban estuary: Elizabeth River, Virginia. *Mar. Chem.* **103**, 276–288 (2007).
61. Dupont, C. L., Moffett, J. W., Bidigare, R. R. & Ahner, B. A. Distributions of dissolved and particulate biogenic thiols in the subtropical Pacific Ocean. *Deep. Res. Part I Oceanogr. Res. Pap.* **53**, 1961–1974 (2006).
62. Gao, Z. & Guéguen, C. Distribution of thiol, humic substances and colored dissolved organic matter during the 2015 Canadian Arctic GEOTRACES cruises. *Mar. Chem.* **203**, 1–9 (2018).
63. Le Gall, A. C. & Van Den Berg, C. M. G. Folic acid and glutathione in the water column of the North East Atlantic. *Deep. Res. Part I Oceanogr. Res. Pap.* **45**, 1903–1918 (1998).
64. Tang, D., Hung, C.-C., Warnken, K. W. & Santschi, P. H. The distribution of biogenic thiols in surface waters of Galveston Bay. *Limnol. Oceanogr.* **45**, <https://doi.org/10.4319/lo.2000.45.6.1289> (2000).
65. Zhang, J., Wang, F., House, J. D. & Page, B. Thiols in wetland interstitial waters and their role in mercury and methylmercury speciation. *Limnol. Oceanogr.* **49**, <https://doi.org/10.4319/lo.2004.49.6.2276> (2004).
66. Leal, M. F. C. & Van den Berg, C. M. G. Evidence for strong copper(I) complexation by organic ligands in seawater. *Aquat. Geochem.* **4**, 49–75 (1998).
67. Boiteau, R. M. et al. Structural characterization of natural nickel and copper binding ligands along the US GEOTRACES eastern Pacific zonal transect. *Front. Mar. Sci.* **3**, <https://doi.org/10.3389/fmars.2016.00243> (2016).
68. Louropoulou, E. et al. Heme b distributions through the Atlantic Ocean: evidence for “anemic” phytoplankton populations. *Sci. Rep.* **10**, 1–14 (2020).
69. Hogle, S. L., Barbeau, K. A. & Gledhill, M. Heme in the marine environment: from cells to the iron cycle. *Metallomics* **6**, 1107 (2014).
70. Rue, E. & Bruland, K. Domoic acid binds iron and copper: a possible role for the toxin produced by the marine diatom *Pseudo-nitzschia*. *Mar. Chem.* **76**, 127–134 (2001).
71. Coale, K. H. & Bruland, K. W. Copper Complexation in the Northeast Pacific. *Limnol. Oceanogr.* **33**, 1084–1101 (1988).
72. Coale, K. H. & Bruland, K. W. Spatial and temporal variability in copper complexation in the North Pacific. *Deep. Res. Part A-Oceanographic Res. Pap.* **37**, 317–336 (1990).
73. Mantoura, R. F. C. & Riley, J. P. Analytical concentration of humic substances from natural-waters. *Anal. Chim. Acta* **76**, 97–106 (1975).
74. Ruzic, I. Theoretical aspects of the direct titration of natural-waters and its information yield for trace-metal speciation. *Anal. Chim. Acta* **140**, 99–113 (1982).
75. Omanović, D. et al. Significance of data treatment and experimental setup on the determination of copper complexing parameters by anodic stripping voltammetry. *Anal. Chim. Acta* **664**, 136–143 (2010).
76. Turner, D. R., Whitfield, M. & Dickson, A. G. The equilibrium speciation of dissolved components in fresh-water and seawater At 25-degrees-C and 1 atm pressure. *Geochim. Cosmochim. Acta* **45**, 855–881 (1981).
77. Bruland, K. W., Rue, E. L., Donat, J. R., Skrabal, S. A. & Moffett, J. W. Intercoupling of voltammetric techniques to determine the chemical

- speciation of dissolved copper in a coastal seawater sample. *Anal. Chim. Acta* **405**, 99–113 (2000).
78. van den Berg, C. M. G. & Donat, J. R. Determination and data evaluation of copper complexation by organic ligands in sea water using cathodic stripping voltammetry at varying detection windows. *Anal. Chim. Acta* **257**, 281–291 (1992).
 79. van den Berg, M. G. C. Determination of the complexing capacity and conditional stability constants of complexes of copper(II) with natural organic ligands in seawater by cathodic stripping voltammetry of copper-catechol complex ions. *Mar. Chem.* **15**, 1–18 (1984).
 80. van den Berg, C. M. G., Merks, A. G. A. & Duursma, E. K. Organic complexation and its control of the dissolved concentrations of copper and zinc in the Scheldt estuary. *Estuar. Coast. Shelf Sci.* **24**, 785–797 (1987).
 81. Ružič, I. Theoretical aspects of the direct titration of natural waters and its information yield for trace metal speciation. *Anal. Chim. Acta* **140**, 99–113 (1982).
 82. Pižeta, I. et al. Interpretation of complexometric titration data: An intercomparison of methods for estimating models of trace metal complexation by natural organic ligands. *Mar. Chem.* **173**, 3–24 (2015).
 83. Omanovic, D. et al. ProMCC: an all-in-one tool for trace metal complexation studies. *Mar. Chem.* **173**, 25–39 (2015).
 84. Heller, M. I. & Croot, P. L. Copper speciation and distribution in the Atlantic sector of the Southern Ocean. *Mar. Chem.* **173**, 253–268 (2015).
 85. Scatchard, G. The attractions of proteins for small molecules and ions. *Ann. N. Y. Acad. Sci.* **51**, 660–672 (1949).
 86. Van Den Berg, C. M. G. Determination of copper complexation with natural organic ligands in seawater by equilibration with MnO₂ II. Experimental procedures and application to surface seawater. *Mar. Chem.* **11**, 323–342 (1982).
 87. Gerringa, L. J. A., Rijstenbil, J. W., Poortvliet, T. C. W., van Drie, J. & Schot, M. C. Speciation of copper and responses of the marine diatom *Ditylum brightwellii* upon increasing copper concentrations. *Aquat. Toxicol.* **31**, 77–90 (1995).
 88. Nuester, J. & Van Den Berg, C. M. G. Determination of metal speciation by reverse titrations. *Anal. Chem.* **77**, 11–19 (2005).
 89. Wong, K. H. et al. Wide-range detection of Cu-binding organic ligands in seawater using reverse titration. *Mar. Chem.* **230**, 103927 (2021).
 90. Van Den Berg, C. M. G., Nimmo, M., Daly, P. & Turner, D. R. Effects of the detection window on the determination of organic copper speciation in estuarine waters. *Anal. Chim. Acta* **232**, 149–159 (1990).
 91. Moriyasu, R. & Moffett, J. W. Determination of inert and labile copper on GEOTRACES samples using a novel solvent extraction method. *Mar. Chem.* **239**, 104073 (2022).
 92. Pernet-Coudrier, B., Waeles, M., Filella, M., Quentel, F. & Riso, R. D. Simple and simultaneous determination of glutathione, thioacetamide and refractory organic matter in natural waters by DP-CSV. *Sci. Total Environ.* **463–464**, 997–1005 (2013).
 93. Laglera, L. M., Downes, J., Tovar-Sánchez, A. & Monticelli, D. Cathodic pseudopolarography: a new tool for the identification and quantification of cysteine, cystine and other low molecular weight thiols in seawater. *Anal. Chim. Acta* **836**, 24–33 (2014).
 94. Waska, H., Koschinsky, A., Ruiz Chanco, M. J. & Dittmar, T. Investigating the potential of solid-phase extraction and Fourier-transform ion cyclotron resonance mass spectrometry (FT-ICR-MS) for the isolation and identification of dissolved metal-organic complexes from natural waters. *Mar. Chem.* **173**, 78–92 (2015).
 95. Gordon, A. S., Dyer, B. J., Kango, R. A. & Donat, J. R. Copper ligands isolated from estuarine water by immobilized metal affinity chromatography: temporal variability and partial characterization. *Mar. Chem.* **53**, 163–172 (1996).
 96. Gordon, A. S., Donat, J. R., Kango, R. A., Dyer, B. J. & Stuart, L. M. Dissolved copper-complexing ligands in cultures of marine bacteria and estuarine water. *Mar. Chem.* **70**, 149–160 (2000).
 97. Ross, A. R. S., Ikonou, M. G. & Orians, K. J. Characterization of copper-complexing ligands in seawater using immobilized copper(II)-ion affinity chromatography and electrospray ionization mass spectrometry. *Mar. Chem.* **83**, 47–58 (2003).
 98. Nixon, R. L. et al. Evidence for the production of copper-complexing ligands by marine phytoplankton in the subarctic northeast Pacific. *Mar. Chem.* **237**, 104034 (2021).
 99. Nixon, R. L., Jackson, S. L., Cullen, J. T. & Ross, A. R. S. Distribution of copper-complexing ligands in Canadian Arctic waters as determined by immobilized copper(II)-ion affinity chromatography. *Mar. Chem.* **215**, 103673 (2019).
 100. Gledhill, M. & Buck, K. N. The organic complexation of iron in the marine environment: a review. *Front. Microbiol.* **3**, <https://doi.org/10.3389/fmicb.2012.00069> (2012).
 101. Kim, H. J. et al. Methanobactin, a copper-acquisition compound from methane-oxidizing bacteria. *Science* (80-). **305**, 1612–1615 (2004).
 102. Kleint, C. et al. Organic Cu-complexation at the shallow marine hydrothermal vent fields off the coast of Milos (Greece), Dominica (Lesser Antilles) and the Bay of Plenty (New Zealand). *Mar. Chem.* **173**, 244–252 (2015).
 103. Moffett, J. W. & Dupont, C. Cu complexation by organic ligands in the subarctic NW Pacific and Bering Sea. *Deep. Res. Part I Oceanogr. Res. Pap.* **54**, 586–595 (2007).
 104. Buck, K. N. et al. The organic complexation of iron and copper: an intercomparison of competitive ligand exchange-adsorptive cathodic stripping voltammetry (CLE-ACSV) techniques. *Limnol. Oceanogr. Methods* **10**, 496–515 (2012).
 105. Oldham, V. E., Swensen, M. & Buck, K. N. Spatial variability of total dissolved copper and copper speciation in the inshore waters of Bermuda. *Mar. Pollut. Bull.* **79.1**, 314–320 (2014).
 106. Arnone, V., González-Santana, D., González-Dávila, M., González, A. G. & Santana-Casiano, J. M. Iron and copper complexation in Macaronesian coastal waters. *Mar. Chem.* **240**, 104087 (2022).
 107. Buck, K. N., Selph, K. E. & Barbeau, K. A. Iron-binding ligand production and copper speciation in an incubation experiment of Antarctic Peninsula shelf waters from the Bransfield Strait, Southern Ocean. *Mar. Chem.* **122**, 148–159 (2010).
 108. Boyle, E. A., Husted, S. S. & Grant, B. The chemical mass balance of the amazon plume-II. Copper, nickel, and cadmium. *Deep Sea Res. Part A, Oceanogr. Res. Pap.* **29**, 1355–1364 (1982).
 109. Windom, H. et al. Trace metal transport in a tropical estuary. *Mar. Chem.* **24**, 293–305 (1988).
 110. Abdel-Moati, A. R. Behaviour and fluxes of copper and lead in the Nile River estuary. *Estuar. Coast. Shelf Sci.* **30**, 153–165 (1990).
 111. Cutter, G. Trace elements in estuarine and coastal waters—U.S. studies from 1986–1990 article. *Rev. Geophys.* <https://doi.org/10.1002/rog.1991.29.s2.639> (1991).
 112. Zhang, J. Geochemistry of trace metals from Chinese river/estuary systems: an overview. *Estuar. Coast. Shelf Sci.* **41**, 631–658 (1995).
 113. Illuminati, S. et al. In-situ trace metal (Cd, Pb, Cu) speciation along the Po River plume (Northern Adriatic Sea) using submersible systems. *Mar. Chem.* **212**, 47–63 (2019).
 114. Plavšić, M. et al. Determination of the copper complexing ligands in the Krka river estuary. *Fresenius Environ. Bull.* **18**, 327–334 (2009).
 115. Padan, J. et al. Organic copper speciation by anodic stripping voltammetry in estuarine waters with high dissolved organic matter. *Front. Chem.* **8**, 628749 (2021).
 116. Pearson, H. B. C., Comber, S. D. W., Braungardt, C. & Worsfold, P. J. Predicting copper speciation in estuarine waters—is dissolved organic carbon a good proxy for the presence of organic ligands? *Environ. Sci. Technol.* **51**, 2206–2216 (2017).
 117. Pavoni, E. et al. Partitioning and mixing behaviour of trace elements at the Isonzo/Soča River mouth (Gulf of Trieste, northern Adriatic Sea). *Mar. Chem.* **223**, 103800 (2020).
 118. Jordi, A., Basterretxea, G., Tovar-Sánchez, A., Alastuey, A. & Querol, X. Copper aerosols inhibit phytoplankton growth in the Mediterranean Sea. *Proc. Natl. Acad. Sci. USA* **109**, 21246–21249 (2012).
 119. Moffett, J. W., Brand, L. E., Croot, P. L. & Barbeau, K. A. Cu speciation and cyanobacterial distribution in harbors subject to anthropogenic Cu inputs. *Limnol. Oceanogr.* **42**, 789–799 (1997).
 120. Hong, S., Candelone, J.-P., Turetta, C. & Boutroun, C. F. Changes in natural lead, copper, zinc and cadmium concentrations in central Greenland ice from 8250 to 149,100 years ago: their association with climatic changes and resultant variations of dominant source contributions. *Earth Planet. Sci. Lett.* **143**, 233–244 (1996).
 121. Sholkovitz, E. R. Flocculation of dissolved organic and inorganic matter during the mixing of river water and seawater. *Geochim. Cosmochim. Acta* **40**, 831–845 (1976).
 122. Jokinen, S. A., Jilbert, T., Tiihonen-Filppula, R. & Koho, K. Terrestrial organic matter input drives sedimentary trace metal sequestration in a human-impacted boreal estuary. *Sci. Total Environ.* **717**, 137047 (2020).
 123. Wilkinson, K. J., Joz-Roland, A. & Buffle, J. Different roles of pedogenic fulvic acids and aquagenic biopolymers on colloid aggregation and stability in freshwaters. *Limnol. Oceanogr.* **42**, 1714–1724 (1997).
 124. Karbassi, A. R., Bassam, S. S. & Ardestani, M. Flocculation of Cu, Mn, Ni, Pb, and Zn during estuarine mixing (Caspian Sea). *Int. J. Environ. Res.* **7**, 917–924 (2013).
 125. Buck, K. N., Ross, J. R. M., Russell Flegal, A. & Bruland, K. W. A review of total dissolved copper and its chemical speciation in San Francisco Bay, California. *Environ. Res.* **105**, 5–19 (2007).
 126. Shank, G. C., Skrabal, S. A., Whitehead, R. F. & Kieber, R. J. Fluxes of strong Cu-complexing ligands from sediments of an organic-rich estuary. *Estuar. Coast. Shelf Sci.* **60**, 349–358 (2004).
 127. Sander, S. G., Buck, K. N. & Wells, M. The effect of natural organic ligands on trace metal speciation in San Francisco Bay: Implications for water quality criteria. *Mar. Chem.* **173**, 269–281 (2015).

128. Wells, M., Buck, K. N. & Sander, S. G. New approach to analysis of voltammetric ligand titration data improves understanding of metal speciation in natural waters. *Limnol. Oceanogr. Methods* **11**, 450–465 (2013).
129. Earley, P. J. et al. Life cycle contributions of copper from vessel painting and maintenance activities. *Biofouling* **30**, 51–68 (2014).
130. Sanial, V. et al. Radium-228 as a tracer of dissolved trace element inputs from the Peruvian continental margin. *Mar. Chem.* **201**, 20–34 (2018).
131. Skrabal, S. A., Donat, J. R. & Burdige, D. J. Pore water distributions of dissolved copper and copper-complexing ligands in estuarine and coastal marine sediments. *Geochim. Cosmochim. Acta* **64**, 1843–1857 (2000).
132. Resing, J. A. et al. Basin-scale transport of hydrothermal dissolved metals across the South Pacific Ocean. *Nature* **523**, 200–U140 (2015).
133. Buck, K. N., Sohst, B. & Sedwick, P. N. The organic complexation of dissolved iron along the U.S. GEOTRACES (GA03) North Atlantic Section. *Deep. Res. Part II Top. Stud. Oceanogr.* **116**, 152–165 (2015).
134. Cotte, L. et al. On the nature of dissolved copper ligands in the early buoyant plume of hydrothermal vents. *Environ. Chem.* **15**, 58 (2018).
135. Klevenz, V., Sander, S. G., Perner, M. & Koschinsky, A. Amelioration of free copper by hydrothermal vent microbes as a response to high copper concentrations. *Chem. Ecol.* **28**, 405–420 (2012).
136. Sander, S. G., Koschinsky, A., Massoth, G. J., Stott, M. & Hunter, K. A. Organic complexation of copper in deep-sea hydrothermal vent systems. *Environ. Chem.* **4**, 81 (2007).
137. Wiramanaden, C. I. E. E., Cullen, J. T., Ross, A. R. S. S. & Orians, K. J. Cyanobacterial copper-binding ligands isolated from artificial seawater cultures. *Mar. Chem.* **110**, 28–41 (2008).
138. Croot, P. L., Moffett, J. W. & Luther, G. W. Polarographic determination of half-wave potentials for copper-organic complexes in seawater. *Mar. Chem.* **67**, 219–232 (1999).
139. Lawrence, M. G. *Voltammetric Characterisation of A Strong Extracellular Copper Binding Ligand from Synechococcus PCC7002*. (University of British Columbia, 1998).
140. Croot, P. L., Moffett, J. W. & Brand, L. E. Production of extracellular Cu complexing ligands by eucaryotic phytoplankton in response to Cu stress. *Limnol. Oceanogr.* **45**, 619–627 (2000).
141. Dupont, C. L., Nelson, R. K., Bashir, S., Moffett, J. W. & Ahner, B. A. Novel copper-binding and nitrogen-rich thiols produced and exuded by *Emiliania huxleyi*. *Limnol. Oceanogr.* **49**, 1754–1762 (2004).
142. Leal, M. F. C., Vasconcelos, M. T. S. D. & Van Den Berg, C. M. G. Copper-induced release of complexing ligands similar to thiols by *Emiliania huxleyi* in seawater cultures. *Limnol. Oceanogr.* **44**, 1750–1762 (1999).
143. Ahner, B. A., Wei, L., Oleson, J. R. & Ogura, N. Glutathione and other low molecular weight thiols in marine phytoplankton under metal stress. *Mar. Ecol. Prog. Ser.* **232**, 93–103 (2002).
144. Wei, L. & Ahner, B. A. Sources and sinks of dissolved phytochelatin in natural seawater. *Limnol. Oceanogr.* **50**, 13–22 (2005).
145. Ahner, B. A. & Morel, F. M. M. Phytochelatin production in marine algae. 2. Induction by various metals. *Limnol. Oceanogr.* **40**, 658–665 (1995).
146. Deveux, A., Achterberg, E. & Gledhill, M. Metal ion-binding properties of phytochelatin and related ligands. *Met. Ions Life Sci.* **5**, 441–481 (2009).
147. Morelli, E. & Scarano, G. Copper-induced changes of non-protein thiols and antioxidant enzymes in the marine microalga *Phaeodactylum tricornutum*. *Plant Sci.* **167**, 289–296 (2004).
148. Miao, A.-J. & Wang, W.-X. Predicting copper toxicity with its intracellular or subcellular concentration and the thiol synthesis in a marine diatom. *Environ. Sci. Technol.* **41**, 1777–1782 (2007).
149. Wang, M.-J. & Wang, W.-X. Cadmium in three marine phytoplankton: accumulation, subcellular fate and thiol induction. *Aquat. Toxicol.* **95**, 99–107 (2009).
150. Wu, Y. & Wang, W.-X. Thiol compounds induction kinetics in marine phytoplankton during and after mercury exposure. *J. Hazard. Mater.* **217**, 271–278 (2012).
151. Wang, M.-J. & Wang, W.-X. Temperature-dependent sensitivity of a marine diatom to cadmium stress explained by subcellular distribution and thiol synthesis. *Environ. Sci. Technol.* **42**, 8603–8608 (2008).
152. Stauber, J. L. & Florence, T. M. Mechanism of toxicity of ionic copper and copper complexes to algae. *Mar. Biol.* **94**, 511–519 (1987).
153. Laglera, L. M. & van den Berg, C. M. G. Copper complexation by thiol compounds in estuarine waters. *Mar. Chem.* **82**, 71–89 (2003).
154. Blainey, P. C., Mosier, A. C., Potanina, A., Francis, C. A. & Quake, S. R. Genome of a low-salinity ammonia-oxidizing archaeon determined by single-cell and metagenomic analysis. *PLoS ONE* **6**, 16626 (2011).
155. Walker, C. B. et al. *Nitrosopumilus maritimus* genome reveals unique mechanisms for nitrification and autotrophy in globally distributed marine crenarchaea. *Proc. Natl. Acad. Sci. USA* **107**, 8818–8823 (2010).
156. Bayer, B. et al. Metabolic versatility of the nitrite-oxidizing bacterium *Nitrospira marina* and its proteomic response to oxygen-limited conditions. *ISME J.* **15**, 1025–1039 (2021).
157. Amin, S. A. et al. Copper requirements of the ammonia-oxidizing archaeon *Nitrosopumilus maritimus* SCM1 and implications for nitrification in the marine environment. *Limnol. Oceanogr.* **58**, 2037–2045 (2013).
158. Sunda, W. G. & Lewis, J. A. M. Effect of complexation by natural organic ligands on toxicity of copper to a unicellular alga, *Monochrysis-lutheri*. *Limnol. Oceanogr.* **23**, 870–876 (1978).
159. Sunda, W. G. *The Relationship between Cupric Ion Activity and the Toxicity of Copper to Phytoplankton* (Massachusetts Institute of Technology, 1975).
160. Guo, J., Lapi, S., Ruth, T. J. & Maldonado, M. T. The effects of iron and copper availability on the copper stoichiometry of marine phytoplankton1. *J. Phycol.* **48**, 312–325 (2012).
161. Croot, P. L., Karlson, B., Van Elteren, J. T. & Kroon, J. J. Uptake and efflux of ⁶⁴Cu by the marine cyanobacterium *Synechococcus* (WH7803). *Limnol. Oceanogr.* **48**, 179–188 (2003).
162. Quigg, A., Reinfelder, J. R. & Fisher, N. S. Copper uptake kinetics in diverse marine phytoplankton. *Limnol. Oceanogr.* **51**, 893–899 (2006).
163. Guo, J. et al. Copper-uptake kinetics of coastal and oceanic diatoms. *J. Phycol.* **46**, 1218–1228 (2010).
164. Hudson, R. J. M. Which aqueous species control the rates of trace metal uptake by aquatic biota? Observations and predictions of non-equilibrium effects. *Sci. Total Environ.* **219**, 95–115 (1998).
165. Kong, L. & Price, N. M. A reduction-dependent copper uptake pathway in an oceanic diatom. *Limnol. Oceanogr.* **65**, 601–611 (2020).
166. Kong, L. & Price, N. M. Functional CTR-type Cu(I) transporters in an oceanic diatom. *Environ. Microbiol.* **21**, 98–110 (2019).
167. Semeniuk, D. M. et al. Plankton copper requirements and uptake in the subarctic Northeast Pacific Ocean. *Deep Sea Res. Part I Oceanogr. Res. Pap.* **56**, 1130–1142 (2009).
168. Semeniuk, D. M., Bundy, R. M., Payne, C. D., Barbeau, K. A. & Maldonado, M. T. Acquisition of organically complexed copper by marine phytoplankton and bacteria in the northeast subarctic Pacific Ocean. *Mar. Chem.* **173**, 222–233 (2015).
169. Shaked, Y., Buck, K. N., Mellett, T. & Maldonado, M. T. Insights into the bioavailability of oceanic dissolved Fe from phytoplankton uptake kinetics. *ISME J.* **14**, 1182–1193 (2020).
170. Lis, H., Shaked, Y., Kranzler, C., Keren, N. & Morel, F. M. M. Iron bioavailability to phytoplankton: an empirical approach. *ISME J.* **9**, 1003–1013 (2015).
171. Jones, G. J., Palenik, B. & Morel, F. M. M. Trace metal reduction by phytoplankton: the role of plasmalemma redox enzymes. *J. Phycol.* **23**, 237–244 (1987).
172. Walsh, M. J., Goodnow, S. D., Vezeau, G. E., Richter, L. V. & Ahner, B. A. Cysteine enhances bioavailability of copper to marine phytoplankton. *Environ. Sci. Technol.* **49**, 12145–12152 (2015).
173. Shank, G. C., Whitehead, R. F., Smith, M. L., Skrabal, S. A. & Kieber, R. J. Photodegradation of strong copper-complexing ligands in organic-rich estuarine waters. *Limnol. Oceanogr.* **51**, 884–892 (2006).
174. Moffett, J. W., Zika, R. G. & Brand, L. E. Distribution and potential sources and sinks of copper chelators in the Sargasso Sea. *Deep Sea Res. Part A. Oceanogr. Res. Pap.* **37**, 27–36 (1990).
175. Cottrell, B. A. et al. Photochemistry of marine and fresh waters: A role for copper-dissolved organic matter ligands. *Mar. Chem.* **162**, 77–88 (2014).
176. Sato, M., Ogata, N., Wong, K. H., Obata, H. & Takeda, S. Photodecomposition of natural organic metal-binding ligands from deep seawater. *Mar. Chem.* **230**, 103939 (2021).
177. Laglera, L. M. & van den Berg, C. M. G. Photochemical oxidation of thiols and copper complexing ligands in estuarine waters. *Mar. Chem.* **101**, 130–140 (2006).
178. Barbeau, K., Rue, E. L., Bruland, K. W. & Butler, A. Photochemical cycling of iron in the surface ocean mediated by microbial iron(III)-binding ligands. *Nature* **413**, 409–413 (2001).
179. Brooks, M. L., Meyer, J. S. & McKnight, D. M. Photooxidation of wetland and riverine dissolved organic matter: altered copper complexation and organic composition. *Hydrobiologia* **579**, 95–113 (2007).
180. Mellett, T. et al. The biogeochemical cycling of iron, copper, nickel, cadmium, manganese, cobalt, lead, and scandium in a California Current experimental study. *Limnol. Oceanogr.* **63**, S425–S447 (2018).
181. Mellett, T. & Buck, K. N. Spatial and temporal variability of trace metals (Fe, Cu, Mn, Zn, Co, Ni, Cd, Pb), iron and copper speciation, and electroactive Fe-binding humic substances in surface waters of the eastern Gulf of Mexico. *Mar. Chem.* **227**, 103891 (2020).
182. Little, S. H., Vance, D., Siddall, M. & Gasson, E. A modeling assessment of the role of reversible scavenging in controlling oceanic dissolved Cu and Zn distributions. *Global Biogeochem. Cycles* **27**, 780–791 (2013).
183. Ryan, B. M., Kirby, J. K., Degryse, F., Scheiderich, K. & McLaughlin, M. J. Copper isotope fractionation during equilibration with natural and synthetic ligands. *Environ. Sci. Technol.* **48**, 8620–8626 (2014).

184. Little, S. H. et al. Paired dissolved and particulate phase Cu isotope distributions in the South Atlantic. *Chem. Geol.* **502**, 29–43 (2018).
185. Whitby, H. et al. Contribution of electroactive humic substances to the iron-binding ligands released during microbial remineralization of sinking particles. *Geophys. Res. Lett.* **47**, e2019GL086685 (2020).
186. Gerringa, L. J. A., Rijkenberg, M. J. A., Schoemann, V., Laan, P. & de Baar, H. J. W. Organic complexation of iron in the West Atlantic Ocean. *Mar. Chem.* **177**, 434–446 (2015).
187. Roshan, S., DeVries, T. & Wu, J. Constraining the global ocean Cu cycle with a data-assimilated diagnostic model. *Global Biogeochem. Cycles* **34**, e2020GB006741 (2020).
188. Stüeken, E. E. Hydrothermal vents and organic ligands sustained the Precambrian copper budget. *Geochem. Perspect. Lett.* 12–16 <https://doi.org/10.7185/GEOCHEMLET.2037> (2020).
189. Di Bonito, M., Lofts, S. & Groenenberg, J. E. in *Environmental Geochemistry: Site Characterization, Data Analysis and Case Histories*. 2nd edn, 237–305 (Elsevier, 2018).
190. Gustafsson, J. P. Modeling the acid-base properties and metal complexation of humic substances with the Stockholm Humic Model. *J. Colloid Interface Sci.* **244**, 102–112 (2001).
191. Koopal, L. K., Saito, T., Pinheiro, J. P. & Van Riemsdijk, W. H. in *Colloids and Surfaces A: Physicochemical and Engineering Aspects*. Vol. 265, 40–54 (Elsevier B.V., 2005).
192. Koopal, L. K., Van Riemsdijk, W. H. & Kinniburgh, D. G. Humic matter and contaminants. General aspects and modeling metal ion binding. *Pure Appl. Chem.* **73**, 2005–2016 (2001).
193. Saito, T., Nagasaki, S., Tanaka, S. & Koopal, L. K. Application of the NICA-Donnan model for proton, copper and uranyl binding to humic acid. *Radiochim. Acta* **92**, 567–574 (2004).
194. Tipping, E., Lofts, C. S. & Sonke, J. E. Humic ion-binding model VII: a revised parameterisation of cation-binding by humic substances. *Environ. Chem.* **8**, 225–235 (2011).
195. De Schamphelaere, K. A. C. & Janssen, C. R. A biotic ligand model predicting acute copper toxicity for *Daphnia magna*: the effects of calcium, magnesium, sodium, potassium, and pH. *Environ. Sci. Technol.* **36**, 48–54 (2002).
196. Arnold, W. R. Effects of dissolved organic carbon on copper toxicity: implications for saltwater copper criteria. *Integr. Environ. Assess. Manag.* **1**, 34–39 (2005).
197. Niyogi, S. & Wood, C. M. Biotic ligand model, a flexible tool for developing site-specific water quality guidelines for metals. *Environ. Sci. Technol.* **38**, 6177–6192 (2004).
198. Playle, R. C. Using multiple metal-gill binding models and the toxic unit concept to help reconcile multiple-metal toxicity results. *Aquat. Toxicol.* **67**, 359–370 (2004).
199. De Schamphelaere, K. A. C. et al. Development and field validation of a predictive copper toxicity model for the green alga *Pseudokirchneriella subcapitata*. *Environ. Toxicol. Chem.* **22**, 2454–2465 (2003).
200. Zitoun, R. et al. Copper toxicity to blue mussel embryos (*Mytilus galloprovincialis*): the effect of natural dissolved organic matter on copper toxicity in estuarine waters. *Sci. Total Environ.* **653**, 300–314 (2019).
201. Hirose, K. Conditional stability constants of metal complexes of organic ligands in sea water: past and present, and a simple coordination chemistry model. *Anal. Chim. Acta* **284**, 621–634 (1994).
202. Hirose, K. Metal-organic matter interaction: ecological roles of ligands in oceanic DOM. *Appl. Geochem.* **22**, 1636–1645 (2007).
203. Hirose, K. Chemical modeling of marine trace metals: Effects of ocean acidification to marine ecosystem. In *2011 Seventh International Conference on Natural Computation*. **4**, 2023–2026 (2011).
204. Prosnier, L., Loreau, M. & Hulot, F. D. Modeling the direct and indirect effects of copper on phytoplankton-zooplankton interactions. *Aquat. Toxicol.* **162**, 73–81 (2015).
205. Hoffmann, S. R., Shafer, M. M. & Armstrong, D. E. Strong colloidal and dissolved organic ligands binding copper and zinc in rivers. *Environ. Sci. Technol.* **41**, 6996–7002 (2007).
206. Waelles, M., Tanguy, V., Lespès, G. & Riso, R. D. Behaviour of colloidal trace metals (Cu, Pb and Cd) in estuarine waters: an approach using frontal ultrafiltration (UF) and stripping chronopotentiometric methods (SCP). *Estuar. Coast. Shelf Sci.* **80**, 538–544 (2008).
207. Dai, M., Yin, Z., Meng, F., Liu, Q. & Cai, W. J. Spatial distribution of riverine DOC inputs to the ocean: an updated global synthesis. *Curr. Opin. Environ. Sustain.* **4**, 170–178 (2012).
208. Ludwig, W., Probst, J.-L. & Kempe, S. Predicting the oceanic input of organic carbon by continental erosion. *Global Biogeochem. Cycles* **10**, <https://doi.org/10.1029/95GB02925> (1996).
209. Aumont, O., Ethé, C., Tagliabue, A., Bopp, L. & Gehlen, M. PISCES-v2: an ocean biogeochemical model for carbon and ecosystem studies. *Geosci. Model Dev.* **8**, 2465–2513 (2015).
210. Santos-Echeandía, J., Caetano, M., Laglera, L. M. & Vale, C. Salt-marsh areas as copper complexing ligand sources to estuarine and coastal systems. *Chemosphere* **90**, 772–781 (2013).
211. Murray, N. J. et al. The global distribution and trajectory of tidal flats. *Nature* <https://doi.org/10.1038/s41586-018-0805-8> (2019).
212. Karavoltzos, S. et al. Copper complexation in wet precipitation: Impact of different ligand sources. *Atmos. Environ.* **80**, 13–19 (2013).
213. Hou, H., Takamatsu, T., Koshikawa, M. K. & Hosomi, M. Copper complexing capacity of throughfall and its environmental effect. *Water, Air, Soil Pollut.* **162**, 229–245 (2005).
214. Witt, M. L. I., Skrabal, S., Kieber, R. & Willey, J. Photochemistry of Cu complexed with chromophoric dissolved organic matter: implications for Cu speciation in rainwater. *J. Atmos. Chem.* **58**, 89–109 (2007).
215. Kanakidou, M. et al. Atmospheric fluxes of organic N and P to the global ocean. *Global Biogeochem. Cycles* **26**, 3026 (2012).
216. Abernathy, R. P. et al. Water-mass transformation by sea ice in the upper branch of the Southern Ocean overturning. *Nat. Geosci.* **9**, 596–601 (2016).
217. Echeveste, P., Croot, P. & von Dassow, P. Differences in the sensitivity to Cu and ligand production of coastal vs offshore strains of *Emiliania huxleyi*. *Sci. Total Environ.* **625**, 1673–1680 (2018).
218. Tang, D., Shafer, M. M., Karner, D. A. & Armstrong, D. E. Response of nonprotein thiols to copper stress and extracellular release of glutathione in the diatom *Thalassiosira weissflogii*. *Limnol. Oceanogr.* **50**, 516–525 (2005).
219. Zahn, J. A. & DiSpirito, A. A. Membrane-associated methane monooxygenase from *Methylococcus capsulatus* (Bath). *J. Bacteriol.* **178**, 1018–1029 (1996).
220. Gledhill, M., Nimmo, M., Hill, S. J. & Brown, M. T. The release of copper-complexing ligands by the brown alga *Fucus vesiculosus* (Phaeophyceae) in response to increasing total copper levels. *J. Phycol.* **35**, 501–509 (1999).
221. Kawakami, S. K., Gledhill, M. & Achterberg, E. P. Effects of metal combinations on the production of phytochelatin and glutathione by the marine diatom *Phaeodactylum tricornutum*. *Biomol. Biotechnol.* **19**, 51–60 (2006).
222. Srna, R. F., Garrett, K. S., Miller, S. M. & Thum, A. B. Copper complexation capacity of marine water samples from southern California. *Environ. Sci. Technol.* **14**, 1482–1486 (1980).
223. Nilsen, S. K. & Lund, W. The determination of weakly and strongly bound copper, lead and cadmium in Oslofjord samples. *Mar. Chem.* **11**, 223–233 (1982).
224. Donat, J. R., Statham, P. J. & Bruland, K. W. An evaluation of a C-18 solid phase extraction technique for isolating metal-organic complexes from central North Pacific Ocean waters. *Mar. Chem.* **18**, 85–99 (1986).
225. Li, L., Liu, J., Wang, X. & Shi, X. Dissolved trace metal distributions and Cu speciation in the southern Bohai Sea, China. *Mar. Chem.* **172**, 34–45 (2015).
226. Wong, K. H., Obata, H., Ikhsani, I. Y. & Muhammad, R. Controls on the distributions of dissolved Cd, Cu, Zn, and Cu-binding organic ligands in the East China Sea. *J. Geophys. Res. Ocean.* **126**, e2020JC016997 (2021).
227. Wiwit et al. Wide-range detection of Cu-binding organic ligands in seawater using reverse titration. *Mar. Chem.* **230**, 103927 (2021).
228. Duinker, J. C. & Kramer, C. J. M. An experimental study on the speciation of dissolved zinc, cadmium, lead and copper in River Rhine and North Sea water, by differential pulsed anodic stripping voltammetry. *Mar. Chem.* **5**, 207–228 (1977).
229. Hasle, J. R. & Abdullah, M. I. Analytical fractionation of dissolved copper, lead and cadmium in coastal seawater. *Mar. Chem.* **10**, 487–503 (1981).
230. Piotrowicz, S. R., Springer-Young, M., Puig, J. A. & Spencer, M. J. Anodic stripping voltammetry for evaluation of organic-metal interactions in seawater. *Anal. Chem.* **54**, 1367–1371 (1982).
231. Buckley, P. J. M. & Van Den Berg, C. M. G. Copper complexation profiles in the Atlantic-Ocean—a comparative study using electrochemical and ion-exchange techniques. *Mar. Chem.* **19**, 281–296 (1986).
232. Kramer, C. J. M. Apparent copper complexation capacity and conditional stability constants in North Atlantic waters. *Mar. Chem.* **18**, 335–349 (1986).
233. Donat, J. R. & van den Berg, C. M. G. A new cathodic stripping voltammetric method for determining organic copper complexation in seawater. *Mar. Chem.* **38**, 69–90 (1992).
234. Moffett, J. W. Temporal and spatial variability of copper complexation by strong chelators in the Sargasso Sea. *Deep. Res. Part I Oceanogr. Res. Pap.* **42**, 1273–1295 (1995).
235. Kozelka, P. B. & Bruland, K. W. Chemical speciation of dissolved Cu, Zn, Cd, Pb in Narragansett Bay, Rhode Island. *Mar. Chem.* **60**, 267–282 (1998).
236. Zitoun, R. et al. The complex provenance of Cu-binding ligands in the South-East Atlantic. *Mar. Chem.* **237**, 104047 (2021).
237. Donat, J. R., Lao, K. A. & Bruland, K. W. Speciation of dissolved copper and nickel in South San-Francisco Bay—a multimethod approach. *Anal. Chim. Acta* **284**, 547–571 (1994).
238. Wells, M. L., Kozelka, P. B. & Bruland, K. W. The complexation of dissolved Cu, Zn, Cd and Pb by soluble and colloidal organic matter in Narragansett Bay, RI. *Mar. Chem.* **62**, 203–217 (1998).

239. Dryden, C. L., Gordon, A. S. & Donat, J. R. Interactive regulation of dissolved copper toxicity by an estuarine microbial community. *Limnol. Oceanogr.* **49**, 1115–1122 (2004).
240. Jacquot, J. E. et al. Assessment of the potential for copper limitation of ammonia oxidation by Archaea in a dynamic estuary. *Mar. Chem.* **162**, 37–49 (2014).
241. Wong, K. H. et al. Organic complexation of copper in estuarine waters: an assessment of the multi-detection window approach. *Mar. Chem.* **204**, 144–151 (2018).
242. Batley, G. E. & Florence, T. M. Determination of the chemical forms of dissolved cadmium, lead and copper in seawater. *Mar. Chem.* **4**, 347–363 (1976).
243. Sukekava, C., Downes, J., Slagter, H. A., Gerringa, L. J. A. & Laglera, L. M. Determination of the contribution of humic substances to iron complexation in seawater by catalytic cathodic stripping voltammetry. *Talanta* **189**, 359–364 (2018).
244. Laglera, L. M. & Van Den Berg, C. M. G. Evidence for geochemical control of iron by humic substances in seawater. *Limnol. Oceanogr.* **54**, 610–619 (2009).
245. Longhini, C. M. et al. Coastal waters contamination by mining tailings: what triggers the stability of iron in the dissolved and soluble fractions? *Limnol. Oceanogr.* **66**, 171–187 (2021).
246. Dulaquais, G. et al. The biogeochemistry of electroactive humic substances and its connection to iron chemistry in the North East Atlantic and the Western Mediterranean Sea. *J. Geophys. Res. Ocean.* **123**, 5481–5499 (2018).
247. Gourain, C. G. *Copper Biogeochemical Cycle and the Organic Complexation of Dissolved Copper in the North Atlantic*. The University of Liverpool (United Kingdom) ProQuest Dissertations Publishing (2020).
248. Karavoltos, S. Organic complexation of copper in throughfall and open field bulk deposition: Influence of the tree canopy of Mediterranean forest ecosystems. *Chemosphere* **167**, 28–35 (2017).

Acknowledgements

We would like to thank Stan van den Berg, Kuo Hong Wong, Sylvia Sander and one anonymous reviewers for their insightful comments on the manuscript. A.R. was funded by the National Science Foundation Polar Programs Postdoctoral Fellowship #2138217, R.M.B. by a Life Sciences Simons Early Career Investigator in Marine Microbial Ecology Award #618401, and C.R. was supported by ISblue project, Interdisciplinary graduate school for the blue planet (ANR-17-EURE-0015) and co-funded by a grant from the French government under the program “Investissements d’Avenir”, and by a grant from the Regional Council of Brittany (SAD program).

Author contributions

A.R. led the writing of the manuscript and generated figures, C.R., H.W., and R.B. helped write and edit the manuscript and contributed to making the figures and tables.

Competing interests

The authors declare no competing interests.

Additional information

Supplementary information The online version contains supplementary material available at <https://doi.org/10.1038/s43247-022-00597-1>.

Correspondence and requests for materials should be addressed to Angel Ruacho or Randelle M. Bundy.

Peer review information *Communications Earth & Environment* thanks Kuo Hong Wong, Sylvia Sander and Susan Little for their contribution to the peer review of this work. Primary handling editors: Joe Aslin, Heike Langenberg. Peer reviewer reports are available.

Reprints and permission information is available at <http://www.nature.com/reprints>

Publisher’s note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The images or other third party material in this article are included in the article’s Creative Commons license, unless indicated otherwise in a credit line to the material. If material is not included in the article’s Creative Commons license and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>.

© The Author(s) 2022