



**HAL**  
open science

## Ionospheric composition of comet 67P near perihelion with multi-instrument Rosetta datasets

Zoe Lewis, Arnaud Beth, Kathrin Altwegg, Anders Eriksson, Marina Galand, Charlotte Götz, P. Henri, Kevin Héritier, Laurence O'Rourke, Ingo Richter, et al.

### ► To cite this version:

Zoe Lewis, Arnaud Beth, Kathrin Altwegg, Anders Eriksson, Marina Galand, et al.. Ionospheric composition of comet 67P near perihelion with multi-instrument Rosetta datasets. 16th Europlanet Science Congress 2022, 0000, à renseigner, Unknown Region. 10.5194/epsc2022-960 . insu-04089894

**HAL Id: insu-04089894**

**<https://hal-insu.archives-ouvertes.fr/insu-04089894>**

Submitted on 5 May 2023

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution| 4.0 International License



## Ionospheric composition of comet 67P near perihelion with multi-instrument Rosetta datasets

Zoe Lewis<sup>1</sup>, Arnaud Beth<sup>2</sup>, Kathrin Altwegg<sup>3</sup>, Anders Eriksson<sup>4</sup>, Marina Galand<sup>1</sup>, Charlotte Götz<sup>5</sup>, Pierre Henri<sup>6,7</sup>, Kevin Héritier<sup>1</sup>, Laurence O'Rourke<sup>8</sup>, Ingo Richter<sup>9</sup>, Martin Rubin<sup>3</sup>, Peter Stephenson<sup>1</sup>, and Xavier Vallières<sup>7</sup>

<sup>1</sup>Department of Physics, Imperial College London, London, UK

<sup>2</sup>Department of Physics, Umeå University, Umeå, Sweden

<sup>3</sup>Physikalisches Institut, University of Bern, Bern, Switzerland

<sup>4</sup>Swedish Institute of Space Physics, Uppsala, Sweden

<sup>5</sup>ESTEC, European Space Agency, Noordwijk, Netherlands

<sup>6</sup>LPC2E, CNRS, Université d'Orléans, Orléans, France

<sup>7</sup>Laboratoire Lagrange, OCA, UCA, CNRS, Nice, France

<sup>8</sup>ESAC, European Space Agency, Madrid, Spain

<sup>9</sup>Institut für Geophysik und extraterrestrische Physik, TU Braunschweig, Braunschweig, Germany

The European Space Agency Rosetta mission escorted comet 67P/Churyumov-Gerasimenko for two years, during which it acquired an extensive dataset, revealing unprecedented detail about the neutral and plasma environment of the coma. The measurements were made over a large range of heliocentric distances, and therefore of outgassing activities, as Rosetta witnessed 67P evolve from a low-activity icy body at 3.8 AU to a dynamic object with large-scale plasma structures and rich ion and neutral chemistry near perihelion at 1.2 AU. One such plasma structure is the diamagnetic cavity, a region of negligible magnetic field surrounding the comet nucleus. It is formed through the interaction of the unmagnetized outwardly expanding cometary plasma with the incoming solar wind. This region was encountered many times by Rosetta between April 2015 and February 2016, as the comet moved towards and away from perihelion.

In this study, we focus on the changing role of chemistry during the escort phase, particularly on trends in the detection of high proton affinity species near perihelion and within the diamagnetic cavity.  $\text{NH}_4^+$  is produced through the protonation of  $\text{NH}_3$  which has the highest proton affinity of the neutral species and is therefore the terminal ion. The ratio of this species to the major ion species  $\text{H}_3\text{O}^+$  can then be an indicator of the importance of ion-neutral chemistry as an ion loss process compared to transport. We use data from the high resolution mode of the ROSINA (Rosetta Orbital Spectrometer for Ion-Neutral Analysis)/DFMS (Double Focussing Mass Spectrometer) instrument, which allows certain ions of the same integer mass per charge ratio to be separated from one another, most importantly  $\text{H}_2\text{O}^+$  and  $\text{NH}_4^+$ . This data is then analysed alongside a range of plasma properties from the RPC (Rosetta Plasma Consortium) suite of instruments, to determine features of the plasma within and outside of the diamagnetic cavity that may systematically impact the ion species detected in the mass spectrometer, for example through the spacecraft potential.

For the same period, ion ROSINA/DFMS ion data are analysed together with simultaneous

observations of the cometary plasma properties by RPC instruments, in order to identify systematic plasma features or biases which may impact the ion composition detected by ROSINA. In addition, the ion composition measurements are compared to ionospheric simulations. We evaluate the relative significance of ion-neutral chemistry and transport such that their impacts on the ion composition in different interaction regions are identified.